Comments on the proposed conservation of usage of the names *Mystacina* Gray, 1843, *Chalinolobus* Peters, 1866, *M. tuberculata* Gray, 1843 and *C. tuberculatus* (J.R. Forster, 1844) (Mammalia, Chiroptera)
(Case 3095; see BZN 56: 250–254)

(1) Martyn Kennedy

Division of Environmental and Evolutionary Biology, Institute of Biomedical and Life Sciences, University of Glasgow, Glasgow G12 8QQ, U.K.

I support Spencer & Lee's application for the conservation of name usage for the New Zealand bats *Mystacina tuberculata* Gray, 1843 and *Chalinolobus tuberculatus* (J.R. Forster, 1844); their argument is compelling. These names have been universally accepted for a century or more (in addition to references cited in the application see Miller (1907), Pierson et al. (1986), Koopman (1994), Hand et al. (1998) and Kennedy et al. (1999)). The name *M. velutina* Hutton, 1872 has been used instead of *M. tuberculata* only by Thomas (1905; as *Mystacops velutinus*) and by Mayer et al. (1999), in both cases on the mistaken grounds described in the application. Because New Zealand has only two known extant bat species they are commonly known by their vernacular and generic names, and the similarity of their specific names has not in fact caused confusion. The stable usage of *Mystacina tuberculata* should continue.

Additional references

- Hand, S.J., Murray, P., Megirian, D., Archer, M. & Godthelp, H. 1998. Mystacinid bats (Microchiroptera) from the Australian Tertiary. *Journal of Paleontology*, **72**: 538–545.
- Kennedy, M., Paterson, A.M., Morales, J.C., Parsons, S., Winnington, A.P. & Spencer, H.G. 1999. The long and short of it: Branch lengths and the problem of placing the New Zealand short-tailed bat, *Mystacina*. *Molecular Phylogenetics and Evolution*, **13**: 405–416.
- Koopman, K.F. 1994. Chiroptera: Systematics. *Handbook of Zoology*, vol. 8, pt. 60.
- Miller, G.S. 1907. The families and genera of bats. *Bulletin of the United States National Museum*, no. 57. 282 pp.
- Pierson, E.D., Sarich, V.M., Lowenstein, J.M., Daniel, M.J. & Rainey, W.E. 1986. A molecular link between the bats of New Zealand and South America. *Nature*, **323**: 60–63.

(2) Kerry-Jayne Wilson

Ecology and Entomology Group, P.O. Box 84, Lincoln University, Canterbury, New Zealand

I lecture in vertebrate ecology at Lincoln University and have had three graduate students do theses on New Zealand bats. I have frequent contact with government agencies and, by means of broadcasts and written articles, with the lay public on matters concerning the ecology and conservation of New Zealand's native biota, including the bat species. I know of nobody who finds the existing scientific names of the bats confusing, and I urge their retention.

(3) Trevor Worthy

Palaeofaunal Surveys, 43 The Ridgeway, Nelson, New Zealand

I would like to go on record as supporting the well-founded arguments and proposals of Spencer & Lee. There is no doubt as to what taxa the names *Mystacina*

Kennedy, M et al. 2000. "On the proposed conservation of usage of the names *Mystacina* Gray, 1843, *Chalinolobus* Peters, 1866, *M. tuberculata* Gray, 1843 and *C. tuberculatus* (JR Forster, 1844) (Mammalia, Chiroptera)." *The Bulletin of zoological nomenclature* 57, 117–117. <https://doi.org/10.5962/bhl.part.20697>.

View This Item Online: <https://www.biodiversitylibrary.org/item/45022>

DOI: <https://doi.org/10.5962/bhl.part.20697>

Permalink: <https://www.biodiversitylibrary.org/partpdf/20697>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.