

- Freund, R.J. and R.C. Littell. 1986. SAS system for regression. SAS Institute Inc., Cary, N.C. 164 pp.
- Gray, T.G.; Shepherd, R.F.; Wood, C.S. 1973. A hot-water technique to remove insect eggs from foliage. Can. For. Serv., Bi-mon. Res. Notes 29(5):29.
- Harris, J.W.E., A.F. Dawson, and R.G. Brown. 1982. The western hemlock looper in British Columbia 1911–1980. Environ. Can., Can. For. Serv., Pac. For. Cent., Inf. Rept. BC-X-234, 18 pp.
- Hopping, G.R. 1934. An account of the western hemlock looper, *Ellopiia somniaria* Hulst, on conifers in British Columbia. Scientific Agriculture XV:12–27.
- Kinghorn, J.M. 1952. Western hemlock looper egg sampling. Can. Dep. Agric., Bi-mon. Prog. Rept. 8(3):3–4.
- Kinghorn, J.M. 1954. The effect of stand composition on the mortality of various conifers, caused by defoliation by the western hemlock looper on Vancouver Island, British Columbia. For. Chron. 30(4):380–400.
- Otvos, I.S. and D.G. Bryant. 1972. An extraction method for rapid sampling of eastern hemlock looper eggs, *Lambdina fiscellaria fiscellaria* (Lepidoptera: Geometridae). Can. Ent. 104:1511–1514.
- Richmond, H.A. 1947. Current trend of the western hemlock looper (*Lambdina f. lugubrosa*) in the coastal forest of British Columbia (Lepidoptera: Geometridae). Entomol. Soc. of British Columbia, Proc. (1946) 43:33–35.
- SAS Institute Inc., 1985. SAS users guide: Statistics, Version 5 Edition. Cary, NC: SAS Institute Inc., 956 pp.
- Shepherd, R.F. and T.G. Gray. 1972. Solution separation and maximum likelihood density estimates of hemlock looper (Lepidoptera: Geometridae) eggs in moss. Can. Ent. 104:751–754.
- Shore, T.L. 1989. Sampling western hemlock looper pupae (Lepidoptera: Geometridae) using burlap traps. J. Entomol. Sci. 24(3):348–354.
- Thomson, M.G. 1949. Determination of larval populations of forest defoliators by frass fall and body weight, with particular reference to the western hemlock looper. Can. Dep. Agric., Sci. Serv., Div. For. Biol., For. Biol. Lab. Victoria, B.C. 49 pp.
- Thomson, M.G. 1957. Appraisal of western hemlock looper infestations. For. Chron. 33(2):141–147.
- Thomson, M.G. 1958. Egg sampling for the western hemlock looper. For. Chron. 34(3):248–256.
- Wyatt, G.R. 1946. Hemlock looper: history of outbreaks on the Pacific coast. Can. Dep. Agric., Science Serv., Div. For. Biol., For. Biol. Lab., Victoria, B.C., Interim Tech. Rep. 8 pp.

A review of mosquito collecting in the Yukon

E.M. BELTON and P. BELTON
 CENTRE FOR PEST MANAGEMENT
 DEPARTMENT OF BIOLOGICAL SCIENCES
 SIMON FRASER UNIVERSITY
 BURNABY, B.C. V5A 1S6

The first formal record of a mosquito collected in the Yukon was in 1904 when J. Keele caught *Anopheles occidentalis* in the Mayo River valley (Dyar 1921). In 1916 three females of *Aedes nearcticus* were collected on Herschel Island, off the north coast of the Yukon, by Frits Johansen of the Canadian Arctic Expedition (Dyar 1919). These species, now known as *An. earlei* and *Ae. impiger* respectively, were identified at the time by Dr. Harrison Dyar at the United States National Museum in Washington.

Dyar, himself, visited the Yukon in June and July of 1919. He travelled from Carcross in the south, along the Yukon valley to Dawson which is less than half way to Herschel Island. He recorded 16 species (Dyar 1920, 1921) including nearly 2,000 specimens of *Ae. cataphylla* which he found to be the dominant species with *Ae. campestris*, *Ae. communis* and *Ae. punctator* also common (Table 1). He described three new species from his Yukon material: *Ae. nearcticus* from Herschel Island, the Northwest Territories and Alaska (Dyar 1919); *Ae. callithotrys* from Whitehorse and Takheena River in the Yukon and from Alaska; and *Ae. mercurator* from 65 specimens collected around Dawson (Dyar 1920). He later synonymised *Ae. callithotrys* with *Ae. campestris* (Dyar 1928).

By the 1920s Whitehorse was attracting tourists and, in 1926, presumably as a result of their complaints, Eric Hearle, who had already reduced the mosquito nuisance in the Lower Fraser Valley of B.C. and the resorts of Banff and Lake Louise in Alberta, was invited to make recommendations on controlling mosquitoes in the Yukon (Hearle 1927). He probably collected adults and larvae when he assessed the problem around Whitehorse.

The Dominion Entomologist, Arthur Gibson, reported on mosquito control in Canada from 1923–1941 to the Annual Meetings of the New Jersey Mosquito Control Association, but did not refer to any work in the Yukon.

From 1947 to 1950, mosquitoes were collected from several arctic and subarctic localities as part of the Northern Insect Survey, a joint endeavour of the Canada Department of Agriculture and the Department of National Defense. An Interim Report by Freeman (1952) of the mosquitoes obtained during the Survey consisted mainly of distribution maps. The 16 species reared from sites in the Yukon were mainly from Dawson or Whitehorse. Vockeroth (1954a) addressed the difficult problem of identifying the females and discussed their distribution. *Ae. nigripes* was not found in the Yukon during the Survey but he thought it was probably present because it is the most abundant species elsewhere in the arctic. Vockeroth (1954b) examined the type specimens of several arctic species. He pointed out that mosquitoes identified up to that time as *Ae. nearcticus* were in fact *Ae. impiger* and that the specimens from Dawson and elsewhere, identified in the Canadian National Collection as *Ae. impiger*, belonged to a new species which he described and named *Ae. implicatus*.

In his guide to the mosquito larvae of Western Canada, Rempel (1950) noted that 8 to 10 of the species in the guide occurred in the Yukon. He did not refer to collecting there himself but he may have seen representative specimens in collections loaned to him from the Canadian National Collection in Ottawa and the U.S. National Museum in Washington.

In the summers of 1949 and 1950, Colin Curtis, an entomologist from the Veterinary and Medical Entomology Laboratory at Kamloops, B.C., collected 21 species around Whitehorse and Watson Lake (Table 1). He noted that although *Ae. cataphylla* was common, the predominant pest mosquitoes were *Ae. communis* followed by *Ae. punctor* and *Ae. pionips* (Curtis 1953).

Dr. D.M. McLean, a medical microbiologist from the University of British Columbia, collected mosquitoes in northern B.C. and the Yukon during several seasons in a survey for mosquito-borne encephalitis viruses. He collected mainly adults and some larvae at about a dozen locations in the boreal forest region from Marsh Lake near Whitehorse in the southeast to an area near the Dempster Highway at 67°N and 137°W. He found seven species infected with viruses: *Ae. canadensis*, *Ae. cinereus*, *Ae. communis*, *Ae. hexodontus*, *Ae. nigripes*, *Ae. punctor*, and *Cs. inornata* (McLean, Judd & Shives 1981; McLean & Lester 1984).

One of the largest collections of mosquitoes from the Yukon was made in 1972 and 1973 by John Nelson, a Master of Pest Management student at Simon Fraser University, Burnaby, B.C. He set up New Jersey light traps and bite sampling stations at 28 sites from Watson Lake in the south to Old Crow within the Arctic Circle (Nelson 1977). Of about 27,000 specimens caught by him in 1972, the commonest biting species were *Ae. pionips*, *Ae. hexodontus*, *Ae. cataphylla*, *Ae. communis*, *Ae. campestris*, and *Ae. nigripes*, in that order, although it probably varied considerably from place to place. In addition to many of the species found by Dyar and Curtis, he listed nine more, five of them verified by Wood, Dang & Ellis (1979) and Wood (1989, personal communication), and four others, three of which are probably correct (Table 1), bringing the number of mosquito species in the Territory to about 30.

REFERENCES

Curtis, L.C. 1953. Observations on mosquitoes at Whitehorse, Yukon Territory. *Diptera: Culicidae*. *Can. Ent.* 85:353-370.

Dyar, H.G. 1919. The mosquitoes collected by the Canadian Arctic Expedition, 1913-18. *Rept. Can. Arctic Expedn. III*, pt C:31-33.

Dyar, H.G. 1920. The mosquitoes of British Columbia and the Yukon Territory, Canada. *Insecutor Inscit.* *menstr.* 8:1-27.

Dyar, H.G. 1921. The mosquitoes of Canada. *Trans. R. Can. Inst.* 13:71-120.

Dyar, H.G. 1928. The mosquitoes of the Americas. *Carnegie Inst. Wash. Publ.* 387:1-616.

Freeman, T.N. 1952. Interim Report of the distribution of the mosquitoes obtained in the Northern Insect Survey. Defense Research Board of Ottawa. *Tech. Rep.* 1. 2pp. 43 maps.

Hearle, E. 1927. Mosquito control activities in Western Canada. *Ann. Rep. Ent. Soc. Ont.* 58:45-50.

McLean, D.M., B.D. Judd & S.K.A. Shives. 1981. Snowshoe hare virus infections in Canadian Arctic mosquitoes during 1980. *Mosq. News* 41:287-290.

McLean, D.M. & S.A. Lester. 1984. Isolations of Snowshoe hare virus from the Yukon mosquitoes, 1983. *Mosq. News* 44:200-203.

Nelson, J. 1977. Mosquito control in the Yukon Territory, Canada. MPM Thesis, Simon Fraser Univ., Can. pp. 111.

Rempel, J.G. 1950. A guide to the mosquito larvae of western Canada. *Can. J. Res. D*, 28:207-248.

Vockeroth, J.R. 1954a. Notes on the identities and distribution of *Aedes* species of northern Canada, with a key to the females (*Diptera, Culicidae*). *Can. Ent.* 86:241-255.

Vockeroth, J.R. 1954b. Notes on northern species of *Aedes*, with descriptions of two new species (*Diptera, Culicidae*). *Can. Ent.* 86:109-116.

Wood, D.M., P.T. Dang & R.A. Ellis. 1979. The Mosquitoes of Canada. *Diptera: Culicidae*. *The Insects and Arachnids of Canada. Part 6. Agric. Can. Publ.* 1686. pp. 390.

Table 1
Mosquitoes recorded from the Yukon Territory

Dyar's species	<i>Cs. incidens</i>
<i>Anopheles earlei (occidentalis*)</i>	<i>Cs. morsitans (dyari*)</i>
<i>Aedes campestris (callithotrys*)</i>	Curtis's additional species
<i>Ae. cataphylla</i>	<i>Ae. canadensis</i>
<i>Ae. cinereus</i>	<i>Ae. dianiaetus</i>
<i>Ae. communis</i>	<i>Ae. flavescens</i>
<i>Ae. excrucians</i>	<i>Ae. hexodontus</i>
<i>Ae. fitchii</i>	<i>Ae. nigripes</i>
<i>Ae. impiger (nearcticus*)</i>	<i>Ae. riparius</i>
<i>Ae. mercurator</i>	<i>Culex territans (apicalis*)</i>
<i>Ae. pionips</i>	Nelson's additional species
<i>Ae. pullatus</i>	<i>Ae. decticus</i>
<i>Ae. punctor</i>	<i>Ae. intrudens</i>
<i>Ae. stimulans</i> (Dyar's reference to it in the Yukon should probably be to <i>Ae. mercurator</i> Wood, Dang & Ellis 1979)	<i>Ae. implicatus</i>
<i>Culiseta alaskaensis</i>	<i>Ae. vexans</i>
<i>Cs. impatiens</i>	<i>Cs. inornata</i>

The following 4 of Nelson's species were not recorded as occurring in the Yukon by Wood, Dang & Ellis 1979 and Wood, 1989, Personal Communication. The last three probably do occur there, but *Ae. sticticus* has not been found north of Terrace, B.C., in Western Canada.

<i>Ae. sticticus</i>	<i>Ae. provocans</i>
<i>Ae. euedes</i>	<i>Cx. tarsalis</i>

* Names in parentheses were used in the publications referred to in the text.



Belton, E. M. and Belton, Peter. 1990. "A review of mosquito collecting in the Yukon." *Journal of the Entomological Society of British Columbia* 87, 35–37.

View This Item Online: <https://www.biodiversitylibrary.org/item/181002>

Permalink: <https://www.biodiversitylibrary.org/partpdf/213880>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: Entomological Society of British Columbia

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.