Transfer of Four Species of *Scabiosa* to *Lomelosia* (Dipsacaceae)

Paolo Caputo* and Emanuele Del Guacchio

Dipartimento delle Scienze Biologiche, sez. Biologia Vegetale,
Università degli Studi di Napoli Federico II, via Foria, 223, I-80139 Napoli, Italy.

*Author for correspondence: pacaputo@unina.it

ABSTRACT. Four species of *Scabiosa* L. are transferred to *Lomelosia* Raf. (Dipsacaceae) to complete the relocation of all scabious taxa with a pitted epicalyx into *Lomelosia*. The new combinations are: *L. deserticola* (Rech. f.) P. Caputo & Del Guacchio, *L. poecilocarpa* (Rech. f.) P. Caputo & Del Guacchio, *L. schimperiana* (Boiss. & Buhse) P. Caputo & Del Guacchio, and *L. transcaspica* (Rech. f.) P. Caputo & Del Guacchio.

Key words: Dipsacaceae, *Lomelosia*, *Scabiosa*.

Scabiosa L., according to its original concept (Linnaeus, 1753), was a broadly comprehensive genus. It included all taxonomic entities later placed in Dipsacaceae by Jussieu (1789) with flat or round, many-flowered capitula. The other two dipsacaceous genera recognized by Linnaeus, i.e., *Dipsacus* L. and *Knautia* L., respectively, included the more or less prickly entities with cylindrical capitula and one single pauci-flowered species (*K. orientalis* L.), which resembled *Lychnis* L. in its gross floral morphology. Later, *Scabiosa* sensu Linnaeus underwent various vicissitudes, mainly restricting its circumscription; taxa were segregated into different genera, and *Scabiosa* was subdivided into infrageneric taxa. Some of these names contributed to distinguish *Scabiosa* from far removed forms. For example, Adanson (1763) segregated the taxa with multiplied, plumose calyx awns into *Pterocephalus* Adans.; Haller (1768) segregated *Succisa* Haller, which has a praemorse rhizome and a 5-lobed epicalyx, while Roemer and Schultes, in their *Systema Vegetabilium* (1818), distinguished *Cephalaria* Schrad. ex Roem. & Schult. for its multiseriate involucre bracts and awned epicalyx. Additionally, Coulter (1824) gave a much wider circumscription for *Knautia*, and Beck (1893) separated *Succisella* Beck mainly on the basis of its urceolate epicalyx and lack of calyx awns.

Around the same time, various scholars, such as Coulter (1824) and Höck (1891), presented a much more restricted circumscription of the genus *Scabiosa* (including only taxa with an epicalyx distally expanded into a more or less membranous rim or corona), segregating *Trochocephalus* (Mert. & W. D. J. Koch) Opiz (= *Scabiosa* sect. *Trochocephalus* Mert.

& W. D. J. Koch, = subgenus *Asterocephalus* (Coulter.) Lack & Rech. f.), *Cyrtostemma* (Mert. & W. D. J. Koch) Spach (= *Scabiosa* sect. *Cyrtostemma* Mert. & W. D. J. Koch), *Sclerostemma* Schott ex Roem. & Schult. (= *Scabiosa* sect. *Sclerostemma* Mert. & W. D. J. Koch), and *Spongostemma* (Rehdb.) Rehdb. (= *Scabiosa* subg. *Spongostemma* Rehdb.). By the 20th century, the taxonomic concept of *Scabiosa* included only those forms with more or less radiant capitula, a 5-awned calyx, and an epicalyx with a membranous corona (Ehrendorfer, 1964; Moore, 1976). *Scabiosa* sect. *Trochocephalus* includes taxa with eight pits at the apex of the epicalyx tube; *Scabiosa* sect. *Cyrtostemma* includes taxa with what were interpreted at the time as eight membranous openings (fenestrae) in an otherwise sclerified corona (Mayer and Ehrendorfer [1999] demonstrated that fenestrae are part of the epicalyx tube, not of the corona); and *Scabiosa* sect. *Scabiosa* includes taxa without pits or fenestrae.

After the studies of Verlaque (1984, 1985, 1986a, 1986b) and Devesa (1984), which demonstrated independent evolutionary histories for the various sections of *Scabiosa*, Greuter and Raus (1985) designated *Lomelosia* Raf. (with priority over *Trochocephalus*) for the species traditionally belonging to *Scabiosa* sect. *Trochocephalus* and selected *Sixalix* Raf. for those species belonging to *Scabiosa* sect. *Cyrtostemma*. However, their work concerned only circum-mediterranean species.

The generic name *Pseudoscabiosa* Devesa (a replacement name for *Scabiosa* sect. *Asterothrix* Font Quer) was subsequently used (Greuter et al., 1986) for four archaic species traditionally either placed within *Scabiosa* sect. *Trochocephalus* or isolated in section *Asterothrix*. One of these species was later segregated by López González (1987) as a new genus *Pterocephalidium* G. López as *P. diandrum* (Lag.) G. López. Later, Soják (1987) transferred several *Scabiosa* taxa from Asia to *Lomelosia*. Finally, Rechinger (1989) and Lack and Rechinger (1991), while describing new species from Iran and adjacent Southwest Asia, did not adopt the nomenclatural changes proposed in Greuter and Raus (1985), and raised *Scabiosa* sect. *Asterocephalus* to subgeneric

status. Moreover, Rechinger (1989) created *Scabiosa* sect. *Olivierianae* Rech. f. to accommodate some annual members of *Scabiosa* subg. *Asterocephalus* with few-flowered capitula (this section was confirmed as monophyletic in a morphological investigation on species of *Lomelosia* by Castro & Caputo, 1999). Finally, on the basis of fruit differentiation and pollen, Mayer and Ehrendorfer (1999) demonstrated quite convincingly that *Lomelosia* is rather distant from the remaining members of *Scabiosa*, whereas *Scabiosa* s. str. and *Sixalix* are sister groups and so close that they hardly deserve separate generic status. Therefore, they proposed to merge *Sixalix* again within *Scabiosa* sect. *Cyrtostemma*. The distance of *Lomelosia* from the other scabious Dipsacaceae, as well as the sister group relationship between *Sixalix* and *Scabiosa*, were confirmed on molecular grounds by Caputo et al. (2004) and Avino et al. (2009), based on four DNA regions from the chloroplast and nucleus. Therefore, *Scabiosa* now encompasses only the eight species located in *Scabiosa* sect. *Cyrtostemma* and ca. 30 in section *Scabiosa* (= *Scabiosa* sect. *Sclerostemma*).

Accordingly, the transfer to *Lomelosia* is here proposed for three scabious taxa described by Rechinger (1989) and Lack and Rechinger (1991) in *Scabiosa* subg. *Asterocephalus*, i.e., *S. deserticola* Rech. f., *S. poecilocarpa* Rech. f., *S. transcaspica* Rech. f., as well as for another eastern *Scabiosa* species present in Iran, *S. schimperiana* Boiss. & Buhse. These four taxa show eight pits at the summit of the epicalyx tube, as indicated by Lack and Rechinger in both the descriptions and illustrations of all four taxa (Lack & Rechinger, 1991); this is a synapomorphic condition for *Lomelosia* (Caputo & Cozzolino, 1994) as well. Such transfer should complete the relocation of all known *Asterocephalus* taxa into *Lomelosia*.

It may be interesting to note alternate taxonomic placements for the transferred names. Jamzad (1993), who keeps a traditional and broad concept of genus *Scabiosa*, regarded *S. deserticola* as a synonym of *S. olivieri* Coult., *S. poecilocarpa* as a synonym of *S. persica* Boiss. var. *rosea* Jamzad, and *S. transcaspica* as a synonym of *S. flavida* Boiss. & Hausskn. On the other hand, Czerepanov (1995) accepted independent species rank for *S. deserticola* and *S. transcaspica*. However, we would like to note that taxonomic concepts employed in dipsacaceous taxa can be quite narrow, as testified by the fragmentation of *Scabiosa* and *Pterocephalus* (López Gonzáles, 1987; Mayer & Ehrendorfer, 2000), and species, especially within the *Scabiosa* group, often rest on small differences (e.g., within the *S. columbaria* L. group).

Lomelosia deserticola (Rech. f.) P. Caputo & Del Guacchio, comb. nov. Basionym: *Scabiosa deserticola* Rech. f., Willdenowia 19(1): 142. 1989. TYPE: "Iraq, Desertum occidentale (Western Desert): in arenosis 2 km E Rutba versus Ramadi, 28.5.1957," K. H. Rechinger 12806 (holotype, W not seen).

Habitat and distribution. *Lomelosia deserticola* is found in the deserts of Syria, Jordan, Iraq, Iran, Turkmenistan, Afghanistan, and Pakistan. A detailed distribution, list of specimens, and photographs are provided by Rechinger (1989) and Lack and Rechinger (1991).

Specimen examined. IRAQ. Desertum occidentale (Western desert), inter Ramadi et Rutba, 40 km a Ramadi occidentem versus, 6–7 June 1957, Rechinger 9817 (B [photo]).

Lomelosia poecilocarpa (Rech. f.) P. Caputo & Del Guacchio, comb. nov. Basionym: *Scabiosa poecilocarpa* Rech. f., Fl. Iranica [Rechinger], 168: 50. 1991. TYPE: "Persia. Bakht.: Inter Dashtak et Cherry 13.VI.1973," M. Iranshahr & M. Moussavi 15657-E (holotype, W-photo!; isotype, Herbarium of Evin, Tehran, Iran, not seen).

Habitat and distribution. *Lomelosia poecilocarpa* is found in dry, open mountain habitats of Chaharmahal and Bakhtiari Province (southwestern Iran), where it is endemic.

Specimen examined. See above under *Lomelosia deserticola*. Photograph in Lack and Rechinger (1991: tab. 45).

Lomelosia schimperiana (Boiss. & Buhse) P. Caputo & Del Guacchio, comb. nov. Basionym: *Scabiosa schimperiana* Boiss. & Buhse, Nouv. Mém. Soc. Imp. Naturalistes Moscou 12: 112–113. 1860. TYPE: "Persia. 1847, Buhse s.n." (lectotype, designated by Rechinger in Lack & Rechinger [1991: 48], G-BOIS, G00330078-photo!).

The original label reports the name "*Scabiosa Schimperiana* Boiss. et Buhse" as well as the unpublished "*Scabiosa Buhseana* Boiss." The specimen was relabeled by K. H. Rechinger in 1988, who added the locality "Gilan: Rudbar" and the indication "holotypus." Note that in Lack and Rechinger (1991), the treatment of this species (and of all the annual *Scabiosa* species) is authored by Rechinger alone.

Habitat and distribution. *Lomelosia schimperiana* is found in open environments of Gilan Province

(northwestern Iran), where it is endemic. A distribution, list of specimens, and photograph are provided by Lack and Rechinger (1991).

Specimen examined. See above under *Lomelosia deserticola*. Photograph in Lack and Rechinger (1991: tab. 43).

Lomelosia transcaspica (Rech. f.) P. Caputo & Del Guacchio, comb. nov. Basionym: *Scabiosa transcaspica* Rech. f., Willdenowia 19(1): 148. 1989. TYPE: Turkmenistan. "USSR, Turcomania, Kizyl Arvat, Karakala, in monte Sundsodagh, 16.5.1891," *P. Sintenis 1694-a* (holotype, W photo; isotype, B, WU not seen).

Habitat and distribution. *Lomelosia transcaspica* is found in the mountains of Turkmenistan, Azerbaijan (including Nakhchivan), and the Caucasus region. A distribution, list of specimens, and photographs are provided in Rechinger (1989) and Lack and Rechinger (1991).

Specimen examined. See above under *Lomelosia deserticola*. Photograph in Lack and Rechinger (1991: tab. 53).

Literature Cited

- Adanson, M. 1763. Familles des plantes. P. 152, Vol. 2. Vincent, Paris.
- Avino, M., G. Tortoriello & P. Caputo. 2009. A phylogenetic analysis of Dipsacaceae (Dipsacales) based on four DNA regions. *Pl. Syst. Evol.* 279: 69–86.
- Beck, G. 1893. Flora von Niederösterreich. Vol. 2(2): 1145. Wien.
- Caputo, P. & S. Cozzolino. 1994. A cladistic analysis of Dipsacaceae (Dipsacales). *Pl. Syst. Evol.* 189: 41–61.
- Caputo, P., S. Cozzolino & A. Moretti. 2004. Molecular phylogenetics of Dipsacaceae reveals parallel trends in seed dispersal syndromes. *Pl. Syst. Evol.* 246: 163–175.
- Castro, O. de & P. Caputo. 1999 [2001]. A phylogenetic analysis of genus *Lomelosia* Rafin. (Dipsacaceae) and allied taxa. *Delpinoa* 41: 29–45.
- Coulter, T. 1824. Mémoire sur les Dipsacées. *Mém. Soc. Phys. Genève* 2(2): 13–60.
- Czerepanov, S. K. 1995. P. 216. Vascular Plants of Russia and Adjacent States (the former U.S.S.R.). Cambridge University Press, Cambridge.
- Devesa, J. A. 1984. *Pseudoscabiosa*, género nuevo de Dipsacaceae. *Lagascalia* 12: 213–221.
- Ehrendorfer, F. 1964. Über stammesgeschichtliche Differenzierungsmuster bei den Dipsacaceen. *Ber. Deutsch. Bot. Ges.* 77: 83–94.
- Greuter, W., H. M. Burdet & G. Long (editors). 1986. Med-Checklist. Conservatoire et Jardin botaniques de la Ville de Genève, Genève. 2: 176–198.
- Greuter, W. & T. Raus (editors). 1985. Med-Checklist Notulae: 11. *Willdenowia* 15: 61–64. Conservatoire et Jardin botaniques de la Ville de Genève, Genève.
- Haller, A. 1768. Historia stirpium indigenarum Helvetiae, Vol. 1: 87. Societatis Typographicae, Bern.
- Höck, F. 1891. Dipsacaceae. Pp. 187–189 in H. G. A. Engler & K. A. E. Prantl (editors), Die natürlichen Pflanzenfamilien, IV, Teil. 4. Verlag von Wilhelm Engelmann, Leipzig.
- Jamzad, Z. 1993. *Scabiosa* L. Pp. 63–106 in M. Assadi, M. Khatamsaz & A. A. Maassoumi (editors), Flora of Iran, Vol. 8. Islamic Republic of Iran, Ministry of Jihad-e Sazandegi, Research Institute of Forests and Rangelands, Tehran.
- Jussieu, A. L. de. 1789. Genera plantarum secundum ordines naturales disposita: 194. Herissant & Barrois, Paris.
- Lack, H. W. & K. H. Rechinger. 1991. *Scabiosa*. Pp. 33–64 in K. H. Rechinger (editor), Flora Iranica, Vol. 168. Akademische Druck- und Verlagsanstalt, Graz.
- Linnaeus, C. 1753. Species plantarum, exhibentes plantas rite cognitatas, ad genera relatas, cum Differentiis Specificis, Nominibus Trivialibus, Synonymis Selectis, Locis Natalibus, Secundum systema sexuale digestas, Vol. 1: 98. Impensis Laurentii Salvii, Stockholm.
- López Gonzáles, G. 1987. *Pterocephalidium*, un nuevo género ibérico de la familia Dipsacaceae. *Anales Jard. Bot. Madrid* 43: 245–252.
- Mayer, V. & F. Ehrendorfer. 1999. Fruit differentiation, palynology, and systematics in the *Scabiosa* group of genera and *Pseudoscabiosa* (Dipsacaceae). *Pl. Syst. Evol.* 216: 135–166.
- Mayer, V. & F. Ehrendorfer. 2000. Fruit differentiation, palynology, and systematics in *Pterocephalus* Adanson and *Pterocephalodes*, gen. nov. (Dipsacaceae). *Bot. J. Linn. Soc.* 132: 47–78.
- Moore, D. M. 1976. Dipsacaceae. Pp. 56–74 in T. G. Tutin, V. H. Heywood, N. A. Burgos, D. M. Moore, D. H. Valentine, S. M. Walters & D. A. Webb (editors), Flora Europaea, Vol. 4. Cambridge University Press, Cambridge.
- Rechinger, K. H. 1989. *Scabiosa* sect. *Olivierianae*, sect. nova (Dipsacaceae). *Willdenowia* 19: 137–151.
- Roemer, J. J. & J. A. Schultes. 1818. Systema vegetabilium secundum classes, ordines, genera, species, Vol. 3: 1, 43. Sumtibus J. G. Cotta, Stuttgart.
- Soják, J. 1987. Versetzung einiger asiatischer *Scabiosa*-Arten in *Lomelosia* (Dipsacaceae). *Sborn. Nár. Muz. Praze, Řada B, Přír. Vědy* 43: 57–58.
- Verlák, R. 1984. A biosystematic and phylogenetic study of the Dipsacaceae. Pp. 307–320 in R. Grant (editor), Plant Biosystematics. Academic Press, Toronto.
- Verlák, R. 1985. Etude biosystématique et phylogénétique des Dipsacaceae. II. Caractères généraux des Dipsacaceae. *Rev. Cytol. Biol. Veg. Botaniste* 8: 117–168.
- Verlák, R. 1986a. Etude biosystématique et phylogénétique des Dipsacaceae. IV. Tribus des Scabioseae (phylum n° 1, 2, 3). *Rev. Cytol. Biol. Veg. Botaniste* 9: 5–72.
- Verlák, R. 1986b. Etude biosystématique et phylogénétique des Dipsacaceae. V. Tribus des Scabioseae (phylum n° 4) et conclusion. *Rev. Cytol. Biol. Veg. Botaniste* 9: 97–176.

Caputo, Paolo and Del Guacchio, Emanuele. 2011. "Transfer of Four Species of *Scabiosa* to *Lomelosia* (Dipsacaceae)." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 21(4), 402–404.
<https://doi.org/10.3417/2009135>.

View This Item Online: <https://www.biodiversitylibrary.org/item/180102>

DOI: <https://doi.org/10.3417/2009135>

Permalink: <https://www.biodiversitylibrary.org/partpdf/218512>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Permission to digitize granted by rights holder

Rights: <https://www.biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.