

A RÉSUMÉ OF THE GENUS TIPUANA (LEGUMINOSAE)

Velva E. Rudd

California State University, Northridge and Smithsonian Institution

In the course of tabulating the numerous species originally assigned to Machaerium, but now excluded from that genus, a number of secondary problems have come to my attention. One such involves the now monotypic genus Tipuana (Bentham) Bentham.

Tipuana was first published as a section of Machaerium (Bentham, in Jour. Bot. Kew Misc. 5: 267. 1853) and it included M. heteropterum Freire Allemão, "M. grandiflorum (Benth. in Hook. Journ. Bot. vol. ii. p. 67), and M. Tipu* a hitherto undescribed species found by Tweedie in the Parana." The asterisk refers to a Latin description of "Machaerium (Tipuana) Tipu" and a paragraph in English, "A large free-flowering handsome tree, known in the Parana by the name of Tipú of Peru, and giving its name to the valley of Tipuana, where the best gold in the Parana is found. Tweedie." The appropriate lectotype of Machaerium section Tipuana would seem to be M. tipu.

In the Latin description of M. tipu the inflorescence and the venation of the legumes are compared to those of "M. grandiflori" and the "Corollae majores, fere Platypodii, flavi." The reference to M. grandiflorum possibly was a lapsus calami, or mentis, with Platypodium grandiflorum Bentham in mind. On the page cited by Bentham, in Hook. Jour. Bot. 2: 67. 1840, there is the original description of Machaerium macrocarpum Bentham, later transferred to Tipuana (Jour. Linn. Soc. 4, suppl.: 72. 1860), and erroneously cited as "Machaerium mucronatum" in Martius, Fl. Bras. 15(1): 259. 1862.

Bentham raised Tipuana to generic status in 1860 with the three species, T. macrocarpa, T. heteroptera, and T. speciosa, the latter an illegitimate name based on Machaerium tipu. In 1898 Otto Kuntze made the correct combination, Tipuana tipu (Bentham) O. Kuntze, and that species, I believe, automatically became the lectotype of the genus Tipuana, rather than T. speciosa Bentham (Ducke, in Arch. Jard. Bot. Rio de Janeiro 5: 137. 1930) or T. macrocarpum (Bentham) Bentham as cited by Burkart (Las Leguminosas Argentinas, ed. 1. 544. 1943) and followed by the Index Nomina Genericorum. More recently, Burkart correctly cited T. tipu as the sole species of Tipuana and, indirectly, T. macrocarpa as Vatairea macrocarpa.

A few additional species were assigned to Tipuana before splitting of the genus was begun; Vatairea was resurrected; Luetzelburgia was described as new; other leguminous genera with superficially similar, terminally winged, samaroid fruit, such as Ferreirea, Machaerium, Nissolia, Paramachaerium, and Vataireopsis, have added to confusion in identification, a fact that is reflected in the synonymy of the taxa and the physical location of pertinent specimens in most herbaria. Tipuana finally stands as a monotypic genus, as indicated by Ducke in 1930 (Arch. Jard. Bot. Rio de Janeiro 5: 135).

Having had to unravel this tangled nomenclature, it may be useful to present a list of the species published under Tipuana and to indicate their disposition. The species are given in alphabetical order, the synonymy chronological, and, what I currently believe to be the correct name is emphasized with capital letters.

TIPUANA (Bentham) Bentham, Jour. Linn. Soc. 4, suppl.: 27. 1860; in Martius, Fl. Bras. 15(1): 259. 1862; in Bentham & Hooker, Gen. Pl. 1: 546. 1865. Lectotype: Tipuana tipu (Bentham) O. Kuntze (Machaerium tipu Bentham).

Machaerium section Tipuana Bentham, Jour. Bot. Kew Misc. 5: 267. 1853. Lectotype: Machaerium tipu Bentham.

Trees; bark gray to brown, with irregular, flat-topped ridges; young stems sericeous, glabrescent; unarmed; leaves alternate to sub-opposite, pinnately compound, about 11-31-foliolate; leaflets alternate to sub-opposite, elliptic or elliptic-oblong, about 2-5 cm. long, 1-2 cm. wide, slightly emarginate, rounded at the base, glabrous above, moderately pubescent below, glabrescent; stipules small, linear, caducous; stipels lacking; inflorescences terminal or axillary, paniculate; bracts and bracteoles minute, caducous; flowers papilionoid, about 2 cm. long; calyx 5-8 mm. long, moderately pubescent to subglabrous, campanulate with 5 subequal lobes; petals yellow to light-orange; stamens 10, diadelphous 9:1, the anthers uniform, dorsifixed; ovary brevi-stipitate, 1-4-ovulate; style glabrous; stigma small, truncate; fruit samaroid, about 6-8 cm. long including stipe 1 cm. long, the basal, fertile body 1.5-2 cm. long, 1-1.5 cm. wide, the terminal wing striate, 3.5-5 cm. long, 2-2.5 cm. wide; $n = 10$.

Distribution: Native to subtropical forests of Bolivia and northwestern Argentina; introduced as an ornamental in warm temperate and subtropical areas of Brazil, Uruguay, Europe, and the United States.

1. T. amazonica Ducke, Arch. Jard. Bot. Rio de Janeiro 1: 35. 1915
= VATAIREA MACROCARPA (Benth) Ducke, fide Ducke, Arch. Jard.
Bot. Rio de Janeiro 5: 192, 197. 1930.
2. T. auriculata Freire Allemão, Trab. Comm. Sc. Expl. Bot. Rio de
Janeiro 21. 1862-66.
Luetzelburgia pterocarpoides Harms, Bericht. Deutsch. Bot. Ges.
40: 178. 1922.
Bowdichia (?) freirei Ducke, Arch. Bot. Rio de Janeiro 3: 133.
1922.
LUETZELBURGIA AURICULATA (Freire Allemão) Ducke, Notizbl. 11. 584.
1932.
3. T. cinerascens (Benth) Malme, Arkiv. Bot. 18, no. 17: 14. 1924.
Andira cinerascens Martius ex Benth, Jour. Linn. Soc. 4, suppl.:
72. 1860, nomen in synon.
T. macrocarpa var. cinerascens Benth, in Martius, Fl. Bras.
15 (1): 260. 1862.
VATAIREA MACROCARPA var. CINERASCENS (Benth) Ducke, Arch. Jard.
Bot. Rio de Janeiro 5: 139. 1930.
4. T. erythrocarpa Ducke, Arch. Jard. Bot. Rio de Janeiro 3: 152,
pl. 11a. 1922.
VATAIREA ERYTHROCARPA (Ducke) Ducke, Arch. Jard. Bot. Rio de Jan-
eiro 5: 139, 192, pl. 12, fig. 25. 1930.
5. T. fusca Ducke, Arch. Jard. Bot. Rio de Janeiro 4: 78. 1925.
VATAIREA FUSCA (Ducke) Ducke, Arch. Jard. Bot. Rio de Janeiro 5:
139, 192. 1930.
6. T. heteroptera (Freire Allemão) Benth, Jour. Linn. Soc. 4,
suppl.: 72. 1860.
Machaerium heteropterum Freire Allemão, Trab. Soc. Vell. 4. 1852.
VATAIREA HETEROPTERA (Freire Allemão) Ducke ex de Assis Iglesias,
Album Floristico, Minist. Agric. Serv. Flor. Brasi. 51. 1940.
7. T. lundellii Standley, Carnegie Inst. Wash. Publ. 461: 65. 1935.
VATAIREA LUNDELLII (Standley) Killip ex Record, Trop. Woods no. 63.
5. 1940.
8. T. macrocarpa (Benth) Benth, Jour. Linn. Soc. 4, suppl.: 72.
1860.
Machaerium macrocarpum Benth, Hook. Jour. Bot. 2: 67. 1840.
VATAIREA MACROCARPA (Benth) Ducke var. MACROCARPA, Arch. Jard.
Bot. Rio de Janeiro 5: 141. 1930.

9. T. macrocarpa var. cinerascens Benth, in Martius, Fl. Bras. 15(1): 260. 1862.
Andira cinerascens Martius ex Benth, Jour. Linn. Soc. 4, suppl.: 72. 1860, nomen in synon.
Tipuana cinerascens (Benth) Malme, Arkiv. Bot. 18, no. 17: 14. 1924.
VATAIREA MACROCARPA var. CINERASCENS (Benth) Ducke, Arch. Jard. Bot. Rio de Janeiro 5: 139. 1930.
10. T. mucronata (Benth) Macbride, Candollea 6: 10. 1934.
Machaerium mucronatum Benth, in Martius, Fl. Bras. 15(1): 259. 1862, nomen in synon., a typographical error for M. macrocarpum Benth, non M. mucronatum Vogel, 1837.
= VATAIREA MACROCARPA (Benth) Ducke.
11. T. (?) praecox Harms, in O. Kuntze, Rev. Gen. 3(2): 72. 1898.
Machaerium praecox (Harms) K. Schumann, Just. Bot. Jahresb. 26: 354. 1900.
Ferreireia praecox (Harms) Malme, Arkiv. Bot. Stockh. 18, no. 17, 18. 1924.
LUETZELBURGIA PRAECOX (Harms) Harms, Ber. Deutsch. Bot. Ges. 40: 177. 1922; 43: 595. 1926.
12. T. sericea Ducke, Arch. Jard. Bot. Rio de Janeiro 4: 79. 1928.
VATAIREA SERICEA (Ducke) Ducke, Arch. Jard. Bot. Rio de Janeiro 5: 139, 192, pl. 12, fig. 27. 1930.
13. T. speciosa Benth, Jour. Linn. Soc. 4, suppl.: 72. 1860. Based on Machaerium tipu Benth = TIPUANA TIPU (Benth) O. Kuntze.
14. T. TIPU (Benth) O. Kuntze, Rev. Gen. 3(2): 72. 1898.
Machaerium tipu Benth, Jour. Bot. Kew Misc. 5: 267. 1853.
Tipuana speciosa Benth, Jour. Linn. Soc. 4, suppl. 72. 1860.
Machaerium fertile Grisebach, Goett. Abh. 19: 127. 1874.
Tipuana tipa Lillo, Contrib. Conoc. Arb. Argent. 58. 1910.



Rudd, Velva E. 1974. "A résumé of the genus *Tipuana* (Leguminosae)." *Phytologia* 28(5), 475–478.

View This Item Online: <https://www.biodiversitylibrary.org/item/47049>

Permalink: <https://www.biodiversitylibrary.org/partpdf/219304>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.