## STUDIES IN THE EUPATORIEAE (ASTERACEAE). CLVII. A NEW GENUS, REVEALIA FROM MEXICO

R. M. King and H. Robinson Smithsonian Institution, Washington, D.C. 20560

The generic revisions of the Eupatorieae for Mexico have been considered complete but a totally undescribed taxon has been collected recently in Guerrero by Drs. James Reveal and R. C. Broome with a group from the University of Maryland, College Park. The region is in the Sierra Madre del Sur, an area that has proven very floristically rich, and which is adjacent to regions where George B. Hinton made many of his collections.

The new genus cannot be mistaken for any of the previously known genera of the tribe though there is a vegetative habit resemblance to Oxylobus arbutifolius (H.B.K.) A. Gray but no close relationship. Actual relation can be seen in details of the anthers. The pollen is unusually large at 30 µ in diameter and the median exothecial cells often have transverse walls without thickenings. Both characters are unique in the Eupatorieae to the group of genera including Cronquistia and Carphochaete. The new genus also shows the terete branches of the style with stigmatic lines toward the inside. The style type has previously been regarded as unique to Carphochaete, but the stigmatic lines do not reach near the tip of the style branch as in Carphochaete. The pappus and achene of the new genus are more similar to Cronquistia. The base of the style is distinctly enlarged which contrasts with Cronquistia but the enlargement is not as great as in Carphochaete.

The new genus differs from both <u>Cronquistia</u> and <u>Carphochaete</u> in a few features such as habit and the obtuse anther appendage without an apical notch. The most important difference is the presence of hairs on the inner surface of the corolla. In this trait the new genus is most similar to <u>Stevia</u>. The heads of <u>Revealia</u> also show asymmetry of the peripheral flowers which have the three outer lobes longer, a character also in some species of <u>Stevia</u>. In precise relationship <u>Revealia</u> seems to diverge from between <u>Carphochaete</u> and <u>Cronquistia</u> and the features in common with <u>Stevia</u> might be parallelisms.

Revealia stevioides R.M.King & H.Robinson, n. gen et sp (Eupatorieae) Asteracearum. Plantae frutescentes patentes usque ad 0.3 m altae, ramis patentibus dense

erecte ramulosis. Caules teretes vel subangulares virides minute rubro-puberuli, internodis plerumque 5-20 mm longis. Folia opposita sessilia anguste oblonga plerumque ca. 12 mm longa et 3-4 mm lata carnosa supra et subtus glandulo-punctata margine sparse minute puberula, nervis secundariis longitudinalis submarginalis indistinctis. Capitula in apicem ramularum solitaria, ca. 2 cm alta, pedicellis brevibus 3-10 mm longis; squamae involucri ca. 10 eximbricatae biseriatae ca. 7 mm longae 3-4 mm latae margine distincte scariosae apice abrupte mucronatae extus glandulopunctatae subtiliter-multistriatae; flores ca. 10 in capitulo; corollae purpureae ca. 18 mm longae angusteinfundibulares extus glanduliferis, faucis intus hirsutis, lobis 5 intus papillosis extus laevibus, lobis 3 exterioribus longioribus ca. 5.5 mm longis, lobis interioribus ca. 4 mm longis; filamenta in parte superiore incrassata ca. 0.4 mm longa, cellulis oblongis, parietibus leniter annulate ornatis; thecae ca. 2.5 mm longae base breviter productae rotundatae, cellulis exothecialibus parietibus transversalibus in parte non noduliferis; appendices breviter ovatae obtusae tenues ca. 0.45 mm latae et 0.35 mm longae margine minute crenulatae; basi stylorum distincte nodulosi glabri; appendices stylorum teretes dense papillosae, lineis stigmaticis in superficiis interioribus. Achaenia prismatica ca. 7 mm longa 5-costata in costis minute setifera; carpopodia brevia, cellulis minutis incrassatis ca. 6-seriatis; pappus brevis coroniformis ca. 0.4 mm longus laceratus. Grana pollinis ca. 30-32µ diam. breviter spinosa. Species typica: Revealia stevioides R.M.K et H.R.


TYPE: MEXICO: Guerrero: Sierra Madre del Sur, along the Millpillan-Atoyac road via Puerto del Gallo, about 60.5 miles northeast of Atoyac and 6.5 miles northeast of Puerto del Gallo, on steep rocky slopes and cliff faces, associated with numerous shrubs and scattered trees at about 10,500 ft elevation. Low spreading shrub to 3.0 m high with long, spreading branches; flowers purple. 19 October 1975. Reveal et al 4319 (Holotype US).

## Acknowledgement

This study was supported in part by the National Science Foundation Grant BMS 70-00537 to the senior author.



Revealia stevioides R.M.King & H.Robinson, Holotype, United States National Herbarium. Photos by Victor E. Krantz, Staff Photographer, National Museum of Natural History.



Revealia stevioides R.M.King & H.Robinson, enlargement of head.



King, Robert Merrill and Robinson, Harold E . 1976. "Studies in the Eupatorieae (Asteraceae). CLVII. A new genus, Revealia from Mexico." *Phytologia* 33(4), 277–280.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/47086</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/219360</u>

Holding Institution New York Botanical Garden, LuEsther T. Mertz Library

**Sponsored by** The LuEsther T Mertz Library, the New York Botanical Garden

**Copyright & Reuse** Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: Phytologia License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.