SIGNIFICANT COLLECTIONS OF LOUISIANA PLANTS. III. DESOTO PARISH

R. Dale Thomas and David Lee Dixon. Biology Department, Northeast Louisiana University, Monroe, La. 71209.

A survey of the vascular plants of DeSoto Parish was made from September 1978 to October 1980 (Dixon 1980). Plants representing 135 families, 526 genera and 1234 species and varieties were collected or found in literature or in herbaria. Two state records and several rare Louisiana palnts were collected.

The state records are both introductions to the state in the Schrophulariaceae and were not attributed to the state by Pennell (1935) or by later writers. The citations follow:

Chaenorrhinum minus (L.) Lange. Along Kansas City Southern Railroad tracks between La. 175 and Friendship Road in Frierson, Sec. 34, T15N, R13W. R. Dale Thomas and David Dixon. 64384 & 1301. 6 May 1979. Specimens were collected again at the same site in 1980. Thomascollected this species from Jefferson, Ouachita, and Caddo parishes in 1979 and Winn Parish in 1980.

Parentucellia viscosa (L.) Caruel. Roadside of La. 172 at Bushneck Bayou, 2.7 miles east of La. 169 west of Keatchie, Sec. 25, T14N, R16W. R. Dale Thomas & David Dixon 71368 & 3118. 30 May 1980.

Numerous uncommon plants were collected along the creeks in the hills of DeSoto Parish. These include <u>Glinus radiatus</u> (R. & P.) Rohrb., <u>Zizia aurea</u> (L.) W. D. J. Koch., <u>Amsonia glaberrima</u> Woods., <u>Aristolochia</u> <u>tomentosa Sims, Solidago auriculata Shuttlew., Cayaponia</u> <u>quinqueloba</u> (Raf.) Shinners, <u>Forestiera ligustrina</u> (Michx.) Poir., <u>Pentodon pentandrous</u> (Schum. & Thonn.) Vatke., <u>Parietaria pensylvanica Muhl. ex Willd, and Ranunculus carolinianus DC. The northernmost collection of <u>Mikania</u> <u>cordifolia</u> (L.) Willd. ever made in the state was made. <u>Sparganium americanum</u> was collected in one creek.</u> Thomas & Dixon, DeSoto Parish plants

Although the practice of spraying herbicides on the railroad rights-of-way has probably drastically decreased the populations of many plants, several rare plants still occur along the railroads in the parish. In addition to Chaenorrhinum minus, there are plants of: Eurytaenia texana T. & G., Spermolepis divaricata (Walt.) Raf., Spermolepis echinata (Huds.) Link., Camelina microcarpa Andrz. ex DC., Thlaspi arvense L., Agrostemma githago L., Vaccaria pyramidata Medicus, Salsola kali L., Boerhaavia coccinea Mill., Cissus incisa (Nutt.) Des. Moul., Solanum dimidiatum Raf., Solanum eleagnifolium Cav., and Solanum rostratum Dun. Scutellaria drummondii Benth., Boerhaavia erecta L., Plantago wrightiana Dcne, and Froelichia gracilis are much more common along Louisiana railroads than originally thought. Camelina microcarpa Andrz. ex DC. has been collected in Bossier and Caddo by Thomas, in Sabine by Carroll, and in St. Charles by Glen Montz. Vaccaria pyramidata Medicus (Saponaria vaccaria (L.) Britt.) was reported from Plaquemines Parish by Langlois in 1979 but had not been collected again (?) until recently collected in Caddo Parish by Thomas.

The cemeteries of the parish were searched for small palnts and yielded <u>Ammoselinum butleri</u> (Walt.) Coult. & Rose, <u>Arenaria patula Michx.</u>, <u>Trifolium</u> <u>carolinianum Michx.</u>, and <u>Phacelia glabra Nutt as</u> well as other more common plants.

Roadbanks yielded Torilis nodosa (Huds.) Link., <u>Artemisia ludoviciana Nutt., Cichorium intybus L.,</u> <u>Conyza bonariensis (L.) Cronq., Helianthus grosse-</u> <u>seratus Martens, Rudbeckia nitida Nutt. var. texana</u> <u>Perdue, Silphium laciniatum L., Silene gallica L.,</u> <u>Astragalus distortus T. & G., Trifolium arvense L.,</u> <u>Phacelia hirsuta Nutt., Anemone caroliniana L.,</u> <u>Ranunculus arvensis L., Ranunculus marginatus d'Urv.,</u> <u>Ranunculus trilobus Desf.</u>

Five uncommon plants were collected in pastures: <u>Acorus calamus L., Carduus nutans L., Silybium marianum</u> (L.) Gaertn., <u>Sisymbrium officinale</u> (L.) Scopoli, and <u>Petalostemum purpureum</u> (Vent.) Rydb.

One healthy plant of <u>Cannabis</u> <u>sativa</u> L. and <u>several</u> populations of <u>Chenopodium</u> <u>botrys</u> L. and <u>Melochia</u> <u>corchorifolia</u> L. were found in waste areas. <u>Melochia</u> is common in south Louisiana but is rare in the northern part of the state.

1981

PHYTOLOGIA Vol. 47, No. 5

Several different uncommon plants were collected in the various wooded habitats of the parish. These include: Solidago ulmifolia Muhl., Burmannia biflora L., Symphoriocarpus orbiculatus Moench., Stellaria graminea L., Lathyrus venosus Muhl. ex Willd., Trillium gracile J. D. Freeman, Uvularia perfoliata L., Malaxis unifolia Michx., Huchera americana L., and Ribes curvatum Small.

The Ophioglossaceae is well represented in DeSoto Parish with four of the five species of Botrychium and all five of the Ophioglossum species found in the state occurring there. One cemetery has Ophioglossum nudicaule, O. engelmanni, O. crotalophoroides, Botrychium lunarioides, and B. biternatum in it (See (Thomas 1971, 1972). Botrychium dissectum var. obliquum, B. biternatum, B. virginianum and Ophioglossum vulgatum occur in woods and O. petiolatum grows in moss around rural church buildings.

LITERATURE CITED

- Dixon, David Lee. 1980. A preliminary survey of the vascular plants of DeSoto Parish, Louisiana. Unpublished Masters of Science Thesis, Northeast Louisiana University, Monroe. 151 pp.
- Langlois, A. B. 1879. Catalog provisiore de plantes, phanerogames et cryptogames de la Basse-Louisiane, Etats-Unis d' Amerique, Pointe-a-la-Hache. 35 pp.
- Pennell, F. W. 1935. The Scrophulariaceae in eastern temeprate North America. The Academy of Natural Sciences of Phialdelphia, Philadelphia. 650 pp.
- Thomas, R. D. 1971. Bulbous adder's tongue common in Louisiana. American Fern Journal 71: 39-41
- Thomas, R. D. 1972. Botrychium lunarioides, Ophioglossum crotalophoroides, and Ophioglossum engelmanni in a Louisiana cemetery. The Southwestern Naturalist 16: 431-432.

370



Thomas, R. Dale and Dixon, D P . 1981. "Significant collections of Louisiana plants. III. Desoto Parish." *Phytologia* 47(5), 368–370.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/47388</u> **Permalink:** <u>https://www.biodiversitylibrary.org/partpdf/219864</u>

Holding Institution New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: Phytologia License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.