

SOLANUM VENATORIS (SECT. PETOTA)
A NEW SPECIES FROM BOLIVIA

C. Ochoa

Department of Taxonomy of the International
Potato Center, P.O. Box 5969, Lima, Perú


Solanum venatoris Ochoa sp. nov.

Herbaceum, tuberiferum. Plantae 30-40 cm altae, caules erecti, gracili, 2.5-3.5 mm crassi, plerumque simplici, sparsim pilosi, pilis difficulter manifesti. Caules anguste alati, alae rectae. Stolones plus quam 1 m longi, 1.5-2.0 mm crassi, albidi; tubercula parva 2.0-2.5 cm diam., rotunda, alba. Folia imparipinnata, breviter lataque, 10.5-13.0 x 8.0-9.5 cm petioli 1.0-2.5 cm longi, folia 2-3-juga, 0-4 foliolis interjectis instructa, rarissime 4-juga, foliolis interjectis 6 vel plus. Foliola supra viridia vel obscurius viridia et sparse pilosa, subtus dilute viridia pilis brevibus et densioribus praedita. Foliolum terminale caeteris sensim majus atque latius, 5.5-6.2 x 3.2-4.0 cm, elliptico-lanceolatum, apice acuminatum, basi rotundatum vel asymmetrice rotundatum. Foliola lateraliter elliptico-lanceolata, apice et base tam quam foliolum terminale, petioluli 1.5-3.0 mm longi, foliola interjecta 1.5-6.0 x 1.0-3.5 mm. Foliola pseudostipulacea parva anguste subfalcata usque ad triangulariter-lanceolata, 2.5-4.0 x 1.5-2.0 mm. Inflorescentia cymoso-paniculata, 5-6-flora, pedunculus 5 cm longus, basi 1.5 mm crassus et sparse pilosus, dilute viridus tam quam pedicelli et calyx; pedunculus ad 1/3 superum vel prope calycem articulatus, pedicellus superior 5-6 mm longus, inferior 12-30 mm longus. Calyx 8-9 mm longus, lobi anguste elliptico-lanceolati, in acumina attenuati, acumina recurva, angusta atque acuta 5.5-7.0 mm longa. Corolla parva, rotata vel pentagona, alba 2.0-2.5 cm diam, stella viride-flava. Antherae anguste lanceolatae 5.5-6.0 mm longae, basi cordatae. Filamenta glabra 0.5-0.8 mm longa. Stylus 10 mm longus 2/3 inferioris papillis obtectum, stigma parvum, styli apice vix crassius. Baccae globosae, virides, difficulter 10 mm diam. Ad seriem tuberosa pertinet, numerus cromosomatum $2n=2x=24$.

Typus: BOLIVIA, Department La Paz, province Inquisivi in itinere Quime-Inquisivi, 2500 m alt. Crecit inter frutices et ad margines silvarum. Martius 1978, C. Ochoa 11917. Holotypus: Herb. Ochoanum (OCH). Isotypi: CIP, US.

Paratypes: BOLIVIA, Department Santa Cruz, province Valle Grande, prope Alto Grande, 2750 m alt., in itinere Valle Grande-Pucará, C. Ochoa et A. Salas 15554 (CIP, OCH, US).

Omnibus, Solana tuberifera sponte crescencia venatores, et qui haec loci visitaverunt, ex toto corde dedico.



Solanum venatoris Ochoa. Holotypus
OCH 11917, ca x 1/3



Ochoa, C. 1984. "Solanum venatoris (sect. Petota) a new species from Bolivia." *Phytologia* 55(5), 297–298.

View This Item Online: <https://www.biodiversitylibrary.org/item/46285>

Permalink: <https://www.biodiversitylibrary.org/partpdf/220071>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.