

<i>Chimaphila maculata</i> ,	<i>Larix Europea</i> , seedlings.	<i>Quercus prinoides</i> ,
<i>Cerastium nutans</i> ,	<i>Lespedeza hirta</i> ,	<i>Quercus ilicifolia</i> ,
<i>Cerastium vulgatum</i> ,	<i>Lobelia inflata</i> ,	<i>Rosa lucida</i> ,
<i>Chimaphila umbellata</i> ,	<i>Luzula campestris</i> ,	<i>Ribes hirtellum</i> ,
<i>Carex stipata</i> ,	<i>Lycopus Virginicus</i> ,	<i>Rubus villosus</i> ,
<i>Castanea vesca</i> ,	<i>Lysimachia quadrifolia</i> ,	<i>Rubus Canadensis</i> ,
<i>Ceanothus Americanus</i> ,	<i>Lycopodium clavatum</i> ,	<i>Rudbeckia hirta</i> ,
<i>Carex scoparia</i> .	<i>Lycopodium complanatum</i> ,	<i>Rumex acetosella</i> ,
<i>Carex Muhlenbergii</i> ,	<i>Lycopodium dendroideum</i> ,	<i>Salix humilis</i> ,
<i>Carex rosea</i> .	<i>Lycopodium lucidulum</i> ,	<i>Sambucus Canadensis</i> ,
<i>Onicis lanceolatum</i> .	<i>Magnolia acuminata</i> ,	<i>Sassafras officinale</i> ,
<i>Comptonia asplenifolia</i> ,	<i>Nyssa multiflora</i> ,	<i>Saxifraga Virginienis</i> ,
<i>Cornus paniculata</i> ,	<i>Oenothera biennis</i> ,	<i>Senecio aureus</i> ,
<i>Cornus alternifolia</i> ,	<i>Oenothera pumila</i> ,	var <i>obovatus</i> ,
<i>Conium maculatum</i> ,	<i>Oxalis Acetosella</i> ,	<i>Sisyrinchium Bermudiana</i> ,
<i>Cratægus coccinea</i> ,	<i>Panicum autumnale</i> ,	<i>Setaria glauca</i> ,
<i>Danthonia spicata</i> ,	<i>Panicum depauperatum</i> ,	<i>Solidago arguta</i> ,
<i>Epigæa repens</i> ,	<i>Panicum latifolium</i> ,	<i>Solidago bicolor</i> ,
<i>Eragrostis capillaris</i> ,	<i>Phleum pratense</i> ,	<i>Solidago casia</i> ,
<i>Erigeron Canadense</i> ,	<i>Pinus strobus</i> ,	<i>Solidago Canadensis</i> ,
<i>Erigeron bellidifolium</i> ,	<i>Pinus rigida</i> ,	<i>Solidago ulmifolia</i> ,
<i>Erigeron Philadelphicum</i> ,	<i>Phytolacca decandra</i> ,	<i>Solidago nemoralis</i> ,
<i>Euphorbia corollata</i> ,	<i>Poa brevifolia</i> ,	<i>Solidago squarrosa</i> ,
<i>Euphorbia maculata</i> ,	<i>Poa pratensis</i> ,	<i>Spiranthes gracilis</i> ,
<i>Fragraria Virginiana</i> ,	<i>Polygonum Pennsylvanicum</i> ,	<i>Thalictrum anemonoides</i> ,
<i>Gaylussacia resinosa</i> ,	<i>Polygonatum biflorum</i> ,	<i>Trifolium repens</i> ,
<i>Gnaphalium decurrens</i> ,	<i>Populus grandidentata</i> ,	<i>Trifolium pratensis</i> ,
<i>Gnaphalium polycephalum</i> ,	<i>Potentilla Canadensis</i> ,	<i>Urticaria perfoliata</i> ,
<i>Hedeoma pulegioides</i> ,	<i>Prunus serotina</i> ,	<i>Verbascum thapsus</i> ,
<i>Hieracium scabrum</i> ,	<i>Pyrola rotundifolia</i> ,	<i>Vaccinium Pennsylvanicum</i> ,
<i>Hieracium venosum</i> ,	<i>Pyrola chlorantha</i> ,	<i>Vaccinium stamineum</i> ,
<i>Hieracium paniculatum</i> ,	<i>Pyrola elliptica</i> ,	<i>Vaccinium vacillans</i> ,
<i>Hypericum corymbosum</i> ,	<i>Pyrola secunda</i> ,	<i>Viola cucullata</i> ,
<i>Hypericum perforatum</i> ,	<i>Pyrus arbutifolia</i> ,	<i>Viola sagittata</i> ,
<i>Juncus tenuis</i> ,	var <i>melanocarpa</i> ,	<i>Vilfa vaginæflora</i> ,
<i>Juncus effusus</i> ,	<i>Pyrus coronaria</i> ,	<i>Vitis cordifolia</i> ,
<i>Juniperus Virginiana</i> ,	<i>Pyrus aucuparia</i> ,	<i>Veronica officinalis</i> ,
<i>Kalmia latifolia</i> ,	<i>Quercus alba</i> ,	<i>Veronica peregrina</i> .—
<i>Lactuca elongata</i> ,	<i>Quercus coccinea</i> ,	J. R. LOWRIE.

POTATO PIERCED BY GRASS.—Mr. A. C. Brigham, of Trumbull, Ohio, sends me a freshly dug potato which has been pierced completely through by a growing stem of grass; probably *Poa pratensis*. The potato is three inches long and two inches thick, of regular shape, and with a smooth, uninjured skin. The grass entered from the side, near the "stem" end of the potato, and issued at the "eye" end, its course being somewhat curved. If it entered vertically, as is probable, it emerged horizontally, being carried forward, perhaps, by the contemporaneous growth of the potato. The original root of the grass was not seen, that part being broken away an eighth of an inch outside of the point of entrance. Dissection reveals the perfect stem of the grass imbedded in the center of the potato. After emergence the grass struck upward in a vigorous shoot five inches beyond the potato, appearing above the ground in half a dozen green leaves. Many long, tender roots extend out from the stem above its point of liberation. The potato suffered no discernible irritation at being thus pierced, and the grass, also, was in no way exhausted by its exploit.—ALBERT A. WRIGHT, Oberlin, Ohio.

NELUMBium LUTEUM.—I just saw specimens of *Nelumbium luteum* from Grass Lake, Lake Co., Illinois. It is a new locality for the plant, which is by no means common here.—M. HARRINGTON, Sycamore, Ill.



Harrington, M . 1878. "Nelumbium luteum." *Botanical bulletin* 3(12), 100–100.
<https://doi.org/10.1086/325199>.

View This Item Online: <https://www.biodiversitylibrary.org/item/27487>

DOI: <https://doi.org/10.1086/325199>

Permalink: <https://www.biodiversitylibrary.org/partpdf/221110>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

MSN

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.