

with coarser appressed hairs in all our specimens. It must be that the first description is wrong on this point. Then, the leaves are in no case "7—8 lines wide," but $\frac{3}{8}$ in. to $\frac{1}{2}$ in.; the Angeles Bay plant having some leaves as wide as $\frac{5}{8}$ in. Furthermore, the description has "Calyx 4-parted . . . Stamens 6 according to Bonpland. . . . Achene lenticular." I have frequently found the calyx 5-parted, stamens as many as 8, and the achenes in the Angeles Bay and the Orizaba plant—the latter from one of the stations cited in Biol. Centr. Am. l. c.—as well as in the Texas plant, are triangular: all, however, of the same size, and all "umbonate," as in the first description.

This species is distinguished from *P. persicaria* L. by its narrower, longer leaves, more slender spikes and smaller achenes.—J. M. HOLZINGER, *Department of Agriculture, Washington, D. C.*

New Mosses of North America.—The following brief diagnoses are published in advance of fuller descriptions in order to secure priority. We hope to prepare shortly the fifth number of our series under the above title, in THE BOTANICAL GAZETTE.

Dichodontium olympicum n. sp.—A *D. pellucida* jam multo robustiore primo visu differt: foliis valde papillois, toto fere ambitu minute denticulatis, capsulaque basi strumosa. Planta humilis, vix 1 cent. alta.—Olympic Mts., Wash. (*L. F. Henderson.*)

Grimmia Hendersoni n. sp.—*G. decipienti* Lindb. (*G. Schultzii* Wils.) proxima sed ab ea pedicello longiore, capsula subcylindrica magis elongata, operculo longius rostrato et foliorum rete basilari multo laxiore facillima distinguenda.—Hood River, Oregon, on dry rocks. (*L. F. Henderson.*)

Encalypta lacera n. sp.—Ab *Enc. vulgari* proxima differt calyptra basi lacerata, peristomio e membrana alba fugacissima lacerata composito et pedicello longiore.—Milwaukie, Oregon, Willamette River, (*L. F. Henderson.*)

Leskea obtusa n. sp.—Formis robustioribus *L. polycarpæ* similis sed ab illis primo aspectu foliis obtusis marginibus planis distincta, A *L. obscura* habitu valde robustiore, foliis majoribus et magis obtusis, costa brevioris et capsula majore et longiore quoque differt.—Bethlehem, Pa., mixed with *Anomodon obtusifolius* (*Rau.*) Chinchuba, La., near Mandeville, on trees (*Langlois.*)—F. RENAULD AND J. CARDOT, *Monaco, and Stenay, France.*



Renauld, Ferdinand and Cardot, Jules. 1892. "New Mosses of North America." *Botanical gazette* 17(9), 296–296. <https://doi.org/10.1086/326841>.

View This Item Online: <https://www.biodiversitylibrary.org/item/92733>

DOI: <https://doi.org/10.1086/326841>

Permalink: <https://www.biodiversitylibrary.org/partpdf/222183>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.