"The maiden hair tree." The American publisher is G. P. PUTNAM'S SONS of New York.—J. M. C.

North American Flora.⁷—Vol. VII, part 3, continues the treatment of the Uredinales and contains the Aecidiaceae from *Prospodium* to *Dichaeoma* by JOSEPH CHARLES ARTHUR, the text for the genus *Gymnosporangium* being contributed by FRANK DUNN KERN. One new genus (*Argomyces*) is proposed, which has a geographical distribution from New Mexico and Texas through Mexico and the West Indies to South America, and is represented at present by four known species. Further new species are characterized in the following genera: *Earlea* (1), *Kuehneola* (1), *Spirechina* (1), and *Xenodochus* (1).—J. M. GREENMAN.

NOTES FOR STUDENTS

Variation curves.—Several years ago papers dealing with variation in the number of parts of flowers, flower heads, inflorescences, etc., were of frequent appearance. As the novelty of the method disappeared, the number of contributors to the knowledge of such variations has decreased, but, as is usually true in such cases, the value of the contributions has correspondingly improved. Several recent studies in this field are of exceptional interest.

VOGLER⁸ gives a large number of counts of ray flowers in *Chrysanthemum*, *Boltonia*, and *Senecio*. In *Chrysanthemum Parthenium* he found a curve having the mode on 21 when the plants were grown on well-manured soil, and on 13 when grown on infertile soil, the curves being strongly skew in each case toward an intermediate point, the mean values lying between 14 and 19. These results agree essentially, therefore, with those of KLEBS⁹ on *Sedum spectabile*. In *Boltonia latisquama* the ray flowers have a wide range of variation (39–81), with the summit of the curve near 55. Three different plants were separately counted in three successive years, and although the different seasons differed considerably, there was no corresponding change in the number of ray flowers. One of these plants had each year the mean number approximately 57, another approximately 54. These permanent differences are probably not to be attributed to genotypic differences in the plants, however, as they originated from a common stock by vegetative division. In *Senecio alpinus* a count of over 3000 heads from two different localities in three different years gave in every case a nearly monomodal curve with the mode on 19, thus convincing the author of the limitations of Ludwig's law that the maxima of such

⁷ North American flora. Vol. VII, part 3, pp. 161–268. The New York Botanical Garden. April 15, 1912.

⁸ VOGLER, P., Variation der Anzahl der Strahlblüten bei einigen Kompositen. Beih. Bot. Centralbl. 25:387–396. 1910.

⁹ KLEBS, G., Studien über Variation. Arch. Entwickl.-Mech. Organ. 24:29–113. 1907.

1912. "North American Flora." *Botanical gazette* 54(1), 77–77.

<https://doi.org/10.1086/330870>.

View This Item Online: <https://www.biodiversitylibrary.org/item/109452>

DOI: <https://doi.org/10.1086/330870>

Permalink: <https://www.biodiversitylibrary.org/partpdf/223472>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.