SPEGAZZINIAN MELIOLA TYPES

F. L. STEVENS

(WITH PLATES XXIV-XXVI)

Through the kindness of Dr. CARLO SPEGAZZINI, I have received recently a number of original packages containing type specimens of *Meliola* described earlier by Dr. SPEGAZZINI. In each case the packet bore on the outside copious penciled notes concerning the specimen and careful, delicate drawings of the more significant structures. This collection of types, together with the notes and drawings in particular, are a fine commentary on the work of Dr. SPEGAZZINI. When we remember the large volume of his descriptive work and reflect that not only his types but other specimens as well are thus thoroughly and carefully annotated and figured, we are in a position more adequately to recognize the great indebtedness of mycology to him.

The drawings of the present collection have not been published, and in view of the comparative inaccessibility of most of these types, it is desirable that they should be made generally accessible to students by publication.

Permission having been received from Dr. SPEGAZZINI, Dr. ALVA PETERSON has faithfully copied for me, for publication, the most important of the drawings. Such copying was necessary, owing to the color of the paper upon which the originals were drawn, and the faintness of the penciling, which prohibited direct photographic reproduction. These drawings have been compared by me with the type material and are published herewith with such comments as seem necessary. A permanent celloidin mount¹ has been made from each specimen and, together with a fragment of the type specimen, is deposited in the herbarium of the University of Illinois. The original specimens have been returned to Dr. SPEGAZZINI at La Plata, Argentine.

MELIOLA ARMATA Speg. (fig. 1).—F. Puigg., Pug. I, no. 231; Sacc. 9:415.

¹ STEVENS, F. L., Phytopathology 6:367. 1916.

On coriaceous leaves, *Myrsine* (?), Apiahy, May 1888; no. 2382 (type).

The type specimen is heavily overgrown with several parasites. There is considerable variation in the character of the mycelium, which is sometimes straight, sometimes quite crooked. I have not been able to see the mycelial setae around the bases of the perithecia from which this species takes its name.

MELIOLA ARGENTINA Speg. (fig. 2).—Fung. Arg., Pug. I, no. 177; Sacc. 1:61.

On Cyperaceae, Buenos Aires, February 1880 (type).

The mycelium is very characteristic, close, dense, somewhat like *M. manca*, but distinguished by its very thick mycelial setae, which are striking, being darker and thicker than the mycelium. They are about $15\ \mu$ thick at base and over $800\ \mu$ long. The capitate hyphopodia are angular, that is, not smooth or echinulate as shown in fig. 2. The type is heavily overgrown by a *Coniothyrium*.

MELIOLA BRASILIENSIS Speg. (fig. 3).—Fung. Arg., Pug. IV, no. 116; Sacc. 1:66.

On leaves of Bignoniaceae (?), Apiahy; no. 1551 (type).

The young perithecia are surrounded by an areola of radiating hyphae. The perithecia also possess short hairs as figured and described by Dr. Spegazzini.

MELIOLA CALVA Speg. (fig. 4).—F. Puigg., Pug. I, no. 233; Sacc. 9:414.

On Laurinaceae, Apiahy, August 1881 (type).

Heavily overgrown with "*Podosporium penicillium* Speg."

MELIOLA CLAVATISPORA Speg. (fig. 5).—F. Puigg., Pug. I, no. 241; Sacc. 9:422.

On leaves of Apocynaceae, Apiahy, April 1881; no. 1701 (type).

Perhaps the most striking character is in the sessile, nearly globular capitate hyphopodia.

MELIOLA CORONATA Speg. (fig. 6).—F. Guar., Pug. I, no. 175; Sacc. 9:428.

On *Luchea divaricata*, Guarapi, July 1883; no. 3847 (type).

The figure shows perithecial hairs to be more conspicuous than they usually are.

MELIOLA CRUSTACEA Speg. (fig. 7).—F. Puigg., Pug. I, no. 235; Sacc. 9:413.

On *Drymis*, Apiahy, 1881 (type).

The mycelium forms a compact crustose colony, with the parts decidedly more crowded even than is shown in fig. 7.

MELIOLA DECIDUA Speg. (fig. 8).—F. Puigg., Pug. I, no. 240; Sacc. 9:426.

On Convolvulaceae (?), Apiahy, April 1888; no. 2344 (type).

The capitate hyphopodia are very irregularly angular.

MELIOLA DELICATULA Speg. (fig. 9).—F. Guar., Pug. II, no. 63; Sacc. 9:415.

On *Myrsinus*, Sierra de Peribebuy, September 15, 1883; no. 3985 (type).

MELIOLA ERIOPHORA Speg. (fig. 10).—F. Guar., Pug. II, no. 62; Sacc. 9:413.

On *Ficus ibapoy*, Paraguay, January 1883 (type).

MELIOLA GLABRIUSCULA Speg. (fig. 11).—F. Aliq. Paul., no. 35; Sacc. 22:48.

On Photiniae (?), Agua branca, Sao Paulo (type).

MELIOLA GLEDITSCHIAE Speg. (fig. 12).—Myc. Argent. VI, no. 1337.

On *Gleditschia amorphoidis*, Puerto Leon, Misiones, July 1909 (type).

MELIOLA GUAREAE Speg. (fig. 13).—Myc. Argent. VI, no. 1338.

On *Guarea balansae*, Puerto Leon, Misiones, August 1909 (type).

MELIOLA HARIOTI Speg. (fig. 14).—F. Guar. nonn. III, no. 78; Sacc. 11:267; Gaill. Bull. Soc. Myc. Fr. 8:186. 1892.

On Bignoniaceae, Paraguay, no. 1291 (type).

MELIOLA LEVIPODA Speg. (fig. 15).—F. Guar. nonn., no. 77 (p. 26); Sacc. 11:264; Bull. Soc. Myc. Fr. 8:181. 1892.

On *Aspidosperma quebracho*, Yaguaron, Paraguay, November 1882; no. 3589 (type).

MELIOLA LUDIBUNDA Speg. (fig. 16).—F. Guar. I, no. 178; Sacc. 9:431.

On *Pilocarpus pinnatus*, Paraguay, January 1882; no. 3489 (type).

MELIOLA MEGALOSPORA Speg. (fig. 17).—F. Arg., Pug. IV, no. 115; Sacc. 1:67.

On *Jodina rhombifolia*, January 1888 (type).

The very coarse hyphae are quite characteristic.

MELIOLA MELASTOMACEARUM Speg. (fig. 18).—F. Puigg., Pug. I, no. 232; Sacc. 9:414.

On Melastomaceae, no. 2485, Apiahy, May 1888 (type).

The mycelium is often less straight than might be assumed from the figure. The oval hyphopodia are characteristic.

MELIOLA OBESA Speg. (fig. 19).—F. Guar., Pug. I, no. 179; Sacc. 9:421.

On Rutaceae, Piragu Bras, July 1883; no. 3834 (type).

MELIOLA OBESULA Speg. (fig. 20).—F. Guar. nonn., no. 75; Sacc. 11:262.

On Rutaceae, Caa-guaza, Brazil, January 1882; no. 3585 (type).

MELIOLA PUIGGARII Speg. (fig. 21).—F. Puigg., Pug. I, no. 228; Sacc. 9:414.

On *Rubus*, Apiahy, May 1888; no. 2722 (type).

MELIOLA PULCHELLA Speg. (fig. 22).—F. Puigg., Pug. I, no. 227; Sacc. 9:414.

On Myrtaceae, Apiahy, 1881; no. 1699 (type).

MELIOLA SAPINDACEARUM (fig. 23).—F. Guar. nonn., III, no. 79; Sacc. 11:266; Bull. Soc. Myc. Fr. 8:184. 1892.

On Sapindaceae, Caa-guazu, Brazil, January 1882; no. 3600.

MELIOLA SPEGAZZINIANA Wint. (fig. 24).—F. Guar., Pug. II, no. 64; Sacc. 9:418.

On Compositae, Paraguari, March 5, 1883; no. 3751 (type).

MELIOLA SORORCULA (fig. 25).—F. Puigg., Pug. I, no. 230; Sacc. 9:418.

On *Baccharis pingrea*, Apiahy, May 1886; no. 2774 (type).

MELIOLA SUBCRUSTACEA Speg. (fig. 26).—F. Puigg., Pug. I, no. 236; Sacc. 9:430.

Apiahy, 1888; no. 2703 (type).

MELIOLA TABERNAEMONTANAE Speg. (fig. 27).—Myc. Argent. VI, no. 1345; Bompland Missiones.

STEVENS on MELIOLA

Stevens, Frank Lincoln. 1917. "Spegazzinian Meliola Types." *Botanical gazette* 64(5), 421–425. <https://doi.org/10.1086/332169>.

View This Item Online: <https://www.biodiversitylibrary.org/item/109348>

DOI: <https://doi.org/10.1086/332169>

Permalink: <https://www.biodiversitylibrary.org/partpdf/223947>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.