or the south Pacific than to America. The American elements that are present are accounted for partly through introduction by winds or migratory birds, and partly as a residue of once more widespread forms that are now extinct except in Hawaii and America. The absence of conifers may similarly be explained by extinction, if they were ever present, or by the absence of suitable soil conditions. The almost complete absence, for example, of granitic or calcareous soils might well explain certain absences. It is noted also that great ocean deeps separate Hawaii from America, whereas it is much shallower between Hawaii and the Orient. It is concluded, therefore, that the Hawaiian flora has been derived for the most part from the southern Pacific region, and that the Hawaiian Islands are a remnant of a northeastern extension of some large land mass, once connected closely with south Pacific lands.—H. C. COWLES.

Studies of cambium.—BAILEY,¹⁴ in a third paper on cambium, has made what he calls a cytological "reconnaissance." In the preceding paper, reviewed in this journal,¹⁵ he called attention to the size variations of cambial initials, and to the unusual opportunity offered by the cambium for the study of a number of fundamental cytological problems. In this preliminary study he has reached the following conclusions. The initials of the cambium, which may attain a length of more than 9000 μ , are uninucleate, and the "working distance" of their nuclei must extend in some cases for a distance of several thousand microns. The nucleo-cytoplasmic ratio may be relatively constant in ray initials, but varies enormously in fusiform initials. All the cambium initials of *Pinus Strobus* contain the diploid number of chromosomes. Small ray initials may contain as large chromosomes as adjacent fusiform initials with a volume 200–1000 times as large. Fusiform initials, which are frequently several hundred times as long as they are wide, divide longitudinally by an extraordinary extension of the cell plate. The various types of cell plate formation described by various cytologists are believed to be merely different phases or stages of a single general type of cytokinesis. These glimpses would seem to justify the writer in his belief that the cambium well deserves intensive cytological investigation.—J. M. C.

Economic plants of Philippines.—In an illustrated report BROWN¹⁶ gives a series of descriptions of the indigenous food-producing plants of the Philippines. Many will be surprised to find the statement that the edible wild plants of these islands are less abundant, more inaccessible, and inferior in

¹⁴ BAILEY, I. W., The cambium and its derivative tissues. III. A reconnaissance of cytological phenomena in the cambium. Amer. Jour. Bot. 7:417–434. pls. 26–29. 1920.

¹⁵ BOT. GAZ. 71:408. 1921.

¹⁶ BROWN, WM. H., Wild food plants of the Philippines. Phil. Dept. Agric. and Nat. Res., Bur. For. Bull. 21:1–165. figs. 81. 1920.

1921. "Studies of Cambium." *Botanical gazette* 72(1), 53–53.

<https://doi.org/10.1086/332884>.

View This Item Online: <https://www.biodiversitylibrary.org/item/109576>

DOI: <https://doi.org/10.1086/332884>

Permalink: <https://www.biodiversitylibrary.org/partpdf/224277>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.