

Five new species of *Adenanthos* (Proteaceae) from Western Australia

By A. S. George

Abstract

The following new species of *Adenanthos* are described and their relationships discussed: *A. acanthophyllus* sp. nov., *A. ellipticus* sp. nov., *A. gracilipes* sp. nov., *A. stictus* sp. nov., and *A. teges* sp. nov.

Introduction

Recognition of the following new species of *Adenanthos* arose during preliminary work on the genus towards the Flora of Western Australia. All belong to the Section *Stenolaema* in which all four anthers are perfect.

Notes on Morphology

In *Adenanthos* the zygomorphic perianth is arranged antero-posteriorly so that there is a posticus and an anticus segment and two lateral ones. The anticus one is the broadest, the posticus the narrowest, with the lateral ones intermediate but more similar to the anticus one. They form a tube in the lower half but separate above. Before anthesis, the posticus segment separates to a lower level than the others but may adhere on either side to the adjacent segment, and the style, except for the pollen presenter, emerges on the more deeply split side. The pollen presenter is held until anthesis within the limb, when the latter opens to free it, bearing the pollen. The limb of the anticus segment recurves more than the others, and its anther may reflex against the claw. A ring of straight, erect hairs usually surrounds the base of the perianth, and another surrounds the ovary.

The gender of *Adenanthos*

The name *Adenanthos*, for many years treated as feminine, is here given masculine gender. The International Code of Botanical Nomenclature (1972), Chapter VI Section 2, Recommendation 75A, states:

“Similarly, all modern compounds ending in *-ceras*, *-dendron*, *-nema*, *-stigma*, *-stoma*, and other neuter words should be neuter. The fact that Robert Brown and Bunge respectively made *Aceras* and *Xanthoceras* feminine is immaterial. An exception should be made for names ending in *-anthos* (or *-anthus*) and *-chilos* (*chilus* or *-cheilos*), which ought to be neuter, since that is the gender of the Greek words *anthos* and *cheilos*, but which have generally been treated as masculine and should have that gender assigned to them.”

In Australia most genera of the latter endings are already treated as masculine, e.g. *Acianthus*, *Angianthus*, *Phyllanthus*, *Calochilus*, *Eriochilus*, etc. In order to further settle nomenclature, the Recommendation has been adopted for some time in Western Australia for *Adenanthos* and is followed here. Similarly, the genus *Anigozanthos* is now treated as masculine (Geerinck, 1970, p. 261).

Adenanthos acanthophyllus A. S. George sp. nov.

Derivation of epithet: Greek, *acantha*, a spine or prickle, and *phyllon*, a leaf.

Frutex multicaulis ad 5 m altus cum lignotubero. *Folio* rhombeo-lunata 15–25 mm longa, 18–40 mm lata, profunde trilobata, lobis denue 2–3-lobatis; lobi triangulari-ovati 4–10 mm longi in spinis pungentibus contracti; laminae crassae aliquantum undulatae marginibus planis venatione furcato-reticulata; petiolis teretibus 5–10 mm longis; folia juvenes pubescentes, deinde glabra.

Flores solitarii in pedunculos 3–4 mm longos. *Bractee* ovato-lanceolatae, acutae, 2–8 mm longae, appresso-pubescentes. *Perianthium* 24–28 mm longum limbo acuto 2–3 mm longo, intus glabrum, extus pilis brevibus longisque pubescens. *Antherae* sessiles 2 mm longae. *Squamae hypogynae* oblongae 2.5 mm longae. *Ovarium* breviter pubescens. *Stylus* glaber. *Fructus* fusiformis, obtuso-truncatus, 5–7 mm longus, breviter pubescens.

Type: Nanga Station, on road to Tamala, 5 miles SW of turnoff from Denham road, Western Australia—26°34'S, 113°55'E, 15 July 1973, A. S. George 11671. Holo: PERTH, iso: CANB, K, MEL, NSW, PERTH.

A broad, dense shrub to 3 m tall, occasionally to 5 m, with many stout, erect or spreading stems arising from a lignotuber. Bark smooth, light grey. Branchlets terete, yellowish, pubescent with short curled hairs and a few \pm straight ones, becoming glabrous after a few years. Leaves deeply 3-lobed, each lobe again 2–3-lobed, or when small simply 3-lobed, rarely entire; rhombic to lunate in outline, 15–25 mm long, 18–40 mm wide, the lobes often unequal, triangular-ovate, 4–10 mm long, contracted into a pungent, acuminate spine; lamina \pm undulate, sclerophyllous, margins flat, venation furcate-reticulate, pubescent when young, becoming glabrous; petiole terete, 5–10 mm long, pubescent.

Flowers pale pink and green, solitary, axillary in the uppermost axils; peduncles 3–4 mm long, bearing a few scattered, ovate bracts less than 1 mm. Involucral bracts ovate-lanceolate, acute to acuminate, the outermost 2 mm long, the innermost 7–8 mm, appressed-pubescent. Perianth 24–28 mm long including the limb of 2–3 mm; tube slightly swollen above base, then narrowed to the limb which is acute; tube moderately pubescent with short and long, straight to curled hairs, the limb with mostly long straight hairs; perianth glabrous within. Anthers sessile, 2 mm long, the connective broad, slightly produced, obtuse. Hypogynous scales oblong, 2.5 mm long, shortly united to the perianth. Ovary shortly pubescent. Style glabrous; pollen presenter 2 mm long, flattened on the lower side but the apex laterally compressed, otherwise smooth and rounded, the stigmatic groove terminal.

Fruit fusiform but swollen, obtuse-truncate, shortly spreading-pubescent, with a ring of straight, erect hairs 5–7 mm long around the base. Bracts slightly enlarged about fruit.

Distribution: central west coast of Western Australia: restricted to the southern end of Nanga Station just south of Shark Bay, where it is relatively frequent.

Other collection: type locality, 26 August 1969, A. S. George s.n. (PERTH).

This unusual species differs from all others of the genus especially in the form of its leaves. The only other species with pungent leaves is *A. pungens* Meisn., in which they are divided into terete lobes. *A. acanthophyllus* is much more robust than other species. Its stems are thick and woody, and the whole shrub is quite dense. Only *A. cygnorum* Diels approaches it in the size of the stems, but it is more brittle than the new species.

Geographically the species is of great interest. It is the only *Adenanthos* in the area, the nearest locality for another species being *A. cygnorum* near the Murchison River nearly 150 km to the south. It is also at the northern extremity of the South-West Botanical Province, in an area where several other taxa of unusual interest occur, e.g. *Newcastelia chrysophylla* C. A. Gardn., *Grevillea rogersoniana* C. A. Gardn. and *Eucalyptus roycei* Carr, Carr & George. These and many other species form a tall, fairly dense shrubland on deep red sand.

Adenanthos ellipticus A. S. George sp. nov.

Derivation of epithet: Latin, *ellipticus*, elliptic, in reference to the leaf-shape.

Frutex patenti-ramosus ad 5 m altus sine lignotubero. *Folia* elliptica, 2–5 cm longa, 5–15 mm lata, plana vel concava, plerumque integra, raro apicibus 2–3-lobatis, 3–5-nervata, breviter tomentosa, in petiolis 4–10 mm longis angustata.

Flores axillares in pedunculos graciles 6–12 mm longos. *Bractae* ovato-ellipticae, obtusae, 1–3 longae, pubescentes. *Perianthium* 22–23 mm longum limbo 4 mm longo, intus glabrum praeter pilos longos post antheram, extus pilis brevibus rectis pubescens. *Antherae* sessiles 2.5 mm longae. *Squamae hypogynae* late obtusae, 1.3 mm longae, ad basin cum perianthio connatae. *Ovarium* doliiforme, breviter pubescens. *Stylus* 29–32 mm longus, dimidio supero patenti-hirsuto, ad basin glaber. *Fructus* doliiformis 6–7 mm longus, parce pubescens.

Type: East Mt. Barren, Fitzgerald River National Park, Western Australia—33°55'S, 120°02'E, 9 September 1971, A. S. George 10968. *Holo*: PERTH, iso: CANB, K, NSW, PERTH. In spirit collection at PERTH, no. 1310/A.

A rather slender, openly-branched *shrub* to 5 m high, without a lignotuber. *Branchlets* erect or ascending, terete, tomentose with somewhat curled, simple hairs, wearing off after several years leaving a smooth bark. *Leaves* elliptic, mostly 2–5 cm long, 5–15 mm wide (sometimes smaller and narrower); flat, (or concave in upper half), entire or with 2–3 obtuse lobes 1–4 mm long; 3–5-nerved (when 5 the inner 3 more prominent); shortly tomentose with curled, \pm appressed hairs; contracted into a terete petiole 4–10 mm long; a small, yellow-orange gland at the apex of the leaf or each lobe.

Flowers dull pink or red, axillary on slender peduncles 6–12 mm long bearing a few scattered bracts up to 1 mm long. *Involucral bracts* about 4, ovate-elliptic, obtuse, 1–3 mm long. *Perianth* 22–23 mm long at anthesis, including the limb of 4 mm long; tube contracted about ovary, broader above, then narrowed towards the limb; pubescent outside with short, straight hairs, glabrous inside except for silky hairs behind the anther. *Anthers* sessile, 2.5 mm long, connective produced into a short acute appendage. *Hypogynous scales* oblong, obtuse, thin, 1.3 mm long, fused to perianth in lower 3/4. *Ovary* doliform, shortly pubescent, the long basal hairs slightly exceeding the ovary. *Style* 29–32 mm long, upper half spreading-hirsute, becoming glabrous below. *Pollen presenter* 2.5 mm long, with a slight kink just above the base, almost terete but the apex laterally compressed, obtuse; stigmatic groove terminal.

Fruit 6–7 mm long, doliform with the truncate apex slightly curved postically; irregularly ribbed, sparsely pubescent, with long hairs about base. Bracts enlarged to 9 mm long in fruit.

Distribution: south coast of Western Australia: restricted to East Mt. Barren in the Fitzgerald River National Park.

Other collections: all from East Mt. Barren: 26 Nov. 1931, W. E. Blackall 1433 and C. A. Gardner 2953; 31 Jan. 1960, A. S. George 572; 21 April 1962, A. S. George 3677A; 25 Oct. 1964, K. Newbey 1451. (All at PERTH)

The affinity of this species is with *Adenanthos cuneatus* Labill. and *A. stictus* A. S. George (see below). However, its leaves are usually entire, have more prominent nervation and lack the small glands over the surface. It is much taller than *A. cuneatus* and lacks a lignotuber which the latter has. *A. cuneatus* also occurs around East Mt. Barren, but the two taxa remain distinct. *A. stictus* is far removed geographically, and though of tall habit is more densely branched than *A. ellipticus*.

Adenanthos gracilipes A. S. George sp. nov.

Derivation of epithet: Latin, *gracilis*, slender, and *pes*, a foot, in reference to the slender peduncles.

Frutex effusus ad 70 cm altus. *Folia* sessilia, 1–3 cm longa, in lobis 3 teretibus angustis divisa, lobi etiam 1-furcati, obtusi, cum glande miniata sub apice, parce appresso-pubescentes, deinde glabra; lobis 3–15 mm longis.

Flores axillares solitarii in pedunculos graciles glabros 3–5 mm longus. *Bractae* ovatae, obtusae, exteriores 1–2 mm, interiores 3–4 mm longae, fere glabrae marginibus ciliatis, cum glande miniata sub apice. *Perianthium* 17–18 mm longum; unguis fere aequales, extus dense pubescentes, intus glabri; limbus 2.5 mm longus, obtusus, cum glande miniata sub apice, extus dense pubescens, intus cum pilis longis post antherum. *Antherae* 1.5 mm longae, connectivis breviter productis, acutis. *Squamae hypogynae* anguste-ovatae, obtusae, minus quam 1 mm longae. *Ovarium* breviter hirsutum. *Stylus* glaber, 23–25 mm longus.

Type: ± 65 km E of Lake King, on Norseman road, Western Australia, 15 Sept. 1964, R. H. Kuchel 1817. *Holo*: PERTH, *iso*: AD, CANB, PERTH.

A low, spreading shrub to 70 cm tall. *Branchlets* terete, reddish, appressed-pubescent, becoming glabrous after 1–2 years. *Leaves* sessile, 1–3 cm long, divided for 1/3–1/2-way into 3 terete, narrow lobes, these again often once-forked; lobes 3–15 mm long, obtuse but with a small orange-red gland below the somewhat uncinat apex; sparsely appressed-pubescent with straight hairs, becoming glabrous but slightly scabrous.

Flowers pink, solitary in the woolly, uppermost axils, on slender, glabrous peduncles 3–5 mm long, the latter sometimes bearing a few minute bracts. *Involucral bracts* ovate, obtuse, outer ones 1–2 mm long, the inner 3–4 mm, with a small red gland below the apex, almost glabrous on both sides, the margins ciliate. *Perianth* 17–18 mm long, the tube swollen about the ovary, narrowed above; claws almost equal, densely pubescent outside with straight, spreading hairs, glabrous within; limb 2.5 mm long, obtuse, with an oval red gland on the outer surface below the apex, densely pubescent outside, with a few long hairs inside behind the anthers. *Anthers* 1.5 mm long, the connective shortly produced, acute. *Hypogynous scales* narrow-ovate, obtuse, less than 1 mm long, shortly united to the perianth. *Ovary* shortly hirsute. *Style* glabrous, somewhat flattened in upper half, 23–25 mm long; *pollen presenter* 1.5 mm long, ± terete but finely ribbed, the apex laterally compressed. *Fruit* not seen.

Distribution: South-Western Australia, between Hyden, Lake King and Salmon Gums.

Other collections: 300 mile peg, Hyden-Norseman road, 26 Oct. 1964, J. S. Beard 3853; sine loc., 1965, W. Rogerson 239. (All at PERTH)

Adenanthos gracilipes is a distinctive species with no very close allies. It is somewhat similar to *A. apiculatus* R.Br. which is of similar, sprawling habit. However the new species has leaves with thicker, widely-spreading lobes, the peduncles are longer and very slender, and the involucre and flowers are different in many respects.

Adenanthos stictus A. S. George sp. nov.

Derivation of epithet: Greek, *stictos*, spotted, in reference to the small glands over the leaf surface.

Frutex ramosus ad 5 m altus ramis erectis vel ascendentibus. *Folia* anguste-ad late-cuneata, 15–35 mm longa petiolis 5–10 mm longis, apicibus 3–6-lobatis lobis triangularibus obtusis 1–4 mm longis cum glandibus miniatis terminalibus; lamina plana, 5–25 mm lata, utrinque sericeo-pubescent tandem ± glabra, glandibus atro-rufis multis punctata.

Flores axillares, solitarii, in pedunculos 7–13 mm longos. *Bractae* ovatae 1–3 mm longae, sericeo-pubescentes, obtusae vel acutae, apicibus canaliculatis cum glandibus parvis. *Perianthium* 23–25 mm longum extus hirsutum intus glabrum, limbo 3–4 mm longo sub-acuto extus hirsuto intus post antheram sericeo. *Antherae* 3 mm longae connectivis brevibus. *Squamae hypogynae* cum perianthio pro 1.5 mm connatae, partibus liberis minus quam 1 mm longis, obtuse-lobatis. *Ovarium* breviter pubescens. *Stylus* 29–32 mm longus, ad basin glaber, supra breviter hirsutus, deinde glaber.

Type: ± 8 miles N of Marchagee on Geraldton Hwy, Western Australia—29°57'S, 116°04'E, 10 Sept. 1970, A. S. George 10379. *Holo*: PERTH, *iso*: AD, CANB, K, MEL, NSW, PERTH, RSA.

A much-branched, grey-green *shrub* to 5 m tall without a lignotuber, the branches erect or ascending. *Branchlets* terete, densely pubescent with \pm straight, appressed hairs, becoming glabrous. *Leaves* narrowly to broadly cuneate, mostly 15–35 mm long including the petiole of 5–10 mm; apex 3–6-lobed, rarely entire when small or with up to 9 lobes in seedling leaves, the lobes triangular, often unequal, 1–4 mm long, obtuse, terminating in a yellow-orange gland; lamina flat, 5–25 mm wide, silky-pubescent on both sides, at length \pm glabrous, dotted with many small dark red-brown glands, more numerous on the upper surface; venation furcate.

Flowers dull reddish-pink and green, axillary, solitary, on slender peduncles 7–13 mm long, the latter bearing scattered, ovate bracts \pm 1 mm long. *Involucral bracts* 4–6, ovate, 1–3 mm long, silky-pubescent, obtuse to acute, the apex canaliculate and tipped with a small orange-yellow gland. *Perianth* 23–25 mm long (including the limb of 3–4 mm), hirsute outside with rather spreading hairs, glabrous inside, the limb sub-acute, hirsute outside, silky inside behind the anther. *Anther* sessile, 3 mm long, connective shortly produced. *Hypogynous scales* united to perianth for 1.5 mm, the free part less than 1 mm long, obtusely lobed. *Ovary* shortly pubescent, the long basal hairs 4 mm long. *Style* 29–32 mm long, glabrous at base, shortly hirsute above, but glabrous towards apex; *pollen presenter* 2–2.5 mm long, scarcely thicker than style apex. *Fruit* 7–8 mm long, oblong but narrowed at apex, obtuse, smooth, sparsely and shortly pubescent, with a basal ring of straight, erect hairs.

Distribution: South-Western Australia, between Coorow and Marchagee, growing in deep sand on heath.

Other collections: W of Coorow, 25 Sept. 1962, J. S. Beard 1933; Marchagee, 14 Jan. 1931, C. A. Gardner s.n.; Watheroo National Park, 7 Oct. 1971, R. D. Royce 9695. (All at PERTH)

Adenanthos stictus is the equivalent on the heaths north of Perth of *A. cuneatus* Labill. on the south coastal heaths. The two are closely related and possibly form a vicarious pair. *A. cuneatus* differs in having a lignotuber; in its lower habit; in the leaves having straight margins and shorter petioles, with a more silky and appressed indumentum which is also more persistent; in the very inconspicuous glands; in the denser, closer, indumentum of the flowers; and in the style having long, spreading hairs as well as short ones. The new growth of *A. cuneatus* is much more red than that of *A. stictus*, and the leaf surface does not wrinkle so much on drying.

***Adenanthos teges* A. S. George sp. nov.**

Derivation of epithet: Latin, *teges*, a mat, in reference to the habit.

Frutex prostratus ad 3 m latus, sine lignotubero. *Ramuli* teretes, dense tomentosi pilibus brevibus longisque crispatis. *Folia* variabilia, ternato-divisa lobis filiformibus teretibus, lateralibus plerumque etiam furcatis; ea ad basin surculorum 6–7 mm longa, supra majora, ad 20 mm longa; ea floralibus simpliciter ternata vel integra, 25–28 mm longa; omnia laxe appresso-hirsuta, cum glandibus miniatis sub apices loborum.

Flores terminales in pedunculos 1–2 mm longos. *Bractee* triangulari-ovatae, 1–3 mm longae, obtusae, pubescentes. *Perianthium* 15–16 mm longum, extus breviter hirsutum, intus glabrum, limbo acuto 2.5 mm longo extus hirsuto intus post antherum sericeo. *Antherae* 1.25 mm longae, connectivis prominenter et anguste productis. *Squamae hypogynae* ca. 1 mm longae, oblongae, obtusae. *Ovarium* breviter pubescens. *Stylus* 27–28 mm longus, glaber.

Type: North-east of Mundaring Weir, \pm 32 km east of Perth, Western Australia—31°57'S, 116°17'E, 7 December 1973, A. S. George 11758. *Holo*: PERTH, *iso*: PERTH, CANB, K, NSW. In spirit collection at PERTH, no. 1523/B.

A prostrate, mat-like *shrub* to 3 m broad, without a lignotuber, the branchlets rarely ascending, not rooting. *Branchlets* terete, densely tomentose with short and long curled hairs, the latter disappearing with age; when young with a few very long hairs also.



George, Alexander S. 1974. "Five new species of Adenanthos (Proteaceae) from Western Australia." *Nuytsia: journal of the Western Australian Herbarium* 1(4), 381–385. <https://doi.org/10.58828/nuy00026>.

View This Item Online: <https://www.biodiversitylibrary.org/item/223201>

DOI: <https://doi.org/10.58828/nuy00026>

Permalink: <https://www.biodiversitylibrary.org/partpdf/228642>

Holding Institution

Western Australian Herbarium

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.