Substrate selection and spatial segregation by two congeneric species of *Eustala* (Araneae: Araneidae) in southeastern Brazil

Hebert da Silva Souza¹, Yuri Fanchini Messas¹, Marcelo de Oliveira Gonzaga² and João Vasconcellos-Neto¹: ¹Departamento de Biologia Animal, Instituto de Biologia, Caixa Postal: 6109, Universidade de Campinas - UNICAMP, 13083-970, Campinas, SP, Brasil. E-mail: hssouza.bio@gmail.com; ²Instituto de Biologia, Universidade Federal de Uberlândia, 38400-902, Uberlândia, MG, Brasil

Abstract. Habitat and microhabitat selection by spiders are influenced by abiotic and biotic factors, including vegetation structure, natural enemies, and prey availability. Some species are highly dependent on particular conditions, such as the presence of substrates where they remain camouflaged, constantly humid sites or the occurrence of plants bearing glandular trichomes. Others are distributed in areas that include a wide range of physical conditions and interact with several types of prey, predators and competitors. In the present study, we evaluated spatial distribution and substrate selection of two sympatric congeneric species with distinct body shapes and colors, *Eustala taquara* (Keyserling 1892) and *E. sagana* (Keyserling 1893), in an area of Atlantic Forest in southeastern Brazil. We focused on the following factors regarding habitat selection: i) distance from the border (forest edge or interior); ii) altitudinal distribution, ranging from 740 to 1294 m; iii) web height above ground level; and iv) plant species used for web attachment. All individuals of both species were located at the forest edge, especially on dry branches. However, they occurred preferentially in different host plants *suaveolens* and *C. sumatrensis*. Dry branch preferences might be important to reduce species conspicuousness to visually oriented predators, such as birds and wasps. Spatial segregation between closely related species possibly minimizes interference interactions, such as competition for particular sites or prey items.

Keywords: Orb weaver spider, coexistence, altitudinal gradient, habitat selection, web building

Microhabitat selection during web construction has important implications for web-building spiders, affecting the frequency of prey interception (Heiling 1999; Herberstein & Fleisch 2003; Heiling et al. 2006), prey species availability (Herberstein 1997), interspecific competition and/or frequency of agonistic interactions with other spiders or organisms (Herberstein 1998; Nahas et al. 2012; Purcell et al. 2012). Microhabitat selection could also expose spiders to inappropriate environmental conditions (Gillespie 1987; Gonzaga et al. 2006), to predators (Gonzaga & Vasconcellos-Neto 2005) and to other sources of mortality, such as infections by fungi (Gonzaga et al. 2006).

The first requirements for web establishment are associated with site structural conditions, such as spatial distribution of branches used in web attachment, and an open space large enough to facilitate construction of the web (Stevenson & Dindal 1982; Rao & Poyyamoli 2001). Spiders, however, often have more specific criteria, and can use several cues to assess site quality before initiating web settlement. For example, Schuck-Paim & Alonso (2001) showed that *Nephilengys* cruentata (Fabricius 1775) (Nephilidae) used signs of previous conspecific presence to evaluate habitat quality. Conspecific silk inside the enclosures employed in the experiments increased settlement likelihood by *N. cruentata* individuals.

Plant characteristics can also be integral in determining initial settlement, establishment success, and permanence of several spider species (Romero & Vasconcellos-Neto 2005). Selectivity for particular host plants has been previously described for many spider species (Romero & Vasconcellos-Neto 2005, 2011; Romero et al. 2008; Hesselberg & Triana 2010). Spiders select plants with specific attributes, e.g., presence of glandular trichomes, adequate branch and leaf densities, pollination by potential prey, color pattern, and structural composition that can serve as suitable shelter, which consequently increases each species potential to improve prey capture success, and reduces the likelihood of capture by natural enemies.

The genus Eustala Simon 1895 (Araneidae), restricted to the Americas, is currently composed of 90 species (Platnick 2014). Individuals of several Eustala species are often distributed throughout shrub and tree vegetation (Poeta et al. 2010), and are the primary prey species of several hunting wasps (Gonzaga & Vasconcellos-Neto 2005; Buschini et al. 2008). In Serra do Japi, an area of Atlantic Forest situated in southeastern Brazil, three species have been previously collected (Indicatti et al. 2012): Eustala perfida Mello-Leitão 1947, which typically occurs in shaded areas of forest interior and constructs elongated orbs close to concavities on tree trunks (Messas et al. 2014); E. sagana (Keyserling 1893); and E. taquara (Keyserling 1892). Habitat selection and spatial distribution of E. perfida was previously studied by Messas et al. (2014), but there is no information available on the remaining species. We hypothesized that E. sagana and E. taquara, two species with distinct body shapes and colors, would exhibit differences in daytime resting substrates. In addition, we evaluated spatial segregation in these species by analyzing habitat selection based on vegetation height and altitude within the distribution area.

METHODS

Study site.—We conducted the present work in the Serra do Japi (23° 11' S, 46° 52' W), a forest reserve located near the city of Jundiaí, São Paulo state, Brazil. The climate is seasonal, with average monthly temperatures from 13.5° C

THE JOURNAL OF ARACHNOLOGY



Figure 1.—Schematic drawing of the methodology employed to analyze the frequency of available sites (vegetation height) in the Serra do Japi, Jundiaí-SP. Four 5-m lines, separated vertically every 0.5 m, were arranged in the vegetation; all plants touched by the lines were included.

in July to 20.3° C in January. The driest period is from June to September, and the rainy season is from December to February (Pinto 1992). The local vegetation is characterized by a semi-deciduous rainforest with canopy height varying between 10–15 m. The higher mountain regions support a semideciduous forest composed of thinner trees (Leitão-Filho 1992). The altitude in the Serra do Japi ranges from 740–1294 m.

Spatial distribution.—All sampling was conducted in March 2011. We examined the distance from the border (forest edge or interior) and the altitudinal distribution of both species. *Eustala taquara* and *E. sagana* altitudinal distribution and edge effects were investigated by sampling forest edge and interior in four altitudinal ranges: 750–800 m, 1000 m, 1250 m, and 1294 m. In this study, we considered the forest edge as a

Table 1.—List of available plant species present on sample plots and frequency and number of individuals of *Eustala sagana* found occupying plants along the edge of the forest in Serra do Japi, Jundiaí – SP, Brazil. The data were collected at an altitude of 800 m during May and July 2011.

Family	Species	Plants		E. sagana	
		п	%	п	%
Anacardiaceae	Lithraea molleoides	1	0.15	0	0
Asclepidadaceae	Asclepias curassavica L.	1	0.15	0	0
Asteraceae	Cyrtocymura scorpioides (Lam.) H. Rob.	125	18.71	0	0
	Eupatorium sp.	32	4.79	0	0
	Vernonanthura ferruginea (Less.) H. Rob.	31	4.64	9	8.18
	Vernonanthura phosphorica (Vell.) H. Rob.	17	2.54	0	0
	Trichogoniopsis adenantha (DC.) R.M.King & H.Rob	12	1.8	2	1.82
	Baccharis dracunculifolia DC.	4	0.6	1	0.91
	Bidens brasiliensis Sherff	4	0.6	3	2.73
	Calea pinnatifida (R. Br.) Less.	8	1.2	0	0
	Conyza bonariensis (L.) Cronq.	5	0.75	0	0
	Conyza sumatrensis (Retz.) E. Walker	15	2.25	35	31.82
	Erechtites valerianifolius (Link ex Spreng.) DC.	0	0	3	2.73
	Mikania sp.	5	0.75	0	0
	sp.1	66	9.88	0	0
	sp.2	4	0.6	0	0
	Acanthospermum hispidum	0	0	1	0.91
	Tagetes minuta L.	0	0	1	0.91
Bignoniaceae	Pirostegia venusta (Ker-Gawl) Miers	2	0.3	0	0
Cecropiaceae	Cecropia sp.	2	0.3	0	0
Euphorbiaceae	Croton floribundus Spreng	1	0.15	0	0
Fabaceae	sp.1	6	0.9	0	0
Lamiaceae	Hyptis suaveolens L.	47	7.04	26	23.64
	Hyptis sp.1	13	1.95	3	2.73
Malvaceae	Triunfetta semitriloba Jacq.	36	5.39	1	0.91
	sp. 1	0	0	3	2.73
	sp. 2	1	0.15	0	0
Melastomataceae	sp. 1	9	1.35	0	0
Piperaceae	Piner sp	2	0.3	0	0
Poaceae	Brachiaria decumbens Stapf.	168	25.15	0	0
	Panicum maximum Jacq	21	3.14	0	0
	Grass	0	0	9	8.18
Pteridophytes		5	0.75	0	0
Rosaceae	Rubus rosifolius BP MS	4	0.6	0	0
Rubiaceae	sp 1	2	0.3	0	0
Solanaceae	Solanum conceinum Schott ex Sendtn.	10	1.5	0	0
	Solanum variabile Mart	3	0.45	0	0
	Sessea brasiliensis Toledo	2	0.3	0	0
Verbenaceae	Lantana camara I	4	0.6	0	0
Others	Rare species	0	0	10	9.09
C there's	Dry branches	0	0	3	2.73
Total		0668	100	110	100

Table 2.—List of available plant species present on sample plots, and frequency and number of individuals of *Eustala taquara* found occupying plants along the edge of the forest in Serra do Japi, Jundiaí – SP, Brazil. The data were collected at an altitude of 1000 m in March 2011.

Family	Species	Plants		E. taquara	
		n	%	п	%
Asteraceae	Ageratum conyzoides L.	1	0.54	0	0.00
	Ambrosia artemisiifolia L.	11	5.95	4	2.65
	Bidens brasiliensis Sherff	2	1.08	3	1.99
	Calea pinnatifida (R. Br.) Less.	2	1.08	0	0.00
	Conyza bonariensis (L.) Crong.	14	7.57	103	68.21
	Cyrtocymura scorpioides (Lam.) H. Rob.	4	2.16	0	0.00
	Erechtites valerianifolius (Link ex Spreng.) DC.	1	0.54	2	1.32
	Eupatorium sp.	2	1.08	0	0.00
	Trichogoniopsis adenantha (DC.) R.M.King & H.Rob	0	0.00	2	1.32
Euphorbiaceae	Croton floribundus Spreng	8	4.32	0	0.00
	Croton urucurana Baill	5	2.70	0	0.00
Fabaceae	Acacia plumosa Lowe	1	0.54	0	0
	Desmodium discolor Vogel	0	0.00	3	1.99
Lamiaceae	Hyptis suaveolens L.	49	26.49	10	6.62
Malvaceae	sp.1	24	12.97	0	0.00
Melastomataceae	Miconia sp.	2	1.08	0	0.00
	sp.1	2	1.08	0	0.00
Myrtaceae	sp.1	0	0.00	2	1.32
Piperaceae	sp.1	8	4.32	0	0.00
Poaceae	Grass – sp.1	2	1.08	0	0.00
	Unidentified	5	2.70	0	0.00
Rosaceae	Rubus rosifolius BP. MS.	2	1.08	0	0.00
Rubiaceae	sp.1	3	1.62	5	3.31
Solanaceae	Solanum concinnum Schott ex Sendtn	2	1.08	0	0.00
	Solanum variabile Mart	2	1.08	0	0.00
	Sesseia brasiliensis Toledo	2	1.08	0	0.00
Verbenaceae	Lantana camara L.	2	1.08	0	0.00
Others	Pteridophytes	13	7.03	0	0.00
	Eudicotyledons	1	0.54	2	1.32
	Dry branches	15	8.11	13	8.61
Total		185	100	151	100

zone of 2 m from the road margin towards the inner forest. This area is covered by herbaceous plants and shrubs located between the road and arboreal vegetation. For each site, individuals from 20×2 m plots (n = 15) were sampled at the forest edge, and individuals from 10×10 m plots (n = 12) were sampled within the forest. Shape and size of the plots in the forest edge were restricted by the area available under the influence of the wide open trails and roads. We sprayed water in vegetation to locate webs during the searching procedure.

We compared the number of *E. taquara* and *E. sagana* individuals at distinct altitudes using a non-parametric Kruskal-Wallis test, as initial analyses showed the data distribution lacked homogeneity of variances. Only plots located at the edge of trails were considered as replicates because both species were absent from plots located in the forest interior. Post-hoc Dunn's multiple comparison tests (Dunn 1964) were applied to compare the abundances between altitudes.

Substrate selection.—We searched for spiders along the forest edge in order to obtain the observed values of web height from ground level, green and dead substrate and plant species occupied. Samples were collected where each species was more abundant, at altitudes of 800 m for *E. sagana* during May and July 2011, and at 1000 m for *E. taquara* in March 2011.

We evaluated whether *E. sagana* and *E. taquara* exhibited specific height preferences (above ground level) by establishing 5 m transects (n = 8) in the two altitudes with higher occurrence of each species. We extended lines at four different heights (0.5 m; 1.0 m; 1.5 m; and 2.0 m), and counted the number of times each line intersected vegetation (Fig. 1). The total number of intersections was used to establish the availability of sites at different sample heights. We used the number of intersections at each height as the expected frequency for spider occupation. These frequencies were compared with the frequencies of observed web heights used by *E. taquara* and *E. sagana* using a *G* test.

Substrate selection for web construction in *E. taquara* and *E. sagana* was investigated in 10×2 m plots (n = 8) also on the forest edge. Within these plots we calculated the number of sites available with green and dry branches (open spaces wide enough to allow web construction) and the proportion of each plant species (Table 1 and 2). We compared the available number of green and dead branches with the number of these kinds of branches occupied by *Eustala* spiders. We also compared the relative abundance of plant species in plots (expected frequencies), at the altitude with highest abundance of each spider species, with the relative abundance of plants used as web-building sites along the forest edge (observed frequencies) applying *G* test.

THE JOURNAL OF ARACHNOLOGY



Figure 2.—Eustala taquara specimens resting on dead (dry) vegetation, with coloration pattern details. Scale bar: 0.5 cm.



Figure 3.—*Eustala sagana* specimens resting on dry vegetation. Detail of a free sector of the web which is lacking capture spirals. The arrow indicates the signal thread that connects the spider to the web center. Scale bar: 1.0 cm.

RESULTS

Results showed both *Eustala* species did not build shelters, but were instead found in shrub branches, which provided attachment points for web construction. Spiders remained in contact with branches, holding a web thread connected to the hub of the orb web. The spider color patterns were very similar to the substrate patterns used as resting positions (based on the human visual system) (Figs. 2, 3).

Spatial distribution.—All *Eustala taquara* (n = 97) and *E*. sagana (n = 46) individuals were located at the forest edge. Both Eustala species were sampled at all altitudes of Serra do Japi, but with different density patterns. A significantly higher abundance of E. sagana was detected in the lower Serra do Japi area, between 750-850 m (91.3% of all sampled individuals) (Kruskal-Wallis: H = 23.08, DF = 3, P < 0.0001; Dunn $Z_{800m} \times$ $I_{1000m} = 3.73, P = 0.001; Z_{800m \times 1250m} = 3.99, P = 0.0004; Z_{800m \times 1294m} = 3.99, P = 0.0004; Z_{1000m \times 1250m} = 0.27, P = 0.0004; Z_{1000m \times 1250m} = 0.27, P = 0.0004; Z_{1000m \times 1250m} = 0.0004; P = 0.0004; P$ 1.0; $Z_{1000m \times 1294m} = 0.27$, P = 1.0; $Z_{1250m \times 1294m} = 0.00$, P =1.0; Fig. 4A). We found 20.6% of Eustala taquara individuals at 750 m; 52.6% at 1000 m; 16.5% at 1250 m; and 10.3% at 1294 m of altitude, with a significantly higher abundance at the intermediate zone (1000 m) (Kruskal-Wallis: H = 19.24; DF = 3; P < 0.001; Dunn $Z_{800m \times 1000m}$ = 3.27, P = 0.006; $Z_{800m \times 1000m}$ $_{1250m} = 0.005, P = 1.0; Z_{800m \times 1294m} = 0.66, P = 1.0; Z_{1000m \times 1294m} = 0.006, P = 0.0$ $_{1250m} = 3.26, P = 0.007; Z_{1000m \times 1294m} = 3.93, P = 0.0005;$ $Z_{1250m \times 1294m} = 0.66, P = 1.0;$ Fig. 4B).

Substrate selection.—Lines intersected vegetation more frequently at the 0.5 m level above the ground, and decreased gradually as heights increased to 2.0 m. However, results

showed *E. sagana* occurrence was highest at 1.0 m (30.6%) and 1.5 m (46.4%) above ground level (G = 115.29; P < 0.0001; n = 117 sites; n = 106 spiders; Fig. 5A). *Eustala taquara* frequencies were significantly higher at 1.0 m (39.6%) and 1.5 m (44.3%) compared to the expected frequencies considering the available sites (G = 97.80; P < 0.0001; N sites = 188; N spiders = 108; Fig. 5B).

Almost all *E. sagana* individuals (96.9%) were observed using dead vegetation branches (Fig. 6A), indicating a preferential use for these sites (G = 197.16; P < 0.0001; Nsites = 596; N spiders = 96). *Eustala taquara* exhibited the same pattern, with most individuals (96.6%) also sampled on dead branches (Fig. 6B), indicating preferred use of these sites (G = 167.27; P < 0.0001; N sites = 315; N spiders = 146).

Host plants.—*Eustala sagana* occurred preferentially on *Hyptis suaveolens* (L.) Poit (Lamiaceae), *Conyza sumatrensis* (Retz.) E. Walker (Asteraceae), and *Vernonanthura ferruginea* (Less.) H. Rob. (Asteraceae) (G = 278.5; P < 0.0001; Fig. 7A; Table 1), while *E. taquara* occurred mainly on *C. bonariensis* (L.) Cronquist (G = 107.70; P < 0.0001, Fig. 7B; Table 2).

DISCUSSION

Both species exhibited similarities in habitat selection, and showed preferential use and restriction to forest borders, dead vegetation branches, and heights between 1.0 and 1.5 m above ground level. In contrast, results showed each species was distributed preferentially at a specific altitude, and on distinct plant species. These results are consistent with our hypothesis that the species are spatially segregated.



Figure 4.—*Eustala sagana* and *E. taquara* densities at different altitudes of the Serra do Japi, Jundiaí-SP. Box plots show median and interquartiles. Whiskers represent $1.5 \times$ interquartile ranges.

Eustala taquara occurred at a higher abundance at 1000 m, while *E. sagana* exhibited higher abundance at lower altitudes (750–850 m). Preferences regarding altitude might be a response to requirements for specific abiotic conditions (temperature, humidity, exposure to solar radiation), or to biotic criteria, including obtainable prey, exposure to predators, or availability of suitable web construction sites (Turnbull 1973; Brown 1981; Janetos 1986; Lubin et al. 1991; Marshall & Rypstra 1999). Similarly, Purcell & Avilés (2007) observed that altitude is an important factor affecting *Anelosimus* Simon 1891 (Theridiidae) species distribution and colony size (within social species of this genus) in Ecuador. The same authors suggested the influence of altitude occurred primarily in response to biotic factors, such as prey size and predator pressure; large insects were more common in



Figure 5.—Available (vegetation height) and observed sites used by *E. sagana* and *E. taquara* in Serra do Japi, Jundiaí-SP.

lowland rain forests, and predator pressures were reduced at higher altitudes. In addition, it is possible that spatial segregation between closely related species is a response to competition (but see Wise (1993) for evidence against competition in spiders). In a future investigations we intend to examine prey availability (types and sizes) for each species, web architecture variation, and possible variation in abiotic conditions along the altitudinal gradient to evaluate the causes of spatial segregation. It is also important to evaluate possible differences in the abundance of plant species along the gradient as a component of habitat selection.

Eustala taquara individuals were associated with herbaceous plants and shrubs that occurred on the forest edge, primarily *C. bonariensis.* This species apparently avoids constructing webs in *H. suaveolens*, which is preferentially used by *E. sagana.* Respective associations of *E. illicita* (O. Pickard-Cambridge 1889) and *E. oblonga* Chickering 1955 with *Acacia collinsii* Safford and *A. melanoceras* Beurl. have previously been characterized (Chickering 1955; Hesselberg & Triana 2010), suggesting habitat selection by some *Eustala* species. Despite the association of *E. illicita* with *A. collinsii*, Hesselberg & Triana (2010) also observed these spiders occupying dead vegetation along the study site, and concluded the relationships between *Eustala* and vegetation species are not obligatory.

64



Figure 6.—Available green and dry (dead) vegetation sites and sites used by *Eustala sagana* and *E. taquara* in Serra do Japi, Jundiaí-SP.

The selection of specific sites for web construction by E. taquara and E. sagana may be important to potentially reduce their conspicuousness and, consequently, exposure to visually oriented predators (such as birds and hunting wasps). We found that individuals of both species were apparently cryptic when resting in dry vegetation during the day. We have no information on their predators in Serra do Japi, but several empirical studies have demonstrated that Eustala species are hunted by Trypoxylon Latreille 1796 (Crabronidae) wasps (Rehnberg 1987; Gonzaga & Vasconcellos-Neto 2005; Buschini et al. 2008, 2010). Eustala is the most common prey found in nests of Trypoxylon albonigrum Richards 1934, for example, in another Atlantic forest area in São Paulo state (Gonzaga & Vasconcellos-Neto 2005; Araújo & Gonzaga 2007). It is important now to evaluate the ability of birds and wasps to perceive the chromatic and achromatic contrasts between the spiders and the substrates selected for web construction.

To conclude, this study showed that two congeneric sympatric species of *Eustala* present some similarities regarding habitat characteristics, such as preferential distribution close to the forest borders and occurrence on dead vegetation branches. However, there were also important differences in the two species' distributions. Their spatial segregation at distinct altitudes might reduce interference interactions between species. Future studies on prey preferences and experimental procedures manipulating the density of each species are essential to understanding the possible impact of each species on the abundance and habitat selection of the other.



Figure 7.—Available plant species and plants occupied by *Eustala* sagana and *E. taquara* in Serra do Japi, Jundiaí-SP.

ACKNOWLEDGMENTS

This project was supported by Instituto Nacional de Ciência e Tecnologia dos Hymenoptera Parasitoides da Região Sudeste (http://www.hympar.ufscar.br) and Fundação de Amparo à Pesquisa do Estado de São Paulo (processes: 2011/02535-3, 2011/15457-0, 2011/15456-4 and 2011/02537-6). We also thank Antonio Domingos Brescovit and Maria Rita M. Poeta for the identification of the spiders, and we appreciate the valuable suggestions of the reviewers.

LITERATURE CITED

- Araújo, M.S. & M.O. Gonzaga. 2007. Individual specialization in the hunting wasp *Trypoxylon (Trypargilum) albonigrum* (Hymenoptera, Crabronidae). Behavioral Ecology and Sociobiology 61:1855–1863.
- Brown, K.M. 1981. Foraging ecology and niche partitioning in orbweaving spiders. Oecologia 50:380-385.
- Buschini, M.L.T., N.A. Borba & A.D. Brescovit. 2008. Patterns of prey selection of *Trypoxylon (Trypargilum) lactitarse* Saussure (Hymenoptera: Crabronidae) in southern Brazil. Brazilian Journal of Biology 68:519–528.
- Buschini, M.L.T., T.R. Caldas, N.A. Borba & A.D. Brescovit. 2010. Spiders used as prey by the hunting wasp *Trypoxylon (Trypargilum) agamemnon* Richards (Hymenoptera: Crabronidae). Zoological Studies 49:169–175.
- Chickering, A.M. 1955. The genus *Eustala* (Araneae, Argiopidae) in Central America. Bulletin of the Museum of Comparative Zoology 112:391–518.
- Dunn, O.J. 1964. Multiple comparisons using rank sums. Technometrics 6:241–252.
- Gillespie, R.G. 1987. The mechanism of habitat selection in the longjawed orb-weaving spider *Tetragnatha elongata* (Araneae, Tetragnathidae). Journal of Arachnology 15:81–90.
- Gonzaga, M.O. & J. Vasconcellos-Neto. 2005. Orb-web spiders (Araneae: Araneomorphae; Orbiculariae) captured by huntingwasps (Hymenoptera: Sphecidae) in an area of Atlantic Forest in south-eastern Brazil. Journal of Natural History 39:2913–2933.

- Gonzaga, M.O., N.O. Leiner & A.J. Santos. 2006. On the sticky cobwebs of two theridiid spiders (Araneae: Theridiidae). Journal of Natural History 40:293–306.
- Heiling, A.M. 1999. Why do nocturnal orb-web spiders (Araneidae) search for light? Behavioral Ecology and Sociobiology 46:43–49.
- Heiling, A.M., K. Cheng & M.E. Herberstein. 2006. Picking the right spot: crab spiders position themselves on flowers to maximize prey attraction. Behaviour 143:957–968.
- Herberstein, M.E. 1997. Implications of microhabitat selection on prey capture for the web spider *Neriene radiata* (Walckenaer) (Araneae: Linyphiidae). Pp. 197–202. *In* Proceedings of the 17th European Colloquium of Arachnology. (P.A. Selden, ed.). British Arachnological Society, Edinburgh.
- Herberstein, M.E. 1998. Web placement in sympatric linyphiid spiders (Arachnida, Araneae): individual foraging decisions reveal inter-specific competition. Acta Oecologica 19:67–71.
- Herberstein, M.E. & A.F. Fleisch. 2003. Effect of abiotic factors on the foraging strategy of the orb-web spider *Argiope keyserlingi* (Araneae: Araneidae). Austral Ecology 28:622–628.
- Hesselberg, T. & E. Triana. 2010. The web of the acacia orb-spider *Eustala illicita* (Araneae: Araneidae) with notes on its natural history. Journal of Arachnology 38:21–26.
- Indicatti, R.P., A.D. Brescovit & J. Vasconcellos-Neto. 2012. Aranhas (Arachnida, Araneae) da Serra do Japi, Jundiaí-SP. Pp. 273–297. *In* Novos Olhares, Novos Saberes Sobre a Serra do Japi: Ecos de sua biodiversidade. (J. Vasconcellos-Neto, P.R. Polli & A.M. Penteado-Dias, eds.). Editora CRV, Curitiba, Brazil.
- Janetos, A.C. 1986. Web-site selection: are we asking the right questions? Pp. 9–48. *In* Spiders: webs, behavior, and evolution. (W.A. Shear, ed.). Stanford University Press, Stanford, California.
- Leitão-Filho, H.F. 1992. A flora arbórea da Serra do Japi. Pp. 40–63. *In* História natural da Serra do Japi: ecologia e preservação de uma área florestal no Sudeste do Brasil. (L.P.C. Morellato, ed.). Editora da Unicamp/Fapesp, Campinas, Brazil.
- Lubin, Y., M. Kotzman & S. Ellner. 1991. Ontogenetic and seasonal changes in webs and websites of a desert widow spider. Journal of Arachnology 19:40–48.
- Marshall, S. & A. Rypstra. 1999. Spider competition in structurallysimple ecosystems. Journal of Arachnology 27:343–351.
- Messas, Y.F., H.S. Souza, M.O. Gonzaga & J. Vasconcellos-Neto. 2014. Spatial distribution and substrate selection by the orb-weaver spider *Eustala perfida* Mello-Leitão, 1947 (Araneae: Araneidae). Journal of Natural History. DOI:10.1080/00222933.2014.909067.
- Nahas, L., M.O. Gonzaga & K. Del-Claro. 2012. Emergent impacts of ants and spider interactions: herbivory reduction in a tropical savanna tree. Biotropica 44:498–505.
- Pinto, H.S. 1992. Clima na Serra do Japi. Pp. 30-39. In História Natural da Serra do Japi: Ecologia e Preservação de uma Área

Florestal no Sudeste do Brasil. (L.P.C. Morellato, ed.). Editora da Unicamp/Fapesp, Campinas, Brazil.

- Platnick, N.I. 2014. The world spider catalog, version 14.5. American Museum of Natural History, Online at http://research.amnh.org/ entomology/spiders/catalog/index.html DOI: 10.5531/db.iz.0001.
- Poeta, M.R.M., M.A.L. Marques & E.H. Buckup. 2010. O gênero Eustala (Araneae, Araneidae) no sul do Brasil: duas espécies novas, descrições complementares e novas ocorrências. Iheringia, Série Zoologia 100:151–161.
- Purcell, J. & L. Avilés. 2007. Smaller colonies and more solitary living mark higher altitude populations of a social spider. Journal of Animal Ecology 76:590–597.
- Purcell, J., J. Vasconcellos-Neto, M.O. Gonzaga, J.A. Fletcher & L. Avilés. 2012. Spatio-temporal differentiation and sociality in spiders. PLoS ONE 7, e34592. DOI:10.1371/journal.pone.0034592.
- Rao, D. & G. Poyyamoli. 2001. Role of structural requirements in web-site selection in *Cyrtophora cicatrosa* Stoliczka (Araneae: Araneidae). Current Science 81:678–680.
- Rehnberg, B.G. 1987. Selection of spider prey by *Trypoxylon politum* (Say) (Hymenoptera: Sphecidae). Canadian Entomologist 119: 189–194.
- Romero, G.Q. & J. Vasconcellos-Neto. 2005. The effects of plant structure on the spatial and microspatial distribution of a bromeliad-living jumping spider (Salticidae). Journal of Animal Ecology 74:12–21.
- Romero, G.Q. & J. Vasconcellos-Neto. 2011. Interações entre aranhas e plantas: associações específicas e mutualismos. Pp. 241–258. In Ecologia de Interações Plantas-Animais: uma abordagem ecológico-evolutiva. (K. Del-Claro & H.M. Torezan-Silingardi, eds.). Technical Books Editora, Rio de Janeiro, Brazil.
- Romero, G.Q., J.C. Souza & J. Vasconcellos-Neto. 2008. Antiherbivore protection by mutualistic spiders and the role of plant glandular trichomes. Ecology 89:3105–3115.
- Schuck-Paim, C. & W.J. Alonso. 2001. Deciding where to settle: conspecific attraction and web site selection in the orb-web spider *Nephilengys cruentata*. Animal Behaviour 62:1007–1012.
- Stevenson, B.G. & D.L. Dindal. 1982. Effect of leaf shape on forest litter spiders: community organization and microhabitat selection on immature *Enoplognatha ovata* (Clerk) (Theridiidae). Journal of Arachnology 10:165–178.
- Turnbull, A.L. 1973. Ecology of true spiders. Annual Review of Entomology 18:305–348.
- Wise, D.H. 1993. Failure of the competitionist paradigm. Pp. 57–96. In Spiders in ecological webs. (D.H. Wise, ed.). First Edition. Cambridge University Press, New York.

Manuscript received 18 April 2014, revised 24 September 2014.

66



Souza, Hebert da Silva et al. 2015. "Substrate selection and spatial segregation by two congeneric species of Eustala (Araneae: Araneidae) in southeastern Brazil." *The Journal of arachnology* 43(1), 59–66. <u>https://doi.org/10.1636/j14-23.1</u>.

View This Item Online: https://doi.org/10.1636/j14-23.1 Permalink: https://www.biodiversitylibrary.org/partpdf/229485

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: American Arachnological Society License: <u>https://creativecommons.org/licenses/by-nc-sa/4.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.