

The Pan-Pacific Entomologist

Vol. XXXVIII

October, 1962

No. 4

BIOLOGICAL AND TAXONOMIC NOTES ON TWO CALIFORNIA SPECIES OF PROTEOTERAS

(Lepidoptera: Tortricidae)

JERRY A. POWELL

University of California, Berkeley

The genus *Proteoteras* includes eight described species in North America, six of which are restricted to Canada and the northeastern United States. Only one, the widespread *P. aesculana* Riley, has been listed in California (Heinrich, 1923). The larvae of the several species for which biological information is available are borers and gall-makers in the seeds and twigs of maples, box-elder, and horse chestnut (Heinrich, 1923; MacKay, 1959; Peterson, 1958; Schaffner, 1950).

PROTEOTERAS ARIZONAE Kearfott

Proteoteras arizonae Kearfott, 1907, Trans. Amer. Ent. Soc., 33:48.

The type locality of this species is Prescott, Arizona, and there were cotypes from Colorado. Heinrich (1923) figured the male genitalia of a specimen from Mesilla, New Mexico, and described a closely related species, *P. obnigrana*, from New Hampshire which is distinguished by wing color and minor genitalic differences, primarily in the shape of the valvae. Recently collected California specimens of this complex have the genitalia form of *obnigrana* but otherwise match the description and worn lectotype male of *arizonae* in the American Museum of Natural History. The olivaceous tinge of the forewing, characteristic of *obnigrana*, is not represented in any western specimens I have seen. A series in the U.S. National Museum from Eureka, Utah compares well with the type of *arizonae* and has genitalic characters (slide No. 8, C.H., 18 Oct., '24) of California specimens. It seems apparent that Heinrich's figure (fig. 300) does not represent *P. arizonae* well; and if, as Heinrich stated, "the differences (between fig. 300 and *P. obnigrana*, fig. 305) are more than could be looked for in one species," the southern New Mexico specimen may represent an undescribed species.

The California specimens of *P. arizonae* are very similar genitally to *P. obnigrana* but appear to differ by the more well developed "shoulders" of the tegumen and by having larger lateral

spines of the valva. The number of these spines may vary from three to five on the same specimen, or one or more may be greatly reduced, but the size of the well developed three may be diagnostic.

California material examined.—Sonoma Co.: Santa Rosa, 1 ♂ VIII-1-38 (E. C. Johnston). Contra Costa Co.: Walnut Creek, 1 ♀ VI-15-61, 1 ♂ VIII-8-61, at lights (J. Powell). San Mateo Co.: Redwood City, 1 ♂ VI-23-59, r. f. *Acer negundo californicum*, emgd. VII-13-59, 2 ♂ same data except VI-30-59, emgd. VII-24, 30-59 (JAP-59F2) (A. E. Pritchard).

Biology.—Specimens collected as larvae in San Mateo County during June 1959 were reared from *Acer negundo californicum*. Host plants have not been recorded in other areas. The larvae tunnel into the terminal twigs but apparently do not cause any gall-like swelling as does *P. willingana* (Kearfott) on boxelder (Peterson, 1958). Feeding begins at what appears to be the base of the current season's growth, ultimately causing all the twig and leaves beyond this point to die. Working downward, larvae burrow out all contents of the hard, green stems; fully developed larval chambers measured 25 to 35 mm. in length and usually tapered somewhat downward. Apparently most of the frass is ejected from the shelter, as only a few particles were found at the lower end of mature tunnels. A small amount also adheres to webbing which closes the entrance area and is visible from the exterior.

Pupation probably normally takes place within the shelter, since two pupae were found *in situ*, one of which had pupated prior to having been collected. However, in the laboratory two individuals left the stems and pupated between folds of the paper at the bottom of the container; and the larva of *P. willingana* leaves the feeding chamber and drops to the leaf duff to pupate according to Peterson (1958). Prior to pupation *P. arizonae* spins several fine silken partitions at varying intervals along the length of the burrow. The cocoon consists of a fine silk structure or lining of the apical area of the chamber, the pupal head being adjacent to the exit spot.

The flight records suggest two generations during a season in the San Francisco Bay area.

About one-half the larvae from the Redwood City rearing lot (JAP-59F2) were parasitized by the Ichneumonid, *Apistephialtes nucicola* (Cushman)¹. A male and four females were reared, emerging July 7 to 13. Townes and Townes (1960) record this

¹ Determined by G. S. Walley, Entomology Research Institute, Ottawa.

PROTEOTERAS AESCULANA Riley

Proteoteras aesculana Riley, 1881, Trans. St. Louis Acad. Sci., 4:321.

This widespread species was recorded from California by Heinrich (1923) without detailed data. It is commonly collected in the San Francisco Bay area, and the records suggest a multi-voltine life cycle. The rearing record given below involves silver maple, an introduced ornamental plant. A specimen in the U.S. National Museum from Corvallis, Oregon was reared from *Acer negundo*; and this host, as well as *Aesculus californicum* may be expected to be native foodplants in California.

California material examined.—Sonoma Co.: Santa Rosa, 1 ♂ V-12-36 (E. C. Johnston); Petaluma, 1 ♂ VI-26-37, 1 ♂ VIII-1-38 (E. C. Johnston). Napa Co.: Napa, 1 ♀ "Apr. 20" (Guedet). Contra Costa Co.: Richmond, 1 ♀ VI-15-59, 1 ♂ X-6-59 (C. D. MacNeill); El Cerrito, 1 ♀ IV-4-60, 2 ♀ VII-7, 19-60 (C. D. MacNeill), IX-27-60 (T. R. Haig); Walnut Creek, 1 ♂ VI-20-62 (J. Powell). Alameda Co.: Berkeley, 4 ♂, 1 ♀ III-10 to 17-59 (G. I. Stage), 1 ♀ III-9-59, 1 ♀ IV-1-59, 1 ♂ IX-25-59, 3 ♂, 2 ♀ III-4 to IV-7-60, 1 ♂ VII-20-60, 1 ♂ II-26-61, 1 ♂, 2 ♀ IV-3, 4-61, at light (J. Powell). San Mateo Co.: Belmont, 1 ♀ VI-30-59, r.f. *Acer dasycarpum* [= *saccharinum*] emgd. VII-27-59 (JAP-59F1) (A. E. Pritchard).

Biology.—Larvae collected at Belmont displayed a similar feeding behavior to that described for *P. arizonae*. However, the tunnels ranged up to 40-46 mm. in length. Pupation may normally occur outside the shelter, since a number of abandoned twig chambers were collected from the trees; and the one individual which was reared pupated on the bottom of the container.

A collection of young larvae (apparently 3rd and penultimate instars) was taken in mid July 1959 on *Acer negundo californicum* near Castro Valley, Alameda County, by A. E. Pritchard. These are presumed to be *P. aesculana*, although the smaller larvae lack the differentiated pinacula and the anal fork. In at least one case a small larva was found feeding in a tunnel obviously made by a mature larva. Such abandoned shelters, often with new plant tissue growing into the chambers, were common at the site; none contained pupal shells. The collection lends evidence to support the assumptions that more than one generation occur each season and that boxelder is a native host in California.

Two mature larvae taken in the Belmont lot (59F1) compare very well in structural details with the description of *P. aesculana* given by MacKay (1959). The head of the California examples is darker, being brown with darker, mottled areas laterally, and an

extensive black postgenal region. The thoracic shield is correspondingly darker, especially on one specimen; and it is considerably darker than the anal shield. Head measurements of the two individuals, length: width, were 0.90: 1.03 mm. and 0.94: 1.15 mm. Abdominal crotchets varied from 38-42.

Acknowledgement is made to the following, whose cooperation enabled study of the specimens in their care: J. F. Gates Clarke, U.S. National Museum; T. N. Freeman, Canadian National Collection; C. D. MacNeill, California Academy of Sciences; and F. H. Rindge, American Museum of Natural History.

LITERATURE CITED

HEINRICH, C.

1923. Revision of the North American Moths of the Subfamily Eucosminae of the Family Olethreutidae. U. S. Natl. Mus., Bull. 123, 298 pp.

MACKAY, M. R.

1959. Larvae of the North American Olethreutidae. Canad. Ent., 91, Suppl. 10, 338 pp.

PETERSON, L. O. T.

1958. The boxelder twig borer, *Proteoteras willingana* (Kearfott) (Lepidoptera: Olethreutidae). Canad. Ent., 90:639-646.

SCHAFFNER, J. V.

1950. Butterflies and Moths. Order Lepidoptera. in: Craighead, Insect Enemies of Eastern Forests. U. S. Dept. Agric., Misc. Publ., 657: 343-505.

TOWNES, H. AND M. TOWNES

1960. Ichneumon-flies of America north of Mexico: 2. Subfamilies Ephialtinae, Xoridinae, Acaentinae. U. S. Natl. Mus., Bull. 216 (2), viii + 676 pp.

OBSERVATIONS ON THE VOLUNTARY DISPLAY OF COREMATA IN *ESTIGMENE ACREA*

(Lepidoptera: Arctiidae)

C. DON MACNEILL

California Academy of Sciences, San Francisco

The existence of coremata or "brush-organs" on the abdominal venter of many adults of Arctiidae and related families has repeatedly been demonstrated (Muller, 1874; Bethune-Baker, 1925; Chretien, 1926; Eltringham, 1934, 1935; Lane, 1957). Such organs have been noted in *Estigmene acrea* (Drury) (Morrison, 1874; Stretch, 1883; Weed, 1883; Berlese, 1909:541). The above records report results of hand manipulation of living captives or of artificial inflation of dissected organs of dead specimens; how-



Powell, Jerry A. 1962. "Biological and taxonomic notes on two California species of *Proteoteras* (Lepidoptera: Tortricidae)." *The Pan-Pacific entomologist* 38, 191–195.

View This Item Online: <https://www.biodiversitylibrary.org/item/226203>

Permalink: <https://www.biodiversitylibrary.org/partpdf/237889>

Holding Institution

Pacific Coast Entomological Society

Sponsored by

IMLS LG-70-15-0138-15

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Pacific Coast Entomological Society

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.