

Algal Collections from Funafuti Atoll

V. J. CHAPMAN¹

IN SEPTEMBER, 1951, H.M.S. "Challenger" visited Funafuti Atoll in the Ellice Islands in connection with seismic experiments. Biological materials were collected from the reef and samples were dredged from the floor of the lagoon. No algal material was collected in the latter but some was obtained from the reef stations, the location of which is shown in Figure 1. In the report accompanying the material (Challenger 2320/12/3) it is stated that the seaweed samples are good random ones. Considerable differences in respect to both abundance and proportions of species are evident at the various stations. No algal collections appear to have been made on previous visits, not even during the celebrated visit to put down the bore through the reef in order to determine the depth of coral (Finckh, 1904). Recently an extensive treatment of the algae of Bikini Atoll has been published (Taylor, 1950) and this has proved very helpful in the present study. The list of species from Funafuti is not as extensive as one might have expected but this may be a result of the lack of a trained botanist among the collectors.

I am grateful to Dr. Drouet for checking one or two doubtful species among the Myxophyceae and to Dr. Yale Dawson for

the identification of *Wurdemannia miniata*. The collection has been deposited at the Auckland Institute and Museum, Auckland, New Zealand.

MYXOPHYCEAE

OSCILLATORIACEAE:

Lyngbya majuscula (Dill.) Harv. Sta. 1 (lagoon side; seaward side); Sta. 2 (seaward side); Sta. 3 (seaward side).

Lyngbya confervoides C. Ag. Sta. 2 (seaward side); Sta. 4 (seaward side); Sta. 5.

Lyngbya semiplena (C. Ag.) J. Ag. Sta. 5.

Oscillatoria nigro-viridis Thw. Sta. 1 (seaward side).

Phormidium corium (Ag.) Gom. Sta. 4 (seaward side).

Hydrocoleus lyngbyaceus Kütz. Sta. 4 (seaward side).

Hydrocoleus coccineus Gom. Sta. 4.

RIVULARIACEAE:

Calothrix confervicola Kütz. Sta. 4 (lagoon side).

CHLOROPHYCEAE

ULVACEAE:

Enteromorpha procera (Ahl.) f. *minuta* Chap. Sta. 5.

SIPHONOCLADACEAE:

Cladophoropsis membranacea (Ag.) Borg. Sta. 2 (seaward).

¹ Auckland University College, Auckland, New Zealand. Manuscript received November 29, 1954.

Cladophoropsis membranacea (Ag.) Borg. var. *repens* Phinney. Sta. 4.

Cladophoropsis zollingeri (Kütz.) Borg. Sta. 2 (seaward).

BOODLEACEAE:

Boodlea siamensis Reinbold. Sta. 1 (lagoon and seaward sides).

Struvea anastomosans (Harv.) Picc. et Grun. ex Piccone. Sta. 2 (seaward).

ANADYOMENACEAE:

Microdictyon setchellianum Howe. Sta. 4; Sta. 5.

Microdictyon japonicum Setchell. Sta. 4 (seaward). The plants were young but appeared to belong to this species.

VALONIACEAE:

Dictyosphaeria cavernosa (Forsk.) Borg. Sta. 2 (seaward); Sta. 3; Sta. 4 (lagoon and seaward sides); Sta. 5.

Dictyosphaeria versluisii Weber v. Bosse. Sta. 5. One plant which agrees with the description in Egerod (1952) is present.

Valonia ventricosa J. Ag. Sta. 1 (lagoon).

Valonia aegagropila C. Ag. Sta. 2 (seaward); Sta. 4 (seaward); Sta. 5.

DASYCLADACEAE:

Neomeris van-bosseae Howe. Sta. 4.

Acetabularia parvula Solms-Laubach. Sta. 1 (seaward).

CODIACEAE:

Avrainvillea pacifica Gepp. Sta. 5.

Halimeda opuntia (L.) Lamx. f. *triloba* Barton. Sta. 2 (lagoon); Sta. 3.

Halimeda opuntia (L.) Lamx. f. *elongata* (Barton) Barton. Dredged in 25 fathoms, 1.5 miles south of Fualifeke.

Halimeda tuna Lamx. var. *platydisca* (Dec.) Barton. Sta. 2 (lagoon side).

Halimeda incrassata (Ell. et Sol.) Lamx. Sta. 4 (lagoon side).

Halimeda gracilis Harv. f. *lata* Taylor. Dredged at 25 fathoms, 1.5 miles south of Fualifeke.

CAULERPACEAE:

Caulerpa urvilliana Mont. v. *typica* Weber-v.-Bosse. Sta. 5.




FIG. 1. Map of Funafuti Atoll showing the stations from which algal samples were collected and the place (D) where the dredged sample was obtained. Scale in nautical miles.

Caulerpa racemosa (Forsk.) J. Ag. v. *uvifera* (Turn.) Weber-v.-Bosse. Sta. 3.

Caulerpa racemosa (Forsk.) J. Ag. v. *uvifera* (Turn.) Weber-v.-Bosse f. *planiuscula* Svedel. Sta. 2 (lagoon side).

Caulerpa cupressoides (Vahl.) Ag. v. *typica* Weber-v.-Bosse. Sta. 3.

PHAEOPHYCEAE

ECTOCARPACEAE:

Ectocarpus indicus Sonder. Sta. 1 (seaward); Sta. 5.

DICTYOTACEAE:

Pocockiella variegata (Lamx.) Papenfuss. Sta. 1 (lagoon and seaward); Sta. 5.

Padina commersonii Bory. Sta. 2 (lagoon side). Probably this species though the material was sterile.

Dictyota dichotoma Lamx. Sta. 2 (seaward side).



Chapman, V. J. 1955. "Algal Collections from Funafuti Atoll." *Pacific science* 9(3), 354–356.

View This Item Online: <https://www.biodiversitylibrary.org/item/235774>

Permalink: <https://www.biodiversitylibrary.org/partpdf/243668>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.