

NEW SPECIES OF CHIONASPIS.

BY R. A. COOLEY, B. S., AMHERST, MASS.

At the request of Prof. T. D. A. Cockerell, through correspondence with Prof. Fernald, I was induced to take up the study of the genus *Chionaspis*, and Prof. Lull the genus *Pulvinaria*. Prof. Fernald prepared and sent out a circular letter to all entomologists whose addresses could be obtained, in this and other countries, and personal letters were also sent to the leading coccidologists, asking for as many species as possible to aid in the preparation of monographs of these two genera. The result has been most gratifying, for already a very large amount of material has been received.

In the material before me the following new species of *Chionaspis* have been found, and are published now in preference to waiting till the monograph is issued. The studies on these insects are being made in Prof. Fernald's entomological laboratory connected with the Massachusetts Agricultural College, where every possible facility is afforded for breeding and studying insects, together with very complete literature of the subject.

Chionaspis Cockerelli, n. sp.

Scale of female.—The female scale is about 3.2 mm. long, straight or very slightly curved, moderately thick in texture, slightly convex, white, with the exuviae pale yellowish-brown, the second skin being covered with secretion.

Female.—The pygidium is distinctly notched at the end, the sides of the notch being formed by the divergent median lobes. These lobes are firmly united at the base and have serrate edges. Two distinct parallel spines arising from the bottom of the notch are about as long as the distance between the inner edges of the lobes at the base. Compared with the other lobes of the pygidium the median ones are larger and extend farther into the body. Each lobe of the second pair is composed of two well rounded and distinct lobules, the incision between them extending to the base of the lobe. The inner lobule is larger and extends posteriorly about even with the median lobes. The third pair of lobes may be present or aborted; when present they are broad and low, with an elongated pore anterior to the base of each. Between the median and second pair is a minute spine, followed by a plate which is about as long as the second pair of lobes, and following these is a conical projection bearing a marginal pore. Outside of the second lobe is a spine, a plate

and a marginal pore, this plate being a little larger than the first one. Following the third lobe, when it is present, or a space when it is absent, there are two spines, one above and one below. These are followed by a plate and a distinct marginal pore, and after an interval interrupted by one or two spines, another plate, and following this another interval, terminated by a group of about three plates.

The spinnerets are in five groups: median, 7-9; anterior laterals, 17-23; posterior laterals, 23-34.

Described from dead and shrunken specimens.

Scale of male.—Length, 1.2 mm.; feebly carinated, white, with the larval skin almost colourless.

Described from a single imperfect specimen.

Male.—Male insect unknown.

The specimens were taken by Mr. Alexander Craw, on palm imported from China to San Francisco, Cal., July 11, 1897.

I take pleasure in naming this insect after Prof. T. D. A. Cockerell, who has made extensive and valuable contributions to our knowledge of the Coccidæ, and has shown me many kindnesses in my work on this group of insects.

Chionaspis aucubæ, n. sp.

Scale of female.—The female scale somewhat resembles that of *Chionaspis Lintneri* in outline, being strongly broadened posteriorly and abruptly rounded at the extremity. It is moderately convex, about 3 mm. in length and about 2 mm. in width. The exuviae at the apex of the scale have the first skin very pale yellow, and the second yellowish or brownish. The second skin is covered with a slight secretion. The scale itself is white and very thick and strong. There is a partial ventral scale at the anterior end.

Female.—As I had only dead and dry specimens of this insect, I made no attempt to describe anything but the pygidium of the female. Median lobes moderate in size, divergent, united at the base, with their inner edges distinctly serrate. Each lobe of the second pair is composed of two rounded lobules, the incision between the two reaching nearly or quite to the base of the lobe. The inner lobule is larger and projects farther posteriorly than the outer, sometimes surpassing the median lobes. The third lobe is simple and sometimes rudimentary. Between the bases of the median lobes is a pair of minute convergent spines. On each side between the median and second lobes are a spine, a plate and a marginal

pore, and between the second and third lobes two spines, one above and one below, followed first by one or two plates, and then by a conical projection bearing a marginal pore. Outside of the third lobe are a spine and from one to three plates, then a slight notch, immediately followed by a marginal pore and after a space two unequal spines and about three plates. Following these plates are a notch and a marginal pore, then after a space a group of about five plates.

Spinnerets arranged in five groups: median, 8-14; anterior laterals, 19-28; posterior laterals, 19-33.

Scale of male.—The male scales are much more numerous than those of the female. They are white, delicate in texture, about 1.2 mm. in length, the larval skin at the anterior end being colourless or slightly yellowish. The scale itself may be parallel sided or slightly broadened posteriorly, and is indistinctly carinated.

Male.—Male insect unknown.

On Aucuba from Japan. Discovered by Mr. Craw in the course of his quarantine work at San Francisco. The scales are grouped together on one side of the leaf beneath, and the edge of the leaf is folded under, almost completely hiding them from view.

Chionaspis wistariæ, n. sp.

Scale of female.—The female scale is about 2 mm. in length, though some specimens are slightly longer, moderately broadened, dirty white in colour and delicate in texture, being a close imitation of the epidermis of the bark on which it rests. The scales usually occur in the longitudinal cracks of the bark, and are partially concealed under the epidermis. They are very often pressed out of the normal form. The exuviae are brownish, and the second skin is covered with secretion.

Female.—The following description of the female was made from dead and shriveled insects. The median pair of lobes is large and conspicuous, the second pair considerably smaller, and the third pair obsolete. The median lobes are darker in colour than any other part of the pygidium, firmly joined at the base, their inner edges parallel and nearly touching each other for about half their length, then diverging at about a right angle, with the exposed edges serrate. The second lobe is composed of two lobules, the inner one being the larger. Within the outer edge of each of the median lobes is a spine, and next to this a short blunt plate, followed by a marginal pore. Between the lobules of the second lobe is a spine, and outside of the second lobe are a plate and

two marginal pores, followed first by a spine and then by a plate, which is about as long as the median lobes, and often forked at the tip. Outside of this plate are two marginal pores, followed by a spine and one or two plates, then after another marginal pore a group of about four plates.

There are five groups of spinnerets : median, 8-15 ; anterior laterals, 19-31 ; posterior laterals, 13-23.

Scale of male.—The male scale, as in all other species of this genus, is elongated in form and white in colour. The sides are nearly parallel, and it is distinctly tri-carinated. Length, about 1 mm. The larval skin resembles the anterior or smaller one of the female scale.

Male.—Male insect unknown.

Discovered by Mr. Craw, July 8, 1897, at San Francisco, on the bark of Wistaria from Japan.

Chionaspis pinifoliae heterophyllae, n var.

Scale of female.—The scale of the female is indistinguishable from that of *pinifoliae*, Fitch, having the same range of form and size, the colour of the scale and exuviae being the same. The scales vary in size from 2 mm. to 3.4 mm., the average length being about 2.5 mm. The scale is white, strongly convex, with the exuviae at the anterior extremity yellow, both skins being naked.

Female.—The description of the female is made from dead and shriveled specimens. At the anterior end of the body are two distinct, curved bristles, which may be the rudiments of the antennae ; these are found also in *pinifoliae*. The last segment terminates in a median notch, the sides of which are formed by the divergent median lobes. The lobes of the second pair are low and inconspicuous, and each one is composed of two lobules of about equal size. Two minute spines, one above and one below, arise from near each median lobe, though back from the edge of the segment. Contiguous to each median lobe is a simple plate, outside of which is a marginal pore. Between the lobules of the second lobe is a distinct spine, and outside of this lobe is a plate with a spine at its base, followed by a marginal pore. Outside of the rudimentary third lobe is a marginal pore, followed by a spine and a plate with a spine at its base. Then follows a pronounced marginal pore, a short interval, another space and a long interval, interrupted only by a spine, and terminated by the fourth and last plate.

There are five groups of spinnerets : median, 4-8 ; anterior laterals, 12-18 ; posterior laterals, 14-16. The chief characters by which *pinifoliae*



Cooley, R. A. 1897. "New species of Chionaspis." *The Canadian entomologist* 29, 278–282. <https://doi.org/10.4039/Ent29278-12>.

View This Item Online: <https://www.biodiversitylibrary.org/item/88720>

DOI: <https://doi.org/10.4039/Ent29278-12>

Permalink: <https://www.biodiversitylibrary.org/partpdf/245154>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: Not provided. Contact Holding Institution to verify copyright status.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.