

Sporidia cylindrica, obtusa, 4—6 cellularia, quaque cellula primitus nucleo 1 magno instructa, recta, hyalina disticha, 24/8. Paraphyses desunt.

Ad baccas putridas *Juniperi nanae* prope Kühtai (Oetz) alpium Tyrolensium, alt. c. 6500'. Dr. Rehm.

61. *Phomatospora Saccardoi* Rehm nov. spec.

Perithecia conica, seriatim sub peridermide nidulantia, dein apice conice elongato protuberantia, atra, minutissima. Asci cylindracei, tenerrimi, 8 spori, 60/5. *Sporidia* obtuse cylindrica, 1 cellularia, 2 nucleis praedita, hyalina, 1 sticha, —7/3. Paraphyses desunt.

Ad caulem putrescentem *Cirsii spinosiss.* prope Alpeiner-Gletscher in valle Stubai alpium Tyrolensium, alt. c. 7000'. Dr. Rehm.

62. *Chätomium Polypori* Rehm nov. spec.

Perithecia sparsa, minima, globosa, atra, scabra, sessilia, setis simplicibus, brevissimis, acutis, fuscis, c. 2 mikr. crassis obsessa. Asci cylindracei, 8 spori, 45/5. *Sporidia* subrotunda vel rotunda, fuscidula, 1 cellularia, fuscidula, 5 mikr. diam., 1 sticha. Paraphyses filiformes, tenerae.

Ad Lenzitem sepiariam prope Augustam Vindelicorum. Britzelmayr.

Sordariae novae.

Auctore C. A. J. A. Oudemans.

Sordaria Winterii mihi. *Perithecia sparsa*, usque ad collum immersa, subglobosa, fusco-nigra, translucientia, glabra, absque collo glabro, recto, conico-obtuso, 150 μ longo, 425 μ in diametro metientia. Asci 8-spori, perfecte cylindracei, breviter pedunculati, apice truncati, membrana duplici apice incrassata et resupinata, 150 μ longi, 9—10 μ crassi, paraphysibus nullis. Sporae monostichae, primitus ellipticae, denique elliptico-oblongae vel oblongae, brevi ante maturitatem 23—25 μ long., 15—20 μ latae, post ascorum destructionem, itaque prorsus maturae 21—22 μ longae, 10—11 μ latae, nigrae, lucidae, vacuolo centrali, neque appendiculatae, neque zona mucilaginis circumdatae. — In fimo Cameli Bactriani. — Accedit ad *S. humanam*, sed differt sporarum forma, paraphysarum absentia, peritheciolorum dimensione et collo.

Sordaria Hansenii mihi. *Perithecia sparsa*, in fimo cuniculorum siccato inter straminis fibrillas collo tantum prominentia, subglobosa, 350 μ in diametrum metientia, in collum breve-conicum, 23 μ longum, setoso-hirtum desinentia, glabra, fusco-nigra, subtranslucentia. Asci lanceolati, 150 μ longi, 12 μ lati, antrorsum breviter, deorsum longius contracti, polyspori, breviter pedunculati. Sporae discifor-



Oudemans, C. A. J. A. 1882. "Sordariae novae." *Hedwigia* 21, 123–124.

View This Item Online: <https://www.biodiversitylibrary.org/item/14427>

Permalink: <https://www.biodiversitylibrary.org/partpdf/246958>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.