THE ISOPOD GENUS *ICHTHYOXENUS* HERKLOTS, WITH DESCRIPTION OF A NEW SPECIES FROM JAPAN.

By HARRIET RICHARDSON,

Collaborator, Division of Marine Invertebrates, United States National Museum.

In 1870 Herklots¹ described the type-species of the genus *Ichthyoxenus* which was found parasitic on the fish *Barbodes maculatus* Bleeker from Java. The mode of life of this isopod is peculiar and interesting. It bores a hole in the body of the fish just back of the lateral fin, where it lives with its mate. It enters this cavity when young. As it grows in size, it is unable to leave the body of the fish and becomes many times larger than the opening through which it entered. This species was named after Mr. Jellinghaus, who was the first to notice the parasite, which he called "songkeat," and who in 1860 wrote a note concerning it to the Société physique des Indes néerlandaises.² The fish was taken from the River Tjikerang, district of Tjilokotot, regency of Bandong.

The second species of the genus, *Ichthyoxenus montanus*, was described by Schiøedte and Meinert³ in 1884. It was found on *Puntius sophores* in the Himalayan Mountains.

Up to the present time these two were the only species known.

In 1908 Maj. P. A. Ouwens⁴ mentioned another host for *Ichthyoxenus jellinghausii*, the fish *Nemacheilus fasciatus* van Hasselt, from the Tji-Seroema near Batavia.

Not long ago Dr. Harold Heath wrote to me concerning some parasites which were found on Japanese fishes from Lake Biwa. They were described as living in the body of the fish, which they entered through a hole bored just back of the lateral fin. The specimens were sent to the United States National Museum by Mr. Will Thompson, and the parasites proved to be a new species of *Ichthyoxenus*. The hosts are *Acheilognathus rhombeum* (Schlegel),

¹ Archives Néerlandaises des Sciences, vol. 5, 1870, pp. 128-137, pl. 5, figs. 10-18.

² Natuurk. Tijdschr. Nederl.-Indië, vol. 22, 1860, p. 378.

³ Nat. Tidsskrift (3), vol. 14, 1884, pp. 303-309, pl. 11, figs. 10-11.

⁴ Natuurk. Tijdschr. Nederl.-Indië, vol. 67, 1908, pp. 29-35.

Gnathopogon elongata (Schlegel), *Acheilognathus tabira* Jordan and Thompson, MSS., *Acheilognathus lanceolatum* (Schlegel), and *Acheilognathus cyonostigma* (Jordan and Fowler). I have since found this parasite on *A. limbatum* Jordan and Snyder from the same locality.

ICHTHYOXENUS JELLINGHAUSII Herklots.

Ichthyoxenus jellinghausii HERKLOTS, Archives Néerlandaises, vol. 5, 1870, pp. 128-137, pl. 5, figs. 10-18.—SCHIEDTE and MEINERT, Naturhistorisk Tidsskrift (3), vol. 14, 1884, pp. 298-303, pl. 11, figs. 5-6.—MAX WEBER, Zool. Ergebnisse, vol. 2, 1892, pp. 557-560, pl. 30, fig. 1.—WILLINK, Natuurk. Tijdschr. Nederl.-Indië, vol. 64, 1905, pp. 156-161.—OUWENS, Natuurk. Tijdschr. Nederl. Indië, vol. 67, 1908, pp. 29-35.

Locality.—A male and a female, collected by Dr. Owen Bryant and Mr. William Palmer in Java, are in the United States National Museum. The host is not known.

FIG. 1.—*ICHTHYOXENUS JELLINGHAUSII*, BUITENZORG. FEMALE. $\times 3\frac{1}{2}$.

This species has been recorded from Bandon and Batavia. The specimen in the United States National Museum is from Buitenzorg, and if the figures of Herklots and Schiedte and Meinert are correct it differs from the type in the following points: The first

FIG. 3.—*ICHTHYOXENUS JELLINGHAUSII*, BUITENZORG. SEVENTH LEG OF FEMALE. $\times 7\frac{1}{2}$.

segment of the thorax in the female has the anterior margin deeply excavate in the middle, the excavation being rectangular; the seventh pair of legs have the carpus and merus very much inflated, with the inner margin produced in a rounded expansion; the merus is also much elongated.

For the purpose of comparison with the new species from Japan, I have figured the specimen in the United States National Museum, which I have identified as *Ichthyoxenus jellinghausii* Herklots. Since I have not seen the type-species and find the Buitenzorg specimens so close to the description and figures of this form, I have thought it best to do this.

FIG. 2.—*ICHTHYOXENUS JELLINGHAUSII*, BUITENZORG. MALE. $\times 5\frac{1}{2}$.

ICHTHYOXENUS JAPONENSIS, new species.

Body of female oblong-ovate, almost twice as long as wide, 13 mm. long, 7 mm. wide. Surface smooth. Color pale yellow, with scattered black dots.

Head small, triangular, 2 mm. long, $2\frac{1}{2}$ mm. wide; front rounded. Eyes large, oblong, about 1 mm. in length and separated by a distance equal to 1 mm. or the length of one eye. First antennæ short, composed of six articles and extending to the middle of the eye or to the end of the fifth article of the second antennæ. Second antennæ composed of seven articles and extending almost to the post-lateral angle of the head.

The first segment of the thorax is about twice as long in the median line as any of the three following, being $1\frac{1}{2}$ mm. in length. The second, third, and fourth segments are subequal. The last three segments gradually decrease in length, the three together measuring $1\frac{1}{2}$ mm. The anterior margin of the first segment is sinuate in the middle, the lateral parts projecting laterally beyond the

head and being posteriorly constricted. Epimera are present on the last six segments, those of the second, third, and fourth segments

being narrow, elongate plates occupying the anterior half of the lateral margin; those of the last three segments are about as long as wide, with the posterior extremities rounded, and are placed just anterior to the lateral margin of their respective segments.

The abdomen is deeply immersed in the thorax, the lateral parts of the first two segments being completely covered on one side, the second segment partly showing on the other side. The first five segments gradually increase in length, all together measuring $2\frac{1}{2}$ mm. The sixth or terminal segment is a little longer than wide, $4\frac{1}{2}$ mm. long, 4 mm. wide, and is posteriorly widely

rounded. Its posterior half is thin, semitranslucent, and colorless, with no markings. The uropods are much shorter than the terminal segment of the abdomen, and do not reach its extremity by a distance equal to 1 mm.; the branches are equal in length, narrow, tapering, with rounded extremities, the outer one being slightly narrower than the inner one. The pleopods are also short, not quite reaching the

FIG. 4.—*ICHTHYOXENUS JAPONENSIS*, LAKE BIWA, FROM *ACHEILOGNATHUS CYONOSTIGMA*. FEMALE. $\times 5$.

FIG. 5.—*ICHTHYOXENUS JAPONENSIS*, LAKE BIWA, FROM *ACHEILOGNATHUS LIMBATUM*. MALE. $\times 5\frac{1}{2}$.

tip of the uropods and being 1 mm. shorter than the extremity of the terminal abdominal segment.

All seven pairs of legs are prehensile. The last pair has the carpus and merus dilated in rounded expansions on the inferior side. The merus is short, being about one-third the length of the ischium. In the last four pairs of legs these articles show an increasing expansion.

Some of the specimens obtained are larger than the one figured and described, but in most of these the abdomen is curled up at the extremity. I have therefore taken the smallest one as the type.

The male differs from the female in size, being much smaller; in the more symmetrical body, longer antennæ, narrower head, and smaller terminal segment of the abdomen.

The type-specimens, male and female, were taken from *Acheilognathus cyonostigma* (Jordan and Fowler), found in Lake Biwa, and two males and two females were taken from the same host at Omi, Lake Yogo.

FIG. 6.—*ICHTHYOXENUS JAPONENSIS*, LAKE BIWA, FROM *ACHEILOGNATHUS CYONOSTIGMA*. SEVENTH LEG OF FEMALE. $\times 14\frac{1}{2}$.

Other specimens are as follows: A male and a female on *Acheilognathus lanceolatum* (Schlegel) from Funayado, Kiusiu, Japan, and a male and a female on the same host from Omi, Lake Yogo; a male and a female on *Acheilognathus tabira* Jordan and Thompson, MSS., from Lake Biwa; one female on *Gnathopogon elongata* (Schlegel) from Lake Biwa; two males and two females from two specimens of *Acheilognathus rhombeum*

(Schlegel) from Lake Biwa; two males and two females from *Acheilognathus limbatum* Jordan and Snyder from Lake Biwa.

This species differs from *I. jellinghausii* in the larger eyes, the larger and more circular form of the terminal segment of the abdomen, the shorter uropoda, the shorter and differently shaped merus of the seventh pair of legs, and the larger head and wider abdomen, compared with the thorax.

Type.—Cat. No. 45617, U.S.N.M.

LIST OF REFERENCES.

- HERKLOTS, J. A. Deux nouveaux genres de Crustacés vivant en parasites sur des poissons-epichthys et ichthyoxenos. Archives Néerlandaises des Sciences exactes et naturelles, vol. 5, 1870, pp. 120-137, pl. 5, La Haye.
- OUWENS, P. A. Nog iets over Ichthyoxenus jellinghausii (Herklots). Natuurkundig Tijdschrift voor Nederlandsch-Indië, vol. 67, 1908, pp. 29-35, Weltevreden.
- SCHIEDTE, J. C., and MEINERT, FR. Symbolæ ad monographiam cymothoarum crustaceorum isopodum familiæ. IV. Cymothoidæ. Trib. II. Cymothoinæ. Trib. III. Livonecinæ. Naturhistorisk Tidsskrift (3), vol. 14, 1884, pp. 221-421, pls. 6-18, Kjøbenhavn.
- WEBER, MAX.¹ Die Süßwasser-Crust. des Indischen Archipels, nebst bemerkungen über die Süßwasser-Fauna im Allgemeinen. Zool. Ergebnisse, vol. 2, 1892, pp. 557-560, pl. 30, fig. 1, Leiden.
- WILLINK, H. D. TIEENK. De "Songkeat" een vischparasiet. Natuurkundig Tijdschrift voor Nederlandsch-Indië, vol. 64, 1905, pp. 156-161, Amsterdam.

¹ I have not been able to obtain this paper.

Richardson, Harriet. 1913. "The isopod *Ichthyoxenus* Herklots, with description of a new species from Japan." *Proceedings of the United States National Museum* 45, 559–562.

<https://doi.org/10.5479/si.00963801.45-1995.559>.

View This Item Online: <https://www.biodiversitylibrary.org/item/31799>

DOI: <https://doi.org/10.5479/si.00963801.45-1995.559>

Permalink: <https://www.biodiversitylibrary.org/partpdf/24713>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.