

BEES OF THE NEVADA TEST SITE¹

Dorald M. Allred

During the years from 1959 to 1965, many bees were collected at the Nevada Test Site as part of the ecological studies conducted by the Department of Zoology and Entomology of Brigham Young University, under contract with the U. S. Atomic Energy Commission. Most of these were recently identified by Dr. G. E. Bohart, Entomology Research Division, Department of Agriculture, Logan, Utah, and Mr. P. H. Timberlake, Riverside, California. The identifications form the basis of this report on the host relationships, seasonal occurrence, and geographic distribution at the test site.

Further studies specifically directed toward the bees, particularly during the blooming periods of the desert plants, undoubtedly would yield additional records. Genera and species are listed alphabetically for convenience. Areas of collection refer to divisions delineated by Allred, Beck, and Jorgensen (1963a and b). Methods of collection used were hand capture, aerial and sweep nets, and can pit-traps (Allred *et al.*, 1963a).

SPECIES AND PLANT ASSOCIATIONS

- Agapostemon cockerelli*: 2♀♀; areas 17, 18; *Malacothrix glabrata*; June, July.
- A. texana*: 2♂♂ 3♀♀; areas 5, 12, CB, *Asclepias erosa*, *Petradoria pumila*, *Stanleya pinnata* (most common host); June, Aug.
- Andrena* sp.: ♂; area CE; host unknown; March.
- Anthidium dammersi*: ♂♀; areas 1, 18; *Erigeron pumilus*; May; June.
- Anthophora californica*: ♂♀; area 5; *Stanleya pinnata*; June.
- A. porterae*: ♂; area 4; host unknown; March.
- A. urbana*: 8♂♂ 24♀♀; areas 5, 12, 17, 18, CB, TM; *Asclepias erosa*, *Astragalus lentiginosus*, *Gayophytum ramosissima*, *Malacothrix glabrata*, *Oenothera californica*, *Petradoria pumila*, *Stanleya pinnata*; most common host was *S. pinnata*, with *A. erosa* and *P. pumila* also frequently represented; most were taken in June, about half as many in July, and few in August.
- Anthophora* new species: 5♂♂ 1♀; area CE; host unknown; March.
- Ashmeadiella aridula*: ♀; area 2; *Pectis papposa*; Aug.
- A. australis*: ♀♀ 2?; area 12, *Penstemon palmeri*; July.
- A. bigeloviae*: ♀; area 10; *Salsola kali*; Sept.
- A. inyoensis*: 4♂♂; area 5; host unknown; May.
- A. opuntiae*: 2♂♂ 3♀♀; areas 12, 13, CB; *Echinocereus engelmannii* (most common host), *Opuntia erinacea*; May, July.
- Bombus morrisoni*: 7♀♀; areas 5, 12; *Petradoria pumila* (most common host), *Stanleya pinnata*, *Viguiera multiflora*; July, Aug.

¹BYU-AEC publication no. COO-1355-19.

- Centris rhodopus*: 19♂♂ 6♀♀; areas 5, 6, CB, 401, *Cleome lutea*, *Eriogonum inflatum*, *Stanleya pinnata* (most common host), *Tamarix pentandra*; June-Sept., mostly July.
- Ceratina nanula*: ♀; area 17; *Malacothrix glabrata*; June.
- Chelostomoides lobatifrons*: ♂; area CB; *Asclepias erosa*; June.
- Dialictus albohirtus*: 33♂♂ 3♀♀; areas 5, 6, 12, CB, EM, T; *Artemisia tridentata*, *Asclepias erosa* (most common host), *Chrysothamnus paniculatus*, *Eriogonum inflatum*, *Eriogonum fasciculatum*, *Larrea divaricata*, *Petradoria pumila*, *Salix gooddingii*, *Stanleya pinnata*, *Tamarix pentandra*; June (mostly), July.
- D. hyalinus*: ♂ 11♀♀; areas 1, 5, 6, 12, 17, CB, EC; *Acamptopappus shockleyi*, *Asclepias erosa*, *Calochortus flexuosus*, *Malacothrix glabrata*, *Penstemon palmeri*, *Stanleya pinnata*, *Viguiera multiflora*; May-Aug.
- D. incompletus*: 12♀♀; areas 12, FM; *Bromus rubens*, *Calochortus flexuosus*, *Penstemon bridgesii*, *Penstemon floridus*, *Penstemon palmeri* (most common host); July.
- D. microlepoides*: 11♂♂ 7♀♀; areas 12, CB, EM; *Artemisia tridentata*, *Asclepias erosa* (most common host), *Eriogonum fasciculatum*, *Penstemon bridgesii*, *Tamarix pentandra*, *Viguiera multiflora*; June (mostly), July.
- D. nevadensis*: 2♀♀ area 12; *Penstemon palmeri*; July.
- D. pruinosum*: ♂; area CB; *Asclepias erosa*; June.
- Dialictus* sp. 1, ? new: 2♀♀; area 12; *Castilleja linariaefolia*, *Chrysothamnus* sp.; July, Aug.
- Dialictus* sp. 2, ? new: ♂; area CB; *Asclepias erosa*; June.
- Dialictus* sp. 3, ? new: ♀; area 401; *Eriogonum fasciculatum*; July.
- Colletes euplophi*: ♂♀; areas 5, 17; *Sphaeralcea* sp., *Stanleya pinnata*; June.
- Colletes* sp., ? new: ♀; area 17; *Sphaeralcea* sp.; June.
- Diadasia australis*: 2♂♂; areas 12, 17; *Opuntia erinacea*; June, July.
- D. diminuta*: 4♂♂ 3♀♀; areas 5, 17; *Sphaeralcea ambigua* (most common host), *Stanleya pinnata*; June.
- D. lutzi*: ♂♀ 1?; areas 5, 17; *Sphaeralcea* sp., *Stanleya pinnata*; June.
- Dianthidium pudicum*: ♂♀; areas 16, 19; *Chrysothamnus viscidiflorus*; June, Aug.
- D. subparvum*: ♂♀; area 12; *Chrysothamnus viscidiflorus*, *Petrorhagia pumila*; July, Aug.
- D. ulkei*: 2♂♂; areas 12, FC; *Artemisia tridentata*; July.
- Dioxys productus*: ♂; area 5; host unknown; March.
- Dufourea* new species 1: ♀; area JA; host unknown; April.
- Dufourea* new species 2: ♀♀; area 12; *Phacelia peirsoniana*; June.
- Epeolus minimus*: ♀; area 5; host unknown; May.
- Evylaeus ruficornis*: ♀; area 19; host unknown; June.
- Halictus hyalinus*: ♀; area 401; *Chrysothamnus parryi*; Aug.
- H. tripartitus*: 7♀♀; areas 12, CB, FM; *Calochortus flexuosus*, *Eriogonum fasciculatum*, *Lupinus palmeri*, *Penstemon* sp., *Petrorhagia pumila*, *Viguiera multiflora*; June, mostly July, Aug.

- Heriades timberlakei*: 5♂♂ 4♀♀; areas 12, 13; *Opuntia erinacea*, *Viguiera multiflora*; July.
- Hesperapis wilmattae*: ♀; area 401; *Malacothrix glabrata*; June.
- Hylaeus asininus*: 2♂♂; area 12; *Chrysothamnus viscidiflorus*; Aug.
- Hypomacrotera subalpinus*: 4♂♂; area 5; *Baileya multiradiata*, *Sphaeralcea ambigua*, *Stanleya pinnata*; June.
- Lasioglossum sisymbrium*: 3♀♀ 5♂♂; areas 5, 12, 18, 19; *Chrysothamnus viscidiflorus*, *Erigeron pumilis*, *Hymenoxys cooperi*, *Penstemon* sp., *Stanleya pinnata*, *Viguiera multiflora*; June-Aug.
- Lithurgus apicalis*: 3♂♂ 1♀; areas 12, 13; *Opuntia erinacea*; July.
- Melectamorpha californica*: 2♂♂ 2♀♀; areas 5, CB; *Asclepias erosa*, *Stanleya pinnata*; June, July.
- Melissodes subagilis*: 7♀♀; areas 5, 12, 17; *Baileya multiradiata*, *Chrysothamnus viscidiflorus*, *Petradoria pumila*, *Sphaeralcea ambigua*, *Stanleya pinnata*; June-Aug.
- M. tristis*: ♀; area 12; *Petradoria pumila*; July.
- Micranthophora hololeuca*: 2♂♂ 2♀♀; area TA; *Sphaeralcea munroana*; June.
- M. phenax*: ♂; area 1; host unknown; April.
- Nomia tetrazonata*: ♂; area CE; *Berula erecta*; July.
- Osmia titusi*: ♂; area 5; host unknown; March.
- Osmia* sp., ? new: ♂♀; areas 5, 17; *Astragalus lentiginosus*, *Descurainia pinnata*; May, June.
- Perdita arcuata*: ♂ 2♀♀; area 5; *Sphaeralcea ambigua*; June.
- P. callicerata*: ♂; area 12; *Chrysothamnus paniculatus*; June.
- P. chloris*: ♀; area 16; *Eriogonum deflexum*; Aug.
- P. fallugia*: 5♂♂; areas 12, 18; *Cowanía* sp. (most common host), *Erigeron pumilus*, *Oenothera californica*; June.
- P. nasuta*: 2♀♀ 7♂♂; area 6; *Eriogonum inflatum*; July.
- P. thermophila*: ♂♀; areas 6, 410; *Eriogonum inflatum*; July.
- Perdita* new species: 8♀♀; areas 16, 401; *Dalea polyadenia*, *Eriogonum deflexum* (most common host); Aug.
- Sphecodes eustictus*: ♀; area 12; *Eriogonum deflexum*; Aug.
- Stelis* new species: ♀; area 10; host unknown; April.
- Tetralonia quadricincta*: ♂; area 4; host unknown; March.
- Tetralonia* new species "al": ♂ 2♀♀; areas 4, 5; *Stanleya pinnata*; March, June.
- Tetralonia* new species "an": ♂; area 4; host unknown; March.
- Tetralonia* new species "mo": ♀; area 4; host unknown; March.
- Tetralonia* new species "pr": 4♂♂ 32♂♂; areas 4, 5; host unknown; March.
- Triepeolus helianthi*: ♂; area 5; *Sphaeralcea ambigua*; June.
- Xylocopa californica*: ♀ 7♂♂; areas 5, M; *Stanleya pinnata*; June (mostly), July.

PLANT-BEE ASSOCIATIONS

Acamptopappus shockleyi
Dialictus hyalinus
Artemisia tridentata

Dialictus albohirtus
D. microlepidoides
Dianthridium ulkei

<i>Asclepias erosa</i>	<i>Eriogonum inflatum</i>
<i>Agapostemon texana</i>	<i>Centris rhodopus</i>
<i>Anthophora urbana</i>	<i>Dialictus albohirtus</i>
<i>Chelostomoides lobatifrons</i>	<i>Perdita nasuta</i>
<i>Dialictus albohirtus</i>	<i>P. thermophila</i>
<i>D. hyalinus</i>	<i>Gayophytum ramosissima</i>
<i>D. microlepoides</i>	<i>Anthophora urbana</i>
<i>D. pruinatum</i>	<i>Hymenoxys cooperi</i>
<i>Dialictus</i> sp. 2	<i>Lasioglossum sisymbrium</i>
<i>Melectamorpha californica</i>	<i>Larrea divaricata</i>
<i>Astragalus lentiginosus</i>	<i>Dialictus albohirtus</i>
<i>Anthophora urbana</i>	<i>Lupinus palmeri</i>
<i>Osmia</i> sp.	<i>Halictus tripartitus</i>
<i>Baileya multiradiata</i>	<i>Malacothrix glabrata</i>
<i>Hypomacrerata subalpinus</i>	<i>Agapostemon cockerelli</i>
<i>Melissodes subagilis</i>	<i>Anthophora urbana</i>
<i>Berula erecta</i>	<i>Ceratina nanula</i>
<i>Nomia tetrazonata</i>	<i>Dialictus hyalinus</i>
<i>Bromus rubens</i>	<i>Hesperapis wilmattae</i>
<i>Dialictus incompletus</i>	<i>Oenothera californica</i>
<i>Calochortus flexuosus</i>	<i>Anthophora urbana</i>
<i>Dialictus hyalinus</i>	<i>Perdita fallugia</i>
<i>D. incompletus</i>	<i>Opuntia erinacea</i>
<i>Halictus tripartitus</i>	<i>Ashmeadiella opuntiae</i>
<i>Castilleja linariaefolia</i>	<i>Diadasia australis</i>
<i>Dialictus</i> sp. 1	<i>Heriades timberlakei</i>
<i>Chrysothamnus paniculatus</i>	<i>Lithurgus apicalis</i>
<i>Dialictus albohirtus</i>	<i>Pectis papposa</i>
<i>Perdita callicerata</i>	<i>Ashmeadiella aridula</i>
<i>Chrysothamnus parryi</i>	<i>Penstemon bridgesii</i>
<i>Halictus hyalinus</i>	<i>Dialictus incompletus</i>
<i>Chrysothamnus viscidiflorus</i>	<i>D. microlepoides</i>
<i>Dianthidium pudicum</i>	<i>Penstemon floridus</i>
<i>D. subparvum</i>	<i>Dialictus incompletus</i>
<i>Hylaeus asininus</i>	<i>Penstemon palmeri</i>
<i>Lasioglossum sisymbrium</i>	<i>Ashmeadiella australis</i>
<i>Melissodes subagilis</i>	<i>Dialictus hyalinus</i>
<i>Chrysothamnus</i> sp.	<i>D. incompletus</i>
<i>Dialictus</i> sp. 1	<i>D. nevadensis</i>
<i>Cleome lutea</i>	<i>Penstemon</i> sp.
<i>Centris rhodopus</i>	<i>Halictus tripartitus</i>
<i>Cowanias</i> sp.	<i>Lasioglossum sisymbrium</i>
<i>Perdita fallugia</i>	<i>Petroradria pumila</i>
<i>Dalea polyadenia</i>	<i>Agapostemon texana</i>
<i>Perdita</i> sp.	<i>Anthophora urbana</i>
<i>Descurainia pinnata</i>	<i>Bombus morrisoni</i>
<i>Osmia</i> sp.	<i>Dialictus albohirtus</i>
<i>Echinocereus engelmanni</i>	<i>Dianthidium subparvum</i>
<i>Ashmeadiella opuntiae</i>	<i>Halictus tripartitus</i>
<i>Erigeron pumilis</i>	<i>Melissodes subagilis</i>
<i>Anthidium dammersi</i>	<i>M. tristis</i>
<i>Lasioglossum sisymbrium</i>	<i>Phacelia peirsoniana</i>
<i>Perdita fallugia</i>	<i>Dufourea</i> sp. 2
<i>Eriogonum deflexum</i>	<i>Salix gooddingii</i>
<i>Perdita chloris</i>	<i>Dialictus albohirtus</i>
<i>Perdita</i> sp.	<i>Salsola kali</i>
<i>Sphecodes eustictus</i>	<i>Ashmeadiella bigeloviae</i>
<i>Eriogonum fasciculatum</i>	<i>Sphaeralcea ambigua</i>
<i>Dialictus albohirtus</i>	<i>Diadasia diminuta</i>
<i>D. microlepoides</i>	<i>Hypomacroteria subalpinus</i>
<i>Dialictus</i> sp. 3	<i>Melissodes subagilis</i>
<i>Halictus tripartitus</i>	<i>Perdita arcuata</i>



Allred, Dorald M. 1969. "BEES OF THE NEVADA TEST SITE." *The Great Basin naturalist* 29, 20–24.

View This Item Online: <https://www.biodiversitylibrary.org/item/33897>

Permalink: <https://www.biodiversitylibrary.org/partpdf/247839>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Brigham Young University

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.