tegerrima, coriacea, basi attenuata, utrinque punctis parùm elevatis, resinosi, et pube brevissimâ cinereâ copiosè instructa, subtùs obsoletè trinervia. Flores in apice ramulorum sæpiùs 5, rariùs 3 v. 7, brevissimè pedicellati. Involucrum semipollicis longitudine. Flosculi purpurei. Pappus cinereus.

Nomen e vocibus græcis *πεντε* quinque, et *φέρω* porto, compositum.

Genus a præcedente distinctissimum pappi radiis multiplici ordine dispositis, atque involucri connivente, 5-floro, undè nomen.

1. *P. foliolosus.*

In Chili. *Caldcleugh.* 2.

ADDITIONS AND CORRECTIONS.

After *TRIXIS corymbosa*, p. 188, insert the three following Species.

T. glabra, foliis petiolatis ovato-oblongis mucronulatis integerrimis glabris, paniculâ foliosâ.

In Mexico prope Laguna Verde. *Schiede et Deppe.* 2.
Floret Martio.

Caulis fruticosus, ramosissimus, lævis. *Rami* diffusi, cylindrici, glabri. *Folia* alterna, petiolata, ovato-oblonga, mucronulata, integerrima, membranacea, utrinque glaberrima, subtùs reticulata, basi rotundata, sesquiuncialia. *Panicula* foliosa, multiflora. *Bracteolæ* subsolitariae, lineares, acutæ, involucri duplò breviores. *Involucrum* 8-phyllum: *foliolis* linearibus, acutis, pubescentibus, erectis, simplici serie digestis.

gestis. *Receptaculum* densè pilosum. *Flosculi* 8, hermaphroditi, bilabiati, lutei; *labio exterior*e ligulato, tridentato; *interiore* bipartito, revoluto. *Antheræ* appendiculâ lineari obtusiusculâ elongatâ cartilagineâ terminatæ, basi bisetosæ: *setis* longis, simplicibus, attenuatis. *Stigmata* soluta, truncata, apice minutè papillosa. *Achenia* angusta, angulata, papilloso-scabra: *disco epigyno* dilatato. *Pappus* capillaris, denticulato-scaber, cinereo-fulvellus.

T. salicifolia, foliis elongato-lanceolatis acuminatis denticulatis subtùs tomentosis basi auriculatis, involucris multibracteolatis pedunculisque glanduloso-pubescentibus.

In Brasiliâ. *Sello.* 2.

Caulis fruticosus, diffusè ramosissimus, teres, striatus. *Rami* hirsuti. *Folia* alterna, sessilia, elongato-lanceolata, acuminata, membranacea, margine denticulis plurimis subsetaceis ornata, suprâ hirsuta, subtùs cinereo-tomentosa, basi attenuatâ et auriculatâ, 4—5 pollices longa, 10 lineas lata. *Panicula* ramosissima, divaricata. *Pedunculi* teretes, densè glanduloso-pubescentes. *Involucrum* 8—10-phyllum, basi bracteolis 5 v. 7, lineari-lanceolatis, acuminatis, patulis munitum: *squamis* lineari-lanceolatis, acuminatis, membranaceis, leviter carinatis, pube copiosâ glandulisque intermixtâ vestitis. *Receptaculum* densè pilosum. *Flosculi* plures (10—12) hermaphroditi, bilabiati, æquales: *fauce* tubo angusto 5-angulo longiore, dilatatâ, intùs villosâ: *labiis* brevioribus; *exteriore* ligulato, acutè tridentato; *interiore* bipartito, revoluto: *segmentis* lanceolato-linearibus, obtusiusculis. *Filamenta* membranacea, canaliculata, glabra. *Antheræ* appendiculâ lineari-lanceolatâ obtusâ subfalcatâ coronatæ, basi *setis* 2 simplicibus longis instructæ. *Stigmata* recurvata, apice truncata, minutè papillosa. *Achenia* undique

undique papilloso-scabra: *disco epigyno* dilatato, concavisculo. *Pappus* capillaris, flavicans.

T. mollissima, foliis petiolatis ovato-lanceolatis acuminatis serratis basi attenuatis utrinque sericeo-villosissimis.

In Brasiliâ. *Sello.* 2.

Frutex erectus, ramosissimus. *Rami* teretes, undique tomentosi. *Folia* alterna, petiolata, ovato-lanceolata, acuminata, serrata, basi acuta, penninervia, utrinque sericeo-villosissima, suprâ demùm viridia, 4-pollicaria. *Flores* diffusè corymboso-paniculati. *Bracteæ* lanceolatae, acutae, integerrimae. *Involucrum* duplici ordine polyphyllum: *foliolis* lanceolatis, acuminatis, sericeo-villosissimis; *interioribus* angustioribus. *Receptaculum* densè pilosum. *Flosculi* plures, hermaphroditi, bilabiati; *labio exteriori* ligulato, tridenticulato, 4-nervio; *interiore* bipartito, revoluti: *segmentis* lanceolatis, acuminatis, binerviis. *Antheræ* appendiculâ lineari, obtusâ breviori cartilagineâ coronatae, basi biseosae: *setis* compressis, vix attenuatis. *Achenia* subpentagona, undique copiosè papilloso-glandulosa: *disco epigyno* dilatato, concavo. *Pappus* capillaris, denticulato-scaber, cinereo-fulvellus.

TRIXIS divaricata, p. 190.

The flowers of this species are most probably entirely white, although in the dried state they have assumed a yellow colour. This species, as well as *cacaloides*, have been already referred to *Trixis* by Sprengel.

After *JUNGIA spectabilis*, p. 227, insert

3. *J. pyramidalis*, foliorum lobis obtusis pubescentibus, stipulis maximis,

maximis, paniculâ amplâ terminali, corollæ labio exteriori tridenticulato.

In Brasiliâ ad ripas fluminis La Plata. *Sello. v.*

Planta pube brevissimâ densè vestita, rubiginosa. *Caulis* leviter angulatus, crassitie digiti minoris. *Folia* alterna petiolata, subrotundo-cordata, 7-loba, 5-nervia, suprâ planiuscula, punctis minutissimis granulosa, subtùs reticulato-venosissima, varicosa, latitudine 2—3-uncialia: *lobis* obtusis, brevibus, grossè inæqualiterque dentatis, margine paululùm revolutis. *Petoli* teretes, 2—3-pollicares, basi dilatâtâ, concavâ. *Stipulæ* 2, maximæ, foliaceæ, lobato-dentatæ, sessiles, basi parùm angustatæ, 5-nerviæ, reticulato-venosissimæ, pollicares v. sesquipollicares. *Panicula* terminalis, ampla, laxa, tripedalis: *ramis* elongatis, multifloris. *Pedicelli* filiformes, densè pubescentes. *Bracteolæ* lanceolatæ, acuminatæ, dentatæ; *ultimis* integerrimis. *Involucra* simplici ordine polyphylla: *foliolis* lanceolatis, acuminatis, membranaceis, striatis, pubescentibus. *Paleæ* foliolis involucris conformes, lanceolatæ, acuminatæ, membranaceæ, concavæ, striatæ, margine scariosæ. *Flosculi* 10 circiter, albi, bilabiati, hermaphroditi; *labio* exteriori ligulato, revoluti, 4-nervio, obtusè tridenticulato, tubo brevior; *interiore* bipartito, spiraliter revoluti: *segmentis* linearibus, obtusis, binerviis, adprimùm margine conglutinis. *Filamenta* capillaria, glabra. *Antheræ* coalitæ, appendiculâ lineari, falcatâ acutiusculâ cartilagineâ coronatæ, basi bidentatæ: *dentibus* lineari-lanceolatis, acutis. *Stylus* capillaris. *Stigmata* soluta, semicylindrica, recurvata, apice dilatato, truncato, minutè papilloso. *Achenia* angustè fusiformia, sulcata, minutè papillosa: *disco* epigyno dilatato, planiusculo. *Pappus* cinereus: *radiis* plurimis (28) elegant

ganter plumosis, simplici ordine digestis, imâ basi connatis, patentibus, æqualibus.

PROSELIA, p. 234.

Two new species of this genus have been discovered by Dr. Gillies, of which he has had the kindness to favour me with specimens. The rays of the pappus in these are disposed in a double series.

TYLLOMA, p. 238.

The flowers in the specimen of *Tylloma limbatum* which I had to examine being hardly sufficiently advanced to admit of my ascertaining their structure with perfect accuracy, I beg to subjoin the following amendments of the character and description from an inspection of more complete specimens, for which I am indebted to the liberality of Dr. Gillies.

Flosculi radii foeminei, bilabiati; *labio interiore* brevissimo, erecto, bifido. *Pappus* capillaris: *radiis* simplici ordine digestis.

Flosculi radii numerosi (18—20) bilabiati, foeminei, staminibus sterilibus; *labio exteriori* amplo, ligulato, obtusè tridenticulato, patulo, 4-nervio, subtùs sericeo-villosissimo, nervis lateralibus longè intramarginalibus simplicibus, intermediis supernè bifurcatis, ramis apice confluentibus; *interiore* lanceolato, acuminato, erecto, bifido, brevissimo; *disci* hermaphroditi, tubulosi, bilabiati: *labiis* erectis, abbreviatis; *exteriore* obtusè tridentato; *interiore* bilobo.

Obs.—Hujus flosculi aurei, nec rosei, radio semipollicari. In alterâ specie ab amicissimo D. Gillies lectâ flosculi albi; *radii* 10, foeminei, absque sterilium rudimentis; *labio exteriori* ligulato, acutè tridentato; *interiore* brevissimo, bipartito, laciniis linearibus, acutis, erectis, inæqualibus.

DIAZEUXIS?

DIAZEUXIS? *serrata*, p. 254.

In a collection of dried plants, which Mr. Lambert lately received from Messrs. Schiede and Deppe, is a specimen in flower of this plant, which proves to be a genus akin to *Serratula*; and it is therefore to be expunged from *Diazeuxis*, to which it had been referred from habit alone, and the following inserted in its place.

2. *D. latifolia*, foliis lanceolatis acuminatis triplinerviis basi attenuatis, involucris squamis ovato-lanceolatis, pappi foeminei radiis fasciculatis.

Ad Caracas. *D. Fanning.* 2.

Præcedenti proxima. *Rami* angulati, sulcati, undique lanâ fugaci vestiti. *Folia* brevissimè petiolata, latè lanceolata v. elliptico-oblonga, acuminata, triplinervia, coriacea, basi attenuata, margine angustissimo, reflexo, denticulato, suprâ nuda, polita, viridia, subtùs lanâ implexâ copiosissimâ niveâ tecta, spithamæa, 2 pollices lata. *Involucra* fæminea sphæroidea: *squamis* ovato-lanceolatis, mucronatis, coriaceis, adpressè imbricatis, extùs lanuginosis, obsoletè striatis. *Flosculi* fæminei filiformes, limbo 5-fidi: *laciniis* linearibus, obtusiusculis. *Stigmata* lingulata, lævia. *Pappi* radiis copiosissimis, fasciculatis, apice penicillatis, cinereis.

—————
MUTISIA, p. 260.

I ought to have added to my remarks on this genus, that the presence or absence of appendages to the scales of the involucre is a character of specific importance only, as may be seen by a comparison of species intimately allied, such as *Mutisia Clematis* and *lanata*, and *M. inflexa* and *linearifolia*. I have now satisfied myself as to the identity of *Mutisia grandiflora* of Humboldt and Bonpland with the *M. Clematis* of Linnæus, from
the

the examination of another and very perfect specimen of the latter species in the herbarium of the younger Linnæus, now incorporated with that of Sir James Edward Smith.

MUTISIA *viciæfolia*, p. 363.

The specimens of this species from Cavanilles, as well as from Ruiz and Pavon, are marked on the tickets as being from Peru; but it is stated by Cavanilles in his *Icones Plantarum* as being a native of Valparaiso in Chile. Should this information be incorrect, and the plant really proves to be a native of Peru, the *M. acuminata* may ultimately be found to be identically the same.

MUTISIA *sinuata*, p. 267.

From a specimen I have received from Dr. Gillies, the rays appear to be of a pale pink, almost approaching to white. The broad base of the leaves, and the continuous wings of the stalks, sufficiently distinguish the *M. subspinosà* from this species, although Dr. Hooker has proposed to unite them. I have also a specimen of this from Dr. Gillies. The rays of *M. subspinosà* are bright yellow.

MUTISIA *linearifolia*, p. 272.

I have specimens of this from Dr. Gillies marked *M. rigida*, *nov. sp.?* The exterior scales of the involucre are furnished with a short abrupt point. I have referred to this species, with a mark of doubt, the *M. linifolia* of Dr. Hooker, which, however, turns out to be a very distinct plant, as I find from examining a specimen with which Dr. Gillies has favoured me. It differs in having perfectly flat leaves, and the exterior scales of the involucre ovate and acuminate.

[The continuation of this Paper will be given in a future Part of the Transactions.]

1830. "ADDITIONS AND CORRECTIONS." *Transactions of the Linnean Society of London* 16, 297–303. <https://doi.org/10.1111/j.1095-8339.1829.tb00137.x>.

View This Item Online: <https://www.biodiversitylibrary.org/item/19440>

DOI: <https://doi.org/10.1111/j.1095-8339.1829.tb00137.x>

Permalink: <https://www.biodiversitylibrary.org/partpdf/25013>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.