

Pseudophoenix ^{Nov. gen.} Hrn. Windl. *Gaußiae affinis*.

Fructus stipitatus drupaceus exsiformis aurantiacus, carpellis 1-3 globosis,
stigmatum residuis basilaribus vel in fructibus lobatis laterilibus vel centrolibus,
epicarpio coriaceo, mesocarpio grumoso, endocarpio tenuiter vitreo-crustaceo.
Semem liberum subglobosum erectum, hilo basilari, raphe adscendentem utriusque
ramis 2-3 manifeste curvatis, albumine aequabili; embryo basilaris.

Fl. ♀ in fructu: calyx parvus pateriformis leviter 3-denticulatus. Petala
3 ovata obtusa viridia refracta. Staminodia 6 manifeste apice atropurpureo.

— Palme mediocris, ercta. foliis pinnatisectis, segmentis duriusculis imae-
bri velda replicatis.

Species 1. P. Sargentii Hrn. Windl. Elliotts Key. Florida.

fern.



Engelmann, George. 1882. "Engelmann, George Apr. 30, 1882 [to S. Watson] [enclosed description]." *George Engelmann letters to Asa Gray*

View This Item Online: <https://www.biodiversitylibrary.org/item/227886>

Permalink: <https://www.biodiversitylibrary.org/partpdf/255873>

Holding Institution

Harvard University Botany Libraries

Sponsored by

Arcadia 19th Century Collections Digitization/Harvard Library

Copyright & Reuse

Copyright Status: Public domain. The Library considers that this work is no longer under copyright protection

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.