

*Descriptions of three new species of* TRICHOPTERYGIA, *found by the Rev. T. Blackburn in the Sandwich Islands; by the Rev. A. MATTHEWS, M.A.*

To receive Trichopterygia from the Sandwich Islands is a great event in Entomology, and the present species would lead one to suppose that the fauna of that region must be of an abnormal and highly interesting type, since they are all more or less aberrant from any previously known forms. They were all found by Mr. Blackburn in the neighbourhood of Honolulu, and were sent by him to Dr. Sharp, who very kindly presented them to me.

Mr. Blackburn deserves much credit for the care with which these insects had been mounted, their form and limbs being clearly displayed. I hope that he will persevere in his researches, and soon send home a fresh and more numerous collection.

*Actidium Sharpianum*, n. sp. Pl. II, fig. 2.

L.c.  $\frac{5}{16}$  l. = .63 mm. Elongatum valde convexum piceo-brunneum pilis brevibus argenteis vestitum; capite modico triangulare, oculis sat magnis prominentibus retro-positis; pronoto angusto cordiformi, antice latiori, confertim tuberculato, leviter marginato, ad basim in disco leviter transverse impresso; elytris oblongis lateribus vix rotundatis, capite atque pronoto parum latioribus plus quam sesqui longioribus, confertim sat leviter asperatis, apicibus latis fere rectis; pedibus atque antennis piceis.

*Head* elongate triangular, closely tuberculate; eyes large and prominent, placed far back; antennæ long, piceous.

*Thorax* small, heart shaped, widest before the middle, as long and rather wider than the head, closely tuberculate, with a faint oval transverse impression near the base, sides slightly margined, basal margin rounded.

*Scutellum* elongate, triangular, closely asperate.



*Elytra* elongate-oblong, rather wider and more than half as long again as the head and thorax, very convex, widest near the middle, sides narrowly margined and very little rounded, apex very broad and nearly straight.

*Abdomen* with the pygidium alone exposed.

*Legs* moderate, robust, piceous, with all the tibiæ strongly dilated.

*Under parts* castaneous, the posterior coxæ slightly laminate.

*Differs* from its congeners in its elongate convex form, shining surface, small size of the thorax, and abrupt truncation of the elytra.

Of this very distinct species, which forms an obvious connection between the shining and the pubescent species of *Actidium*, a single example was found by Mr. Blackburn on salt mud near Honolulu.

*Obs.* I do not think that the truncated apex of the elytra is sufficient to separate this species from the genus of *Actidium*. The truncation itself is not a true truncation in the common acceptance of the term, the elytra are not shortened, but merely terminate in a straight apex.

#### PTILIODES, n. gen.

*Corpus* elongatum depressum.

*Caput* porrectum oculis magnis, antennis 11-articulatis articulis duobus basalibus permagnis fere paribus, sex sequentibus inter se paribus brevibus plus minusve linearibus, nono precedentibus longiori parum incrassato, decimo magno valde incrassato antice truncato, undecimo magno valde incrassato ovato sat obtuso, duobus his ultimis coronâ mediali setarum perbrevium ornatis.

*Pronotum* plus minusve quadratum.

*Scutellum* triangulare.

*Elytra* valde abbreviata, dimidium abdominis vix tegentia.

*Alæ* amplæ sub elytris visæ.

*Abdomen* elongato-conicum longe exsertum.

*Subtus*—

*Caput* parte gulari magnâ convexâ, oculis permagnis valde prominentibus.



*Prothorax* sat longus, prosterno modico leviter falcato, receptaculis coxarum magnis circularibus contingentibus.

*Pectus mesosterno* modico, carinato carinâ antice productâ, episternis sat magnis triangularibus; receptaculis coxarum magnis carinâ divisis; *metasterno* magno ad latera extenso, inter coxas posteriores late producto margine apicali rectâ.

*Venter* conicus sat acuminatus segmentis sex compositus, segmento apicali parvo dentato vel emarginato.

*Pedes* elongati graciles.

*Coxæ* anteriores contingentæ; intermediae carinâ divisæ; posteriores magnæ robustæ longe distantes, vix aut non laminatæ.

*Ptiliodes Blackburni*, n. sp. Pl. II, fig. 3.

L.c.  $\frac{5}{16}$  l. = .63 mm. Oblongus sat elongatus, sat convexus castaneus pilis argenteis densissime vestitus, haud nitidus; capite brevi lato antice rotundato; pronoto tranverso, capite parum longiori et latiori, confertim tuberculato, ad basim latiori, angulis posterioribus fere rectis; elytris capite atque pronoto parum longioribus haud latioribus modice asperatis, lateribus fere parallelis, apicibus rotundatis; pedibus atque antennis brevioribus late flavis.

*Head* broad, rather short, much rounded in front, closely tuberculate, not shining; eyes moderate, scarcely prominent; antennæ short, robust, bright yellow.

*Thorax* transverse, quadrate, scarcely longer but rather wider than the head, widest at the base, pubescent, not shining, closely tuberculate, sides moderately rounded, posterior margin nearly straight, with the angles right angles.

*Scutellum* small, triangular, closely asperate.

*Elytra* oblong, not wider and scarcely longer than the head and thorax, with the sides nearly straight, finely and closely asperate, each apex separately and strongly rounded, pale, with the extreme edge white.

*Abdomen* moderately exserted.

*Legs* long, bright yellow.

*Under parts* castaneous, with the mouth and five apical segments of the abdomen yellow.

*Differs* from other species in its truly oblong form, castaneous colour and dense silvery pubescence.

*Habitat.* A single specimen of this distinct and pretty species



was found by Mr. Blackburn under the bark of a tree near Honolulu.

*Obs.* The genus *Ptiliodes* must be placed immediately before *Ptinella*, from which it may be distinguished by its oblong depressed form, quadrate thorax, and diverse formation of the antennæ. I have not been able to examine its anatomy any farther than a superficial view of the underside would permit.

*Ptinella Pacifica*, n. sp. Pl. II, fig. 4.

L.c.  $\frac{6-7}{16}$  l. = .75—.87 mm. Elongato-ovalis, sat depressa sat nitida, obscure-castanea pilis flavis sparse vestita; capite magno lato; pronoto capite sat latiori fere pariter longo, tuberculis parvis remotis equidistantibus, interstiisque nitidis leviter reticulatis per totum ornato, atque impressione indistinctâ in disco notato, lateribus marginatis valde rotundatis, ad media latoribus, angulis posterioribus sat obtusis; elytris capite atque pronoto sat longioribus vix latoribus, pone media latoribus, sat fortiter asperatis, apicibus latis valde rotundatis; pedibus atque antennis longis robustis late flavis.

*Head* rather short, much rounded in front, shining, closely reticulated, with a few distant minute tubercles; eyes small and prominent, antennæ long, robust, bright yellow.

*Thorax* moderate, as long but broader than the head, ornamented with minute remote and equidistant tubercles, with the interstices shining, finely and closely reticulate, sides margined and much rounded, with all the angles rather obtuse, there is a faint impression on the disc, which may have been accidental.

*Scutellum* moderate, triangular, rather long, deeply asperate.

*Elytra* rather long, as long and as wide as the head and thorax, widest behind the middle, rather deeply asperate in irregular transverse rows, with the interstices shining and finely reticulate, apices much rounded, with the extreme edge pale.

*Abdomen* with five segments exposed, apex obtuse and pale.

*Legs* long, bright yellow.

*Under parts* castaneous, with the mouth and apex of the abdomen yellow.

*Differs* from all its congeners in the large size and rounded form of the thorax.

Two examples of this species were found by Mr. Blackburn under the bark of a tree on the mountains near Honolulu.



Matthews, A. 1882. "Descriptions of three new species of Trichopterygia, found by the Rev. T. Blackburn in the Sandwich Islands." *Cistula entomologica* 3, 39–42.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/40224>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/26176>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Smithsonian

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.