
REPORT ON A COLLECTION OF JAPANESE CRANE-
FLIES (TIPULIDÆ), WITH A KEY TO THE SPECIES
OF PTYCHOPTERA.

BY CHARLES P. ALEXANDER, ITHACA, N.Y.

(Continued from Page 210.)

Gonomyia (Gonomyia) superba, sp. n.

Antennæ, brown; color, brown and yellow; vein, Sc ends slightly beyond the origin of Rs.

Male.—Length, 5–5.5 mm.; wing, 4.9 mm.

Female.—Length, 5.9 mm; wing, 5.2–5.5 mm.

Male.—Rostrum yellow, palpi brown; antennæ brown, including the basal segments; front, vertex and occiput dull yellow, the vertex clearer yellow behind.

Pronotum, clear light yellow above; on the sides, a short, dull brown stripe from the cervical sclerites down to above the fore coxa. Mesonotum, præscutum very light yellowish brown, with rich chestnut-brown stripes, a median stripe, broad and dark in front, narrow behind, and again enlarged at its end divided by a pale, narrow, median stripe; lateral stripes short, beginning behind the pseudosutural pits crossing the transverse suture and suffusing the lobes of the scutum; lateral edge of the præscutum, in front, yellowish; behind, brown; scutellum pale, whitish; the base and lateral edges tinged with brownish, post notum brown. Pleuræ clear yellowish white, an irregular dark brown mark behind and above the base of the coxa; sternum yellow, the sides of the mesosternum, between the fore and middle legs, brown, separated by a broad median pale mark; the propleural stripe begins on the prosternum as a rounded mark which sends out a narrow caudal prolongation. Halteres light yellow. Legs: coxæ and trochanters light yellow, margins of the segments more or less brown; femora and tibiæ light brown; tarsi somewhat darker brown. Wings, hyaline or nearly so; veins brown, costa more yellowish. Venation (see fig. 14, pl. III): Sc ending slightly beyond the origin of Rs; basal deflection of Cu¹ about at the fork of M.

Abdomen, tergum, light yellow, each segment with a large brown mark on basal half, the caudal margin of this mark much

rounded; sternum light yellow. Hypopygium (see fig. 1 and 2, plate X). Pleurites short and broad, the caudal end produced into one fleshy and three chitinized appendages, as follows: Viewed from above, a fleshy lobe in front, the inner dorsal margin produced entad and dorsad into a slightly curved slender spine; behind the fleshy lobe arises a stout hook, very strong at the base, constricted before the middle, the tip slender and pointed, this hook directed entad and caudad; from the outer ventral angle of the pleurite arises a long, straight chitinized appendage, directed entad and caudad, narrow basally and more enlarged apically. The guard of the penis is long, pale, ending in a long, slender, tube-like point. On either side of the penis guard arises an elongate, very slender, chitinized hook, which is straight for about three-fifths its length and then bent strongly inward; viewed from the side, these hoops are bent very strongly ventrad and then caudad. Summarized, the hypopygium bears eight chitinized slender arms, all except two (which are probably homologous with the second gonapophyses) being borne by the pleurites.

Female.—Very similar to the male, but larger.

Vial No. 1.—Tokio, Japan; Aug. 1912. One ♂.

Vial No. 5.—Nishigahara, Japan; Apr. 25, 1912; 5 ♂, 4 ♀.

Holotype, ♂; Vial No. 1.

Allotype, ♀; Vial No. 5.

Paratypes, 5 ♂, 3 ♀; Vial No. 5.

Types in author's collection; Paratypes in U.S. National Museum and Cornell University Collections.

G. superba differs from *nubeculosa* Meij. (Java). (Tijd. voor Entomol., vol. 44, p. 48, 49; fig. 36, 1911) in the unspotted wings; from *metatarsata*, (l.c., p. 48, fig. 35) in its closed cell 1st M₂, etc.

Gonomyia (Leiponeura) insulensis, sp. n.

Pleuræ without longitudinal stripes; vein Sc ends far before the origin of Rs.

Female.—Length, 3.9–4 mm.; abdomen, 2.6 mm.; wing, 4 mm.

Female.—Rostrum yellow, palpi brown; antennæ, segment one yellowish, remainder dark brown; front, vertex and occiput yellow, the vertex suffused with dark colored.

Mesonotal præscutum yellowish, with three brown stripes, the median one broad, not divided by a pale median vitta, extending to the suture, the lateral stripes are broad, narrow, uniform in width until they cross the suture (not expanded behind), lateral margin of the sclerite dull yellow, the ground color between the brown stripes is very reduced; scutum, lobes dark brown, median line yellowish; scutellum yellow, a brown median spot in front; postnotum brown. Pleuræ, mesopleuræ brown in front, extending from the lateral margin of the præscutum down to and suffusing the mesosternum on the sides; metasternum pale brown. Halteres dull yellow. Legs: coxæ and trochanters yellow, suffused with brown in front; femora, tibiæ and tarsi brown, a little darker toward the tip. Wings subhyaline, veins brown. Venation (see fig. 12, plate III); Sc. ending far before the origin of Rs; R^{2+3} almost parallel to R^1 .

Abdominal tergites yellowish-brown; sternites light yellow.

Vial No. F.—Tokio, Japan; August, 1912; 1 ♀.

Holotype, ♀; in Vial F.

Type in author's collection.

The three species of *Gonomyia* described by de Meijere as *Atarbæ* (Tijd. voor Entomol.; vol. 44, 1911) are all members of the subgenus *Leiponeura* Skuse. These species are *Gonomyia nebulosa* (l.c., p. 42, fig. 25); *pilifera* (l.c.; p. 43, fig. 26) and *diffusa* (l.c.; p. 43, 44). They have nothing in common with *Atarba* and are quite distinct from any members of the *Leiponeura* group, that I know of, in their clouded wings. *G. insulensis* differs from all of the above species in its unmarked wings.

Genus *Erioptera* Meigen.,

Subgenus *Acyphona* Osten-Sacken.

Of this subgenus, two species were included, both of which are herein characterized as new. The only described Palæarctic species, *Acyphona maculata* Meigen, of Europe, differs from the Japanese species, as follows: Wing pattern, in *maculata* large, rounded brown markings mostly with grey centers; the body-shade is much lighter in *maculata* and there are several important differences in hypopygial characters, these being shown by the following key:

1. 9th tergite broad and thin, at its apex deeply notched; two chitinized teeth at the base of the pleura on the ventral side.....2.

9th tergite provided with two chitinized hooks at its apex; no chitinized teeth at the base of the pleura on the ventral side; [horns of the second gonapophyses long, widely separated at the base] (Japan).....*asymmetrica*, sp. n.

2. Base of pleura on sternal side provided with a chitinized plate which is bidentate, the proximal tooth free, the distal one joined to the pleura; 2nd gonapophyses short, chitinized at tip and on sides; apex merely notched.

(Europe).....*maculata* Meigen.

Base of pleura on sternal side provided with a small chitinized tooth, minutely denticulate; 2nd gonapophyses long, the tips long and widely separated (Japan).....*incongruens*, sp. n.

Erioptera (Acyphona) incongruens, sp. n.

Small species; light brown, with narrow dark brown pleural stripes; wings thickly spotted with brown.

Male.—Length, 5 mm.

Male.—Rostrum and palpi brown. Antennæ long, segment one brownish-yellow; segments two to eight light yellow; remainder with increasing amounts of brown at their tips, the apical segments all brownish. Front, vertex and occiput dark brown.

Thoracic pronotum brownish-yellow, brown on the sides. Præscutum reddish-brown, with a double median brown stripe; humeral region brighter yellow; sides of the sclerite darkened; scutum, scutellum and postnotum brown. Pleuræ reddish-brown with narrow dark-brown lines, the most dorsal one continuing from behind the fore coxa underneath the wing to the postnotum; the second beginning on the mesosternum running above the middle coxa, becoming very narrow and indistinct before the root of the halter; the last stripe on the metasternum over the hind coxa. Halteres light yellow. Legs: coxa brown; trochanters brownish-yellow. (The legs are all detached and loose in the vials; most of these have the femora largely brown, basal third mostly paler, yellowish; a post median yellow ring, tip usually pale; tibiæ and

tarsi clear light yellow, sometimes infuscated at the tips; tibiae often with a sub-basal annulus. In the vial were several specimens of *E. asymmetrica*, a closely allied form, and most of the legs evidently belong to that species. Two legs in the vial are very different and may belong to this little species, this being rendered probable by the size; in these the entire legs are clear, light yellow, the femora with a rather narrow subapical dark brown ring).

Wings spotted with brown.

Abdomen: Tergum dull brownish yellow, apex and lateral margins of the sclerites brown. Hypopygium unsymmetrical as in the genus, the 9th abdominal segment being twisted one-half around. Suture between the 9th tergite and the 9th sternite not indicated. The 9th tergite is broad and long, its hind margin produced caudad in a wide, thin plate which is broadly and rather deeply notched at its middle; no chitinized hooks at its apex. The pleurites are convex outerly (produced into two apical appendages), the base (dorsal) produced entad and cephalad in a long, chitinized hook; the ventral edge of the pleura near the sternum possesses a small chitinized organ which is directed caudad and is provided with two or three denticulae; of the two apical appendages, the ventral one is chitinized, the dorsal one is fleshy, the second gonapophyses are close together, the chitinized tips rather long and deeply divided. (See plate X, figs. 5 and 6).

Holotype, ♂. Vial 6, April 25, 1912; Tokio, Japan.

Erioptera (Acyphona) asymmetrica, sp. n.

Resembles *incongruens* closely, but is larger, the coloration darker, especially on the pleurae and usually on the abdomen. Wings hyaline, spotted with brown, varying considerably in the intensity and size of the markings; in some the dots are small, not confluent, in the darker specimens the spots on the costal half of the wing tend to flow together to form large blotches. The male genitalia of the two species is remarkably different. (See plate III, fig. 15, wing.)

The hypopygium is, as in the genus, asymmetrical, the usual dorsal portions of the 9th sclerites being switched around on a level with the pleural sutures of the remaining segments. (See fig. 7-9, plate X), suture between 9th tergite and sternite obliterated, 9th

tergite broad and long with a cross-shaped mark; near its tip set with two small, semicircular, chitinized pieces which are produced into sharp points on the proximal ends. Pleurites short and stout, at the base on the dorsal side, produced into a long, slender, chitinized arm which is directed entad, two apical appendages, the more ventrad being chitinized, especially at the tips, the dorsal apical appendage fleshy. Between the tergite and the unarmed sternite, nearly in the median plate, is a rectangular, subchitinized organ, bearing at its outer angles chitinized hooks, bent ventrad and inward, these hooks minutely denticulated at tip.

♂.—Length, 5.8 mm.; wing, 6.3 mm.

♀.—Length, 6.4–7.1 mm.

Holotype.—Vial 6, April 25, 1912; Tokio, Japan.

Allotype.—Vial 6, April 25, 1912; Tokio, Japan.

Paratypes.—Vial 6 and L; 4 ♀, 2 ♂, April 25, 1912; Aug. 1912, Tokio, Japan.

Subgenus Erioptera, Meigen.

Erioptera (Erioptera) elegantula, sp. n.

Wings with brown spots.

Male.—Length, 5.4 mm.; wing, 7.7–7.9 mm.

Female.—Length, 6–6.5 mm.; wing, 7–8.3 mm.

Male.—Rostrum and palpi dark brown, antennæ with basal segments brown, flagellar segments short, dark brown; front, vertex and occiput dark brown.

Pronotum dark brown above, lighter colored on the sides. Mesonotum dark brown, the region before the pseudosutural pits more yellowish; scutum, scutellum and postnotum dark brown. Pleuræ dark brown. Halteres pale. Legs: coxæ dark brown; trochanters brown; femora dark brown; tibiæ dark brown, a little paler at the extreme base; tarsi dark brown. Wings subhyaline with greyish-brown marks, as follows: A large rounded spot at origin of Rs, a second at Sc², a third at end of Sc¹ running down over cross-vein r; a fourth spot at tip of R¹ and a smaller one at tip of R²; cord broadly margined with the same color; less distinct clouds at ends of the other veins and along most of these veins. Venation, (see fig. 3, plate III.)

Abdomen dark brown, densely clothed with long whitish hairs. Hypopygium. 9th tergite broad at base, narrowed at the middle,

the tip rather expanded with a deep V-shaped incision, the lobes rounded. Pleurites long, cylindrical, not very convex on outer face; three apical appendages, the more dorsal being somewhat fleshy, brown, elongate-cylindrical, narrowed basally, provided with long hairs, and, at its tip, with a slender hook directed cephalad; the median apical appendage is longest, chitinized, very strongly so at its tip; tip broadly expanded and concave, this concavity provided with minute denticulæ; the ventral apical appendage is shorter than the median one, fleshy, cylindrical, narrowed at base. Viewed from beneath, the 9th sternite is straight on its caudal margin, pleurites very broad at base, produced entad and almost meeting on the median line on the sternum; second gonapophyses long, slender, acicular, the tips barely projecting beyond the caudal level of the 9th sternite.

Female.—Similar, but averages larger in size.

Vial No. 1.—Tokio, Japan; 2 ♂, 2 ♀.

Vial No. 16.—Tokio, Japan; 2 ♀ (small, but apparently of the same species.)

Holotype.—♂. Vial No. 1, I.

Allotype.—♀. Vial No. I.

Paratypes.—1 ♂, 3 ♀, Vials I and 16.

Types in author's collection.

E. elegantula differs from *E. javensis* Meij. (Tijd voor Entomol., vol. 44, p. 45, 46, fig. 28, 1911) and *E. notata* Meij. (l.c., p. 46, figs. 29–31) in its spotted wings.

Genus Molophilus Curtis.

Molophilus pegasus, sp. n.

Antennæ of the male short; color of body brown.

Male.—Length, 4.2 mm.; wing, 4.3 mm.

Female.—Length, 4.9 mm.; wing, 5.1 mm.

Male.—Rostrum and palpi dark brown; antennæ light yellow, the flagellar segments with the exception of the first, a little more brownish; antennæ short, extending about to the base of the wings, segments of flagellum cylindrical; front, vertex and occiput brown.

Pronotum above, light yellow, darker on the sides. Mesonotal præscutum reddish-brown, with a broad, dark brown median stripe, and less distinct but broader lateral stripes, which begin

behind the pseudosuture, broaden out behind and fuse with the median stripe near the transverse suture; scutum, lobes brown, median line paler; scutellum lighter colored, yellowish medially, brown on the sides; postnotum brown. Pleuræ brown except dorsally, where there is a pale band running from the pronotum back to the wing basis. Halteres light yellow. Legs: coxæ and trochanters pale yellow, femora short, incrassated beyond the base, brown, paler basally; tibiæ and tarsi brown. Wings slightly tinged with yellowish-grey; veins yellow. Venation (see fig. 11, plate III).

Abdomen, tergites dark brown; sternites rather lighter brown, extreme apices of the sclerites pale. Hypopygium (see figs. 3 and 4, plate X); 9th tergite and sternite completely fused so that no pleural suture remains; viewed from beneath, the 9th sternite projects backward, its caudal margin rather squarely truncated; the outer ventral pleural arm is straight, fleshy, rather thickly covered with long hairs; just entad of the outer arm and nearer to the base of the pleurite, arises the inner ventral pleural arm, which is elongate, slender, its tip strongly chitinized and denticulated at the extreme end and bent inward; the guard of the penis is a pointed, chitinized organ, nearly as long as the outer pleural arm. Viewed from the side, outer ventral arm of the pleurite directed caudad; inner ventral arm with the tips conspicuously arcuated and bent ventrad; just above the base of the inner arm arises the dorsal pleural appendage, very broad at the base, its tip chitinized and directed slightly dorsad, on the dorsum of the pleurite are two protuberances clothed with long hairs. Viewed from above, the pleurites are very broad, so that the space between them on the median line is narrow; about midway of their length, on the inner face, is a strong protuberance, directed inward; it is strongly chitinized and almost touches its mate of the opposite side.

Female.—Similar, but larger; the abdomen is dark brown, the genital segment much brighter, yellowish-brown.

Vial No. 19.—Tokio, Japan; June 25, 1912; 1 ♀.

Vial No. 20.—Tokio, Japan; June 25, 1912; 1 ♀.

Vial No. K.—Tokio, Japan; Aug. 1912; 1 ♂.

Holotype.—1 ♂, Vial K.

Allotype.—1 ♀, Vial 20.

Paratype.—1 ♀ Vial 19.

Types in author's collection; paratype in U.S. National Museum collection.

M. pegasus differs from *bicolor* Meij. (Java) (Tijd. voor Entomol.; vol. 44, p. 45, fig. 27) in its darker brown body-color and darker legs.

Genus Conosia Van der Wulp.

Conosia irrorata Wiedmann.

The following papers since Kertész (1902) may be cited:

1904.—*Conosia irrorata* de Meij; Bijdragen tot de Dierkunde; p. 92.

1911.—*Conosia irrorata* de Meij; Tijdschrift voor Entomologie; vol. 44, p. 51.

1911.—*Conosia irrorata* Brun.; Rec. Indian Museum; vol. 6, part. 5, p. 283.

1912.—*Conosia irrorata* Brun.; Fauna Brit. India, Dipt. Nemat., p. 497.

One female in vial 47; Tokio, Japan. The wing pattern is figured on pl. III; fig. 13.

EXPLANATION OF PLATE X.

Fig. 1.—Hypopygium of *gonomyia superba*; dorsal aspect; x, y, z = chitinized pleural appendages.

Fig. 2.—Hypopygium of *gonomyia superba*; lateral aspect, sternum uppermost; lettering as in fig. 1.


Fig. 3.—Hypopygium of *Molophilus pegasus*; lateral aspect; t = 9th tergite; s = 9th sternite.

Fig. 4.—Hypopygium of *Molophilus pegasus*; dorsal aspect; p = pleura.

Fig. 5.—Hypopygium of *Erioptera (Acyphona) incongruens*, sp. n.; dorsal aspect.

Fig. 6.—Hypopygium of *Erioptera (Acyphona) incongruens*; 9th tergite, dorsal aspect.

Fig. 7.—Hypopygium of *Erioptera (Acyphona) asymmetrica*; 9th tergite, dorsal aspect.



EXPLANATION OF PLATE X.—Continued.

Fig. 8.—Hypopygium of *Erioptera (Acyphona) asymmetrica*; lateral aspect; p = pleura; s = 9th sternite; t = 9th tergite.

Fig. 9.—Hypopygium of *Erioptera (Acyphona) asymmetrica*; dorsal aspect; gonapophyse.

Fig. 10.—Hypopygium of *Limnophila japonica*; dorsal aspect; h = anal tube.

Fig. 11.—Hypopygium of *Limnophila satsuma*; ventral aspect.

Fig. 12.—Hypopygium of *Limnophila inconcussa*; dorsal aspect; h = anal tube; pl = pleura.

Fig. 13.—Hypopygium of *Liogma kuwanai*; lateral aspect; t = 9th tergite; pc = penis-guard.

Fig. 14.—Hypopygium of *Liogma kuwanai*; ventral aspect of the base of the tripartite penis-guard.

Fig. 15.—Hypopygium of *Liogma kuwanai*; dorsal aspect.

(TO BE CONTINUED.)

A NEW PYROMORPHID FROM TEXAS.

BY WM. BARNES, M.D., AND J. MCDUNNOUGH, PH.D., DECATUR, ILL.

Acoloithus novaricus, sp. nov.

Very similar to *falsarius* Clem., having the wings of the same dull black colour. The distinguishing feature is that the collar is unbroken reddish-orange, whereas in *falsarius* this colour is confined to the lateral areas, the centro-dorsal portion being black. Expanse, 14 mm.

Habitat: Kerrville, Texas; Shovel, Mt. Texas (July), 2 ♂'s. Type and cotype coll. Barnes. 4 ♂'s (Texas). Cotypes, Tring Museum, England.

Dr. K. Jordan, with whom we have recently had some correspondence concerning this group, has called our attention to this species and expressed the desire that we describe it. We take pleasure in doing so, as the characteristic feature seems very constant.

September, 1913



Alexander, Charles P. 1913. "Report on a collection of Japanese crane-flies (Tipulidae), with a key to the species of Ptychoptera [part]." *The Canadian entomologist* 45, 285–295.

View This Item Online: <https://www.biodiversitylibrary.org/item/22231>

Permalink: <https://www.biodiversitylibrary.org/partpdf/27102>

Holding Institution

MBLWHOI Library

Sponsored by

MBLWHOI Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.