

# Pollination and Pollinarium of *Dipodium punctatum* (Sm.) R. Br.

BY PETER BERNHARDT† AND PAMELA BURNS-BALOGH\*

*Dipodium punctatum*, the Hyacinth Orchid, is a common leafless saprophyte and one of the few orchid species found flowering over the summer months in Victoria (Willis 1978). Despite its wide distribution throughout the state and its large colourful flowers, virtually nothing is known of its floral biology. We present our recent observations in the hope it will encourage further field studies on this attractive plant.

A clump of six mature shoots of *D. punctatum* was found along a picnic trail off the Serra Road track in the Grampians State Forest in early January 1983. Five shoots had open flowers but one shoot remained in bud.

On 12/1/83 at 1:30 pm a native bee entered the site and was observed visiting the open flowers of *D. punctatum*. The bee visited the flowers on three stalks before it was captured. This insect was identified as a female, *Chalicodoma* (*Hackeriapis*) *derelicta* Cockerell (Megachilidae) by Ms Judith King, of the Department of Entomology at the Univ. of Queensland at Santa Lucia. The bee carried a viscidium of *D. punctatum* just above its antennae but below the middle ocellus (Figure 1). However, the pollen of other species were found on the bee's body. Stylidiaceae pollen (probably *Stylidium graminifolium* Swart, as it was still in flower in the Grampians) was densely deposited between the juncture of the head and the thorax (Figures 1 and 2). The pollen of *Eucalyptus* sp. and *Bursaria spinosa* Cav. were found in the scopae of the hind legs and on the

underside of the abdomen (Figure 3). Although some Australian orchids offer food hairs or a "pseudopollen" of sloughed petal cells (Jones 1981) none were found on the body of *C. derelicta*.

This is the first time that a member of the genus *Chalicodoma* has been observed visiting orchid flowers and carrying a viscidium. *Chalicodoma derelicta* has been recorded previously on the bushpea genera *Jacksonia* and *Dilwynia* (Papilionaceae) and on *Wahlenbergia* (Campanulaceae) by Ms J. King. Armstrong (1979) reviews observations of *Chalicodoma* species foraging on the flowers of Myrtaceae, Papilionaceae and Proteaceae.

The second author examined the pollen masses (pollinarium) with a Scanning Electron Microscope at the Smithsonian Institute in Washington, D.C. Each pollinarium was composed of four structures (fig. 4). There were four *pollinia* fused in two sets. The pollen grains within each *pollinia* were clustered in groups of four (tetrads). Each tetrad was united to other tetrads by a common wall to form hard, compact masses. There were two *caudicles*, one for each *pollinia* set, that served to attach the *pollinia* sets to the stipe. The *caudicles* consisted of elastoviscin produced by degenerative, sporogenous cells within the anther. The *stipe* was two branched (one branch for each *caudicle*) which served to attach the viscidium to the *caudicle*. Unlike these *caudicles*, the *stipe* was cellular. Finally, the *viscidium* ("sticky pad") adhered to the bee after the pollinarium was removed from the anther. The viscidium was shoe-like, rounded at the apex, blunt at its base and cellular like its *stipe* (Figure 5).

The four *pollinia* and their two

† Plant Cell Biology Research Centre, C/o School of Botany, Univ. of Melbourne, Parkville, Vic. 3052.

\* Dept. of Botany, NHB 166, Smithsonian Institution, Washington, D.C. 20650.



caudicles were derived from the orchid flower's solitary anther. In contrast, the viscidium and its stipe were derived from the rostellum (Dressler 1981).

Although *D. punctatum* is a terrestrial orchid its pollinia had little in common with the majority of Australian ground orchids placed in the tribe, Diuridae (*sensu* Dressler 1981). For example, the pollinia of *Pterostylis*, *Diuris* and *Thelymitra* are soft and granulate. Their tetrads are threaded together by cohesion strands and not by common walls. *Caladenia*, *Eriochilus* and *Glossodia* produce granulate pollen too but these grains are separate and never united as tetrads. The hard pollinia of *D. punctatum* are composed of tetrads united by common walls and, therefore, most closely resembled members of the tropical-subtropical Tribe, Vandeeae, in

the subtribe, Cyrtopodiinae, such as *Eulophia* and *Cymbidium* (Dressler 1981).

This story did not have a happy ending. It was the intention of the first author to return to the Serra Road site over a seven day period to collect more insects that may have pollinated *D. punctatum*. However, the following day the author found that someone had found the plants and picked every flowering stalk, uprooting two!

#### Acknowledgements

The authors would like to thank Dr Susan Duigan for analysis of the Stylidiaceae pollen and Ms J. King for identifying the bee. Funding was provided by the Australian Department of Education (CPPER). This article is dedicated to the memory of Mrs Edith Coleman whose observations on pollination remain the touchstone for those interested in Australian terrestrial orchids.




Fig. 1. The head of *Chalicodoma derelicta* bearing the viscidium of *Dipodium punctatum* and the pollen of a member of the Stylidiaceae  $\times 15$ . A = antenna; O = middle ocellus; SY = Stylidiaceae pollen; V = viscidium.





Fig. 2. Pollen grain of Stylidiaceae  $\times 350$ . Removed from the juncture of the head and thorax.




Fig. 3. Pollen of *Eucalyptus* sp. (E) and *Bursaria spinosa* (B)  $\times 192$  taken from the scopal hairs of the hind legs.

#### REFERENCES

- Armstrong, J. A., (1979). Biotic pollination mechanisms in the Australian flora — a review. *New Zealand J. of Bot.* 17: 467-508.
- Dressler, R. L. (1981). *The Orchids*. Harvard Univ. Press, Cambridge, Massachusetts and London, England.




Fig. 4. Scanning Electron Micrograph of the pollinarium of *D. punctatum*  $\times 21$ . C = caudicle; P = two pollinia composing one set; ST = stipe; V = viscidium.



Fig. 5. SEM of the base of the pollinarium showing how the branches of the stipe (ST) emerge from the shoe-like viscidium (V)  $\times 55$ .

- Jones, D. L. (1981). The pollination of selected Australian orchids. 40-43, *In*, Proceedings of The Orchid Symposium Held As a Satellite Function of the 13th International Botanical Congress, Sydney, Australia 1981. Editors Lawler, L. and Kerr, R. D. Harbour Press, Sydney.
- Willis, J. H. (1978). *A Handbook to Plants in Victoria. I. Monocotyledones*. Melbourne Univ. Press.

## Australian Natural History Medallion Fund

Amount on hand January 1983, \$1,958.50  
Mrs Ellen Lyndon, 3rd Donation, \$24.00



Bernhardt, Peter. and Burns-Balogh, Pamela. 1983. "Pollination and pollinarium of *Dipodium punctatum* R.Br." *The Victorian Naturalist* 100, 197–199.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/123451>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/271575>

**Holding Institution**

Field Naturalists Club of Victoria

**Sponsored by**

Atlas of Living Australia

**Copyright & Reuse**

Copyright Status: Permissions to digitize granted by rights holder.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.