

**Tetrastichus doteni**, new species.

*Female*.—Length about 2 mm. Dark olive green, shining; the face collapsing after death; antennæ brown, joints of funicle subequal in length, the first about as long as the pedicel; mesoscutum finely lineolated, the lines mostly longitudinal; scutellum with similar but much finer sculpture; parapsidal furrows very deep; median lobe of mesoscutum with setigerous punctures, arranged in three more or less distinct rows along each lateral margin, the outer line near the parapsidal furrow; disk of metanotum finely reticulately lineolated; median groove of mesoscutum distinct; the pair of grooves on the scutellum more distinct; median carina of propodeum developed into a broadened plate, widening behind, its apex being the median portion of the apical carina of the propodeum; on each side of this plate the surface is sculptured to resemble faintly developed, thimble-like punctures; prepectus lineolated; mesopleuræ in part faintly lineolated, the rest smooth, polished; metapleuræ with sculpture similar to that on the propodeum, but stronger; marginal vein not much longer than the submarginal, stigmal less than half as long as marginal; coxæ and femora greenish, tibiæ bark brown, with more or less metallic reflections; knees, tips of tibiæ, and the tarsi testaceous; last joint of tarsi dark; abdomen longer than the head and thorax combined.

*Male*.—Length about 1.25 mm. Similar to the female, but the tibiæ with more testaceous; joints 2-4 of funicle subequal in length, the first shorter; pedicel about as long as the second joint of funicle; sculpture of propodeum more indistinct than in female.

*Habitat*: Reno, Nevada.

*Type*: Cat No. 14362, U. S. National Museum.

From a series reared by Mr. S. B. Doten a type female and a type male and five paratypes of each sex have been selected. Mr. Doten writes that in cages the larvæ of this parasite feed readily on *Habrobracon hebetor* and on *Meraporus*, and occasionally on *Pteromalus*.

---

**A PREOCCUPIED NAME IN SPHECOIDEA.**
**Notogonidea**, new name.

*Notogonia* Costa, Ann. mus. zool. Napoli vol. 4, 1867 (1864) p. 82; not *Notogonia* Perty, Mitth. Naturf. Ges. Bern, 1850, p. 20.

*Type*: (*Tachytes nigra* Van de Linden) = *Larra pompiliiformis* Panzer.

S. A. ROHWER.



Rohwer, S. A. 1911. "A preoccupied name in Sphecoidea." *Proceedings of the Entomological Society of Washington* 13, 234.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/20330>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/271790>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Smithsonian

**Copyright & Reuse**

Copyright Status: NOT\_IN\_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.