

38. *Viola rugulosa* Greene.

*Viola rugulosa* Greene Pitt. 5: 26. 1902.

*Viola Rydbergii* Greene Pitt. 5: 27. 1902.

Often widely spreading from long thick branching stolons; stems stout, 4-6 dm. high; radical leaves usually 3-8, long-petioled, the blades cordate-ovate, abruptly short-acuminate, often 10 cm. wide, densely hirsutulous beneath, sparsely so along the veins above; lower stem leaves similar, the upper successively smaller and on shorter petioles, the blades becoming ovate-acuminate, the pubescence of the lower surface extending along the petioles and upper part of the stem; stipules lanceolate nearly entire; sepals linear-lanceolate; petals as in *V. canadensis*; capsules ovoid, seeds dark brown. petaliferous flowers often appearing in summer and autumn.

Rich wood lands, Iowa, Minnesota and west to the Rocky Mountains. May-Oct.

[From M.S. of *Viola* for Britton II. Fl. new edition.]



Brainerd, Ezra. 1885. "Brainerd, Ezra undated [notes]." *Walter Deane correspondence*

**View This Item Online:** <https://www.biodiversitylibrary.org/item/164242>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/288586>

**Holding Institution**

Harvard University Botany Libraries

**Sponsored by**

IMLS

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.