EUSTACHYS RETUSA (POACEAE), THE FIRST REPORT IN FLORIDA AND
A KEY TO *EUSTACHYS* IN FLORIDA

Stanley D. Jones & J.K. Wipff

S.M. Tracy Herbarium (TAES), Department of Rangeland Ecology and
Management, Texas A&M University, College Station, Texas 77843-2126
U.S.A.

ABSTRACT

Eustachys retusa (Lag.) Kunth, Chlorideae:Poaceae, previously un-
reported in Florida has been found in the southern coastal plain of the
panhandle in Okaloosa County.

KEY WORDS: *Eustachys*, *Eustachys retusa*, tribe Chlorideae,
Poaceae, Florida

Eustachys Desv. in Nouv. is composed of ten species, principally of the New
World tropical savannas (Clayton & Renvoize 1986). Six species of *Eustachys*
are now found in Florida: *Eustachys distichophylla* (Lag.) Nees, *E. floridana*
Chapm., *E. glauca* Chapm., *E. neglecta* (Nash) Nash, *E. petraea* (Swartz)
Desv., and *E. retusa* (Lag.) Kunth. Long & Lakela (1971), Wunderlin (1982),
Clewell (1985), nor Anderson (1984, 1986, 1988) have listed *E. retusa* as occur-
ring in Florida. It has been recorded in Georgia, New York, South Carolina,
and Texas (McKenzie *et al.* 1987). *Eustachys retusa* can be confused with
small or depauperate specimens of *E. distichophylla*. *Eustachys distichophylla*
is usually a much larger and more robust plant with more numerous panicle
branches, much longer, flexuous spikes, and upper, acute, sterile florets. The
following key will separate the taxa of *Eustachys* found in Florida (modified
from McKenzie *et al.* 1987).

KEY TO *EUSTACHYS* IN FLORIDA

1. Lateral veins of the fertile lemma glabrous. *E. glauca*
- 1' Lateral veins of the fertile lemma pubescent. 2

- 2.(1') Keel of fertile lemma glabrous. 3
- 3.(2.) Spikelets shorter than 2.1 mm; sterile floret widely cuneate, truncate. *E. retusa*
- 3' Spikelets 2.4 mm or longer; sterile floret oblanceolate, acute. .
E. distichophylla
- 2' Keel of fertile lemma ciliate. 4
- 4.(2') Spikelets shorter than 2.5 mm. *E. petraea*
- 4' Spikelets longer than 2.5 mm. 5
- 5.(4') Spike usually 1-3; spikelets 3 mm or longer; fertile lemma awn 0.6 mm or shorter. *E. floridana*
- 5' Spikes usually 4-9; spikelets 3 mm or shorter; fertile lemma awn 0.6 mm or longer. *E. neglecta*

Specimens collected: UNITED STATES. Florida: Okaloosa Co.: 12 July 1991, J.K. Wipff 2108 & S.D. Jones (FLAS,GA,TAES,TEX,SWSL,VDB,US,USF). Northwest corner of Leonard Burnes Road and U.S. 90, between U.S. 90 and the railroad tracks; 10.8 miles (17.3 kilometers) NE on U.S. 90 from its jct. with FL 87, NE of Milton. This taxon was abundant along U.S. 90 in an open disturbed roadside with the soils of the Lakeland-Troup-Alpin association. The site is nearly level at about 200 feet (61 m) elevation with acid sand throughout and a very thick sandy surface layer over a loamy subsoil. The geology of the site is of the Citronelle (Pc) formation; Pliocene series (mid-upper Miocene). Associated taxa include *Paspalum notatum* Flüggé, *Cynodon dactylon* (L.) Pers., *Aristida* sp., *Eragrostis* spp., *Richardia brasiliensis* Gomes., *Senna* sp., *Rubus* sp., *Ipomoea quamoclit* L., and *I. coccinea* L.

ACKNOWLEDGMENTS

We thank Larry E. Brown (SBSC), Gretchen D. Jones, and Robert I. Lonard (PAUH) for their reviews of this manuscript.

LITERATURE CITED

- Anderson, L.C. 1984. Noteworthy plants from north Florida. *Sida* 10:295-297.
- . 1986. Noteworthy plants from north Florida II. *Sida* 11:379-384.
- . 1988. Noteworthy plants from north Florida III. *Sida* 13:93-100.

- Clayton, W.D. & S.A. Renvoize. 1986. *Genera Gramineae Grasses of the World*. Her Majesty's Stationery Office, London, England.
- Clewell, A.F. 1985. *Guide to the Vascular Plants of the Florida Panhandle*. Florida State University Press, Tallahassee, Florida.
- Long, R.W. & O. Lakela. 1971. *A Flora of Tropical Florida*. University of Miami Press, Coral Gables, Florida.
- McKenzie, P.M., L.E. Urbatsch, & C. Aulbach-Smith. 1987. *Eustachys caribaea* (Poaceae), a species new to the United States and a key to *Eustachys* in the United States. *Sida* 12(1):227-232.
- Wunderlin, R.P. 1982. *Guide to the Vascular Plants of Central Florida*. University Presses of Florida, Tampa, Florida.

Jones, S D and Wipff, J. K. 1992. "Eustachys retusa (Poaceae), the first report in Florida and a key to Eustachys in Florida." *Phytologia* 73, 274-276.

<https://doi.org/10.5962/bhl.part.29314>.

View This Item Online: <https://www.biodiversitylibrary.org/item/48977>

DOI: <https://doi.org/10.5962/bhl.part.29314>

Permalink: <https://www.biodiversitylibrary.org/partpdf/29314>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.