BOTANICAL MUSEUM LEAFLETS

HARVARD UNIVERSITY

CAMBRIDGE, MASSACHUSETTS, AUGUST 8, 1939

Vol. 7, No. 9 ✓

ORCHID STUDIES, X

BY

LOUIS O. WILLIAMS

ORCHIDS FROM AMERICA AND ASIA

Altensteinia Rosei (*Ames*) *L. O. Williams* comb.
nov.

Aa Rosei Ames in Proc. Biol. Soc. Wash. 35 (1922)
81.

The genus *Aa* Reichenbach filius is referable to the
older *Altensteinia* Reichenbach filius. This species has
not been previously transferred.

Prescottia tubulosa (*Lindl.*) *L. O. Williams*
comb. nov.

Cranichis tubulosa Lindley Gen. & Sp. Orch. Pl.
(1840) 451—Kränzlin in Arkiv för Botanik 16 (1920)
1.

Prescottia pachyrhiza Richard & Galeotti in Ann.
Sci. Nat. ser. 3, 3 (1845) 31.

? *Prescottia Lindeniana* Richard & Galeotti in Ann.
Sci. Nat. ser. 3, 3 (1845) 31.

Galeoglossum prescottiioides Richard & Galeotti in
Ann. Sci. Nat. ser. 3, 3 (1845) 31.

Prescottia Galeottii Reichenbach filius in Linnaea
19 (1847) 377.

A photograph of the type of *Cranichis tubulosa* and
of Lindley's analytical sketch of the species leaves no

doubt that it belongs to the genus *Prescottia* and that it is the same as the plant commonly determined as *Prescottia pachyrhiza* Richard & Galeotti.

The record of *Prescottia Lindeniana* which is available seems to indicate that the species is referable to *P. tubulosa*, but the record is not satisfactory for certain identification.

***Cheirostylis longiflora* (Reichb.f.) L. O. Williams**
comb. nov.

Anecochilus longiflorus Reichenbach filius in Seemann Fl. Vit. (1868) 294.

Odontochilus longiflorus Benthams & Hooker filius ex Drake Ill. Fl. Ins. Pacif. (1886) 312—L. O. Williams in Bot. Mus. Leaflet Harv. Univ. (Orch. Fiji) 5 (1938) 112.

? *Odontochilus upoluensis* Kränzlin in Mitteil. Instit. Allgem. Bot. Hamb. 5 (1922) 236.

In a restudy of the generic affinity of this plant it appears that it should have been placed in the genus *Cheirostylis* because of the connate bases of the sepals and because of the form of the column.

FIJI ISLANDS: terrestrial, dense forest, southern slope of Mount Seatura, Mbua, Vanua Levu, at 600-700 meters altitude, flowers white, April 27, 28, 1934, *Smith 1623*; terrestrial, swampy rain forest, central plateau between Wainimala and Singatoka Rivers, Wainisavulevu Divide, Taunaisali, Tholo East, Viti Levu, at 3800 feet altitude, flower white, August 18, 1937, *St. John 18372*; "Feejee Islands", 1832-42, *Wilkes U. S. South Pacific Exploring Expedition*.

***Physosiphon Lansbergii* (Reichb.f.) L. O. Williams**
comb. nov.

Masdevallia Lansbergii Reichenbach filius in Nederl. Kruidk. Arch. 4 (1859) 317—Kränzlin in Fedde Repert. Beihefte 34 (1925) 104.

Physosiphon Lansbergii exhibits the generic charac-

ters found in *P. tubatus* (Lodd.) Lindley, the type species of the genus *Physosiphon*.

VENEZUELA: Prope coloniam Tovar, 1854-5, *Fendler 1369* (Gray Herbarium); epiphytic on palm trunks in humid forest, El Portachuelo between Maracay and Ocumare, Aragua, flowers yellow, May 8, 1925, *Pittier 11820* (in Pittier's Herbarium, United States National Herbarium, record in Ames Herbarium); epifita, selvas pluviales de Rancho Grande (Parque Nacional), Aragua, 1300 metros, 11 Abril 1937, *Pittier 13966* (Pittier's Herbarium, record in Ames Herbarium).

Epidendrum Deamii Schlechter in *Beihefte Bot. Centralbl.* 36, Abt. 2 (1918) 402.

Epidendrum tessellatum Bateman ex Lindley in *Bot. Reg.* 24 (1838) Misc. p. 7, non Roxb. (1795).

VENEZUELA: in crotch of *Protium heptaphyllum*, Rancho Grande, Aragua, at 800 meters altitude, May 26, 1938, *Williams 10131*.

The above collection, which was sent by Professor H. Pittier, seems to be the first record of the species in South America.

***Epidendrum oncidioides Lindley* var. *gravidum* (Lindl.) Ames, Hubbard & Schweinfurth** in *Bot. Mus. Leaf. Harv. Univ.* 3 (1935) 104.

Epidendrum gravidum Lindley in *Journ. Hort. Soc.* 4 (1849) 114—Ames in *Sched. Orch.* 4 (1923) 42, t. 2.

Encyclia gravis Schlechter in *Beihefte Bot. Centralbl.* 26, Abt. 2 (1918) 472.

Epidendrum argentinense Spegazzini in *Anales Mus. Hist. Nat. Buenos Aires* 28 (1916) 135, fig. 2—Hauman in *Anales Soc. Cien. Argentina* 90 (1921) 147.

Encyclia saltensis Hoehne in *Arq. Bot. Estado S. Paulo* n.s. 1 (1938) 19, t. 10.

The publication of *Encyclia saltensis*, by Dr. Hoehne, for a plant which I had been determining for my correspondents in Argentina as *Epidendrum argentinense* lead

to a study of the species which shows it to be one of the varieties of the widespread *Epidendrum oncidoides* Lindley.

Dr. Hoehne has apparently overlooked Spegazzini's publication of *Epidendrum argentinense* for otherwise it is difficult to understand how he could propose the plant as new under one of the generic segregates of *Epidendrum*.

***Pseuderia ramosa* L. O. Williams sp. nov.**

Herba epiphytica, scandens. Caule ramoso. Folia lanceolata, acuminata, multinervia, basi vaginis caulem obtegentes. Inflorescentia brevis, plusminusve quinqueflora; bracteae lanceolatae vel ovatae, obtusae vel acutae. Sepalum dorsale anguste ellipticum, acutum vel obtusum, tri- vel quinquenervium. Sepala lateralibus elliptica, acuta vel obtusa, arcuata, tri- vel quinquenervia. Petala anguste elliptica, obtusa, arcuata, tri- vel quinquenervia. Labellum obovatum, acutum, integrum, apice breviter pubescenti.

A scandent epiphytic herb of unknown size, the specimen at hand 3 dm. long. Stem coarse, branching, the main stem up to 8 mm. in diameter; the lateral branches smaller, mostly about 3 dm. long, the nodes 1-2 cm. apart. Leaves lanceolate, acuminate, thin when dry, 7-15 cm. long, 1.5-3 cm. broad, many-nerved; leaf-sheaths covering the stems. Inflorescence short, lateral, 3.5 cm. or less long, commonly rupturing the leaf-sheaths at about the middle, usually about 5-flowered; bracts lanceolate to ovate, obtuse or acute, 4-10 mm. long, 2-4 mm. broad. Dorsal sepal narrowly elliptic, acute or obtuse, about 12 mm. long and 3 mm. broad, 3- to 5-nerved. Lateral sepals elliptic, acute or obtuse, strongly arcuate, 10-12 mm. long, about 3 mm. broad, the apex somewhat callose-thickened, 3- to 5-nerved. Petals narrowly ellip-

tic, obtuse, arcuate, 3- to 5-nerved, about 10 mm. long and 2 mm. broad. Lip obovate, acute, simple, about 8 mm. long and 5 mm. broad, with one strong central, canaliculate, carinate callus extending from the base to beyond the middle, apex densely short-pubescent. Column of the genus.

Pseuderia ramosa seems to be most closely allied to *P. vanikorensis* Ames, a species which occurs in the Santa Cruz Islands. The following characters will serve to distinguish them.

Pseuderia ramosa

Lip obovate, about 8 mm. long.
Petals 3- to 5-nerved, about 10 mm. long.

Pseuderia vanikorensis

Lip elliptic, about 5 mm. long.
Petals 1-nerved, about 7.5 mm long.

Pseuderia ramosa is the first species of the genus to be reported from the Samoan Islands. It extends the range from the Fiji Islands, where two species of the genus are known, to Samoa which now constitutes the known eastern limit of the genus.

SAMOA: epiphytic on tree trunks, wet scrub-forest, top of Pioa, Island of Tutuila, flowers cream-white, at about 500 meters altitude, January 2, 1932, *Christophersen 3533* (TYPE in Herb. Bishop Museum, Honolulu; ISOTYPE in Herb. Ames No. 50309).

Dendrobium nanarauticolum *Fukuyama* in Tokyo Botanical Magazine 51 (1937) 900, fig.

CAROLINE ISLANDS, Ponape Island: Kaporujo, grows on tree trunks on the top of the mountain, rare, March 13, 1936, *Takamatsu 689* (in Herb. Bishop Museum and Herb. Ames).

This is a second collection of Professor Fukuyama's recently described species. *Dendrobium nanarauticolum* belongs to the group of the subgenus *Bolbodium* which is characterized by diphyllous pseudobulbs. Professor Fukuyama did not indicate to which species he con-

sidered the plant to be allied to nor have I been able to find one which is closely allied to it.

Glomera (Euglomera) **carolinensis** *L. O. Williams*
sp. nov.

Herba epiphytica, tenella. Radices filiformes, elongatae, glabrae. Caules leviter complanati, basi foliorum vaginis verrucosis obtecti. Folia linearia vel lineari-lanceolata, acuta, apice bilobata, paulo obliqua. Inflorescentia subcapitata; bracteae lanceolatae. Sepalum dorsale lineari-oblongum, obtusum vel acutum, plusminusve quinquenervium. Sepala lateralina connata, oblonga, basimentum saccatum formantia. Petala oblanceolata, obtusa, paulo obliqua, trinervia. Labellum saccatum, mento saccato circumdato; lamina late ovata, obtusa, paulo carinata.

A slender epiphytic herb up to 2 dm. tall. Roots filiform, elongated, glabrous. Stems somewhat flattened, covered by the persistent verrucose leaf-sheaths. Leaves erect, linear to linear-lanceolate, acute, apex unequally bilobed, slightly oblique, 3.5–7.5 cm. long, 4–8 mm. broad; persistent leaf-sheaths verrucose, especially above, about 1.3–2 cm. long. Inflorescence subcapitate, about 6-flowered, subtended and partly enclosed by a large, scarious, triangular bract; each flower subtended by a smaller lanceolate bract; flowers small for the genus. Dorsal sepal linear-oblong, obtuse or acutish, about 5-nerved, 6 mm. long and 1.5 mm. broad. Lateral sepals connate except about 1 mm. at the tips, forming a strongly saccate mentum at the base, the joined sepals oblong, about 5 mm. long and 3 mm. broad. Petals oblanceolate, obtuse, slightly oblique, 3-nerved, about 6 mm. long and 1.7 mm. broad toward the tip. Lip saccate, the sac contained within the saccate mentum; blade of the lip broadly ovate, obtuse, somewhat carinate, 1.5

mm. long, 1.2 mm. broad; sac oblong, 2 mm. long and 1.3 mm. broad. Column of the genus, about 1.5 mm. long.

Glomera carolinensis seems to be most closely allied to *G. bambusifolia* Schlechter, a New Guinean species. From this species, however, *G. carolinensis* may be distinguished by the comparatively narrow perianth parts and the different formation of the sac and the blade of the lip. *Glomera bambusifolia* is a much larger plant, up to 6.5 dm. tall, while *G. carolinensis* does not exceed 2 dm. in height.

Glomera carolinensis is the only species of the genus which has been reported from Micronesia. The nearest generic allies, in point of distance, would seem to be in New Guinea which is somewhat more than a thousand miles away.

CAROLINE ISLANDS, Ponape Island: Tolomail, grows on trees in open forest, February 11, 1936, *Takamatsu* 988 (TYPE in Herb. Bishop Museum, Honolulu; ISOTYPE in Herb. Ames No. 50281).

***Bulbophyllum Christophersenii* L. O. Williams**
sp. nov.

Herba epiphytica, repens. Pseudobulbi conferti, pyriformis, unifoliati. Folia oblanceolata, cuneata, obtusa, coriacea, petiolata. Inflorescentia uniflora, quam folia paulo brevior. Sepalum dorsale lanceolatum, acutum, basi trinervium. Sepala lateralibus lanceolata, acuta vel obtusa, basi trinervia. Petala linearia, acuta, uninervia. Labellum cordato-lanceolatum, obtusum, basi auriculatum, carinatum.

Epiphytic herbs from a repent rhizome. Pseudobulbs crowded on the rhizome, pyriform, 1–1.5 cm. long, up to 8 mm. in diameter, monophyllous. Leaves oblanceolate, cuneate, obtuse, coriaceous, petiolate, 6–18 cm. long, 1–2 cm. broad. Inflorescence 1-flowered, somewhat

shorter than the leaves, pedicel comparatively short, about 1–2 cm. long, the 6–8 cm. long pedicellate ovary appearing to be the pedicel. Dorsal sepal lanceolate, acute, about 12 mm. long and 4 mm. broad, 3-nerved at the base, several-nerved above. Lateral sepals lanceolate, acute or obtuse, 3-nerved at the base, several-nerved above, about 14 mm. long and 4 mm. broad. Petals linear, acute, 1-nerved, 3.5–4 mm. long, 0.5 mm. broad. Lip cordate-lanceolate, obtuse, auriculate at the base, thick and fleshy, about 4 mm. long and 2 mm. broad at the base. Column about 4 mm. long, with two terminal arms about 1 mm. long.

Bulbophyllum Christophersenii has no near allies which are known to me, certainly there are no known species in Polynesia which are allied to it.

SAMOA: epiphyte in open woodlands, east of Olo, Island of Savaii, flowers purple, at 700–800 meters altitude, August 8, 1931, *Christopherson & Hume* 2297 (TYPE in Herb. Bishop Museum, Honolulu; ISOTYPE in Herb. Ames No. 59298).

***Bulbophyllum Nagelii* L. O. Williams sp. nov.**

Herba epiphytica, repens. Pseudobulbi anguste-obpyriformes, unifoliati. Folia anguste-oblonga vel lanceolata, obtusa, coriacea. Inflorescentia densiflora, spicata; bracteae latae lanceolatae, acutae. Sepala lanceolata, acuminata, carinata, extus bracteolis parvis ornata. Petala lineari-lanceolata, acuta, uninervia, paulo serrulata. Labellum arcuatum, bicarinatum. Columna cum duobus stelidiis.

Repent epiphytic herbs. Stems slender, about 2–4 mm. thick. Pseudobulbs narrowly obpyriform, about 1.5–3.5 cm. long, 0.8–1.5 cm. thick, monophyllous. Leaves narrowly oblong to lanceolate, obtuse, abruptly contracted at the base, coriaceous, 7–12 cm. long, 1.4–2.3 cm. broad. Inflorescence densely flowered, spicate;

mostly about 3 cm. long; bracts narrowly lanceolate, acute, about 8 mm. long; rachis slightly thickened, depressed at the base of each flower. Flowers small, about equalling the subtending bracts in length. Sepals lanceolate, acuminate, carinate, 3-nerved, 6–7 mm. long, about 3 mm. broad at the base. Petals linear-lanceolate, acute, 1-nerved, minutely serrulate, about 4 mm. long and 1 mm. broad. Lip strongly arcuate, about 3 mm. long and 2–2.5 mm. broad, very fleshy except the lateral margins of the disc; with two carinae which merge into a single carina toward the apex. Bracteole opposite the petals, inconspicuous, about 0.5 mm. long. Column about 2 mm. long, with two lanceolate apical stelidia; column-foot prominent, about 2 mm. long.

Bulbophyllum Nagelii is not closely allied to any Mexican or Central American species of the genus but is perhaps closest to *B. pachyrachis* (A. Rich.) Grisebach, which it resembles somewhat in habit and in the structure of the flower. *Bulbophyllum Nagelii* may be distinguished from *B. pachyrachis* by its linear-lanceolate petals and from all known Mexican *Bulbophyllums* by the monophyllous pseudobulbs.

It is indeed a pleasure to name this plant for Mr. Otto Nagel. Mr. Nagel, who has collected in many remote regions of Mexico as well as the more accessible areas, has doubtless collected more specimens of the Mexican Orchidaceae than any other man. Regardless of the difficulties under which he has often worked, his specimens have always been superlatively good.

MEXICO: on oak trees in oak-pine forest, foot of Volcano Popocatepetl, north of Tetla del Volcán toward Santiago, longitude $98^{\circ}45'$ west, latitude $18^{\circ}54'$ north, at about 1800 meters altitude, July 12, 1932 and May 15, 1933, Nagel & Juan G. [onzáles] 1022 and 2228; pine and oak forest, southwest slope of the Volcano Popocatepetl, Tetla del Volcán, Morelos, longitude $98^{\circ}45'$ west, latitude $18^{\circ}54'$ north, at 1800 meters altitude, flowers dull green infused with

purplish-red, May 21, 1938, *Williams & Nagel 3864* (TYPE in Herb. Ames No. 50433).

Govenia liliacea (*La Llave & Lex.*) *Lindl.* var. ***Purpusii*** (*Schltr.*) *L. O. Williams comb. nov.*

Govenia Purpusii Schlechter in *Beihefte Bot. Centralbl.* 25, Abt. 2 (1918) 412; *Beihefte Fedde Repert.* 59 (1921) t. 61, fig. 241.

Govenia liliacea var. *Purpusii* resembles a small and immature *Govenia liliacea*, but all of the plants of the variety which I have seen seem to bear but a single leaf with the possible exception of one plant on the Conzatti and González specimens which may have two leaves as in the species. The structure of the flower is the same in the variety as in the species except that in the variety the apex of the lip tends to be plicate.

MEXICO: Cerro de San Felipe, Oaxaca, at 3000 meters altitude, May 22, 1898, *Conzatti & González 703*; Matrata, Vera Cruz, May 7, 1937, *Matuda 1285*; cool rocky places, Sierra de Tequila, Jalisco, July 5, 1893, *Pringle 4419*; Cerro Verde, Oaxaca, July 1908, *Purpus 3613*.¹

Vandopsis undulata (*Lindl.*) *J.J. Smith* in *Nat. Tijds. Ned.-Ind.* 72 (1912) 7.

Vanda undulata Lindley in *Journ. Linn. Soc. Bot.* 3 (1859) 42.

Stauroopsis undulata Benthams & Hooker filius in *Fl. Brit. Ind.* 6 (1890) 27—Hooker filius in *Ann. Roy. Bot. Gard., Calcutta* 5 (1895) 36, t. 55—King & Pantling in *Ann. Roy. Bot. Gard., Calcutta* 8 (1898) 205, t. 275.

¹ The original number cited by Schlechter in his description is 2613 but that is probably a typographical error as there is a drawing of the type in the Ames Herbarium, prepared by Schlechter, and it bears the number 3613.

Fieldia undulata Reichenbach filius Xenia Orch. 2 (1862) 38.

Stauroopsis polyantha W. W. Smith in Notes Roy. Bot. Gard., Edinburgh 13 (1921) 220.

Recently I have had the opportunity to examine authentic Chinese specimens of *Stauroopsis polyantha* W. W. Smith and am unable to distinguish it from *Vandopsis undulata*. W. W. Smith records that the flowers of the Chinese and Burmese plants are almost twice as large as those of the Indian plant. While the flowers of the Chinese plants, which I have seen, are slightly larger than those of the Indian plants the difference is not great and hardly of noteworthy character.

Taeniophyllum (Subg. Geissanthera) **Hosokawae** (Fukuyama) L. O. Williams comb. nov.

Microtorchis Hosokawae Fukuyama in Tokyo Botanical Magazine 51 (1937) 903, fig. 4-5.

The figures of this plant given by Fukuyama leave little doubt that the plant belongs to the subgenus *Geissanthera* (Schltr.) L. O. Williams of the genus *Taeniophyllum*. This would seem to be the fourth species belonging to this subgenus, the others are: *Taeniophyllum bracteatum* L. O. Williams, *T. papuanum* (Schltr.) L. O. Williams and *T. tubulosum* (J. J. Sm.) L. O. Williams.

Taeniophyllum Hosokawae is of further interest in that it is a considerable extension of the range of the subgenus.

MICRONESIA: Island of Ponape, *Hosokawa* 5956.

Taeniophyllum Parhamiae L. O. Williams sp. nov.

Herba parva epiphytica, acaulescens. Radices planae, uninerviae. Pedunculi breves, glabri vel verrucosi. Inflorescentia brevis, distichiflora. Sepalum dorsale lanceolatum, obtusum vel acutum, trinervium. Sepala lateralia

lanceolata, acuta, trinervia. Petala linearia vel lineari-oblonga, apice obtusa vel truncata, vix dentata, uninervia. Labellum saccatum; lamina anguste triangulari-subcordata, acuta, basi saccata, paulo scrotiformi.

Small acaulescent epiphytic herbs. Roots flattened, strongly 1-nerved, up to 25 cm. long or more. Peduncles short, glabrous but verrucose, at least when dry, up to 1 cm. long. Inflorescence probably 2-3-flowered, distichous; bracts acute, nearly 1 mm. long, semi-peltate, usually somewhat gibbous at the base. Perianth parts ocellate with crystalline inclusions, free to the base. Dorsal sepal about 3 mm. long and 1.2 mm. broad, lanceolate, obtuse or acute. Lateral sepals about 2.7 mm. long and 1.2 mm. broad, lanceolate, acute, 3-nerved, alate dorsally along the median nerve. Petals linear to linear-oblong, 1-nerved, apex obtuse or truncate, often dentate. Lip saccate; blade narrowly triangular-subcordate, acute, apical margins and the tip somewhat inrolled, about 2.5 mm. long and 1.5 mm. broad near the base; saccate base or spur of the lip slightly scrotoform, forming a 90° angle with the blade, about 2 mm. long and 1 mm. broad near the middle.

Taeniophyllum Parhamiae seems to be most closely allied to *T. decipiens* Schlechter, which occurs in the Samoan Islands, rather than to any species known to occur in Fiji.

It is with pleasure that this plant is named for Mrs. H. B. Richenda Parham and her two daughters, Miss Helena F. L. Parham and Miss Beatrice F. H. C. Parham, who have shown a keen interest in the native orchids of the Fiji Islands.

Fiji: epiphyte on Croton, Pender Street, Suva, June 26, 1938, *Parham 1*; in the old garden on breadfruit, Croton and coffee trees, Suva, August 24, 1938, *Parham 3* (TYPE in Herb. Ames No. 50094).

Williams, Louis O. 1939. "Orchid Studies, X: Orchids from America and Asia." *Botanical Museum leaflets, Harvard University* 7(9), 137–148.

<https://doi.org/10.5962/p.295135>.

View This Item Online: <https://www.biodiversitylibrary.org/item/31894>

DOI: <https://doi.org/10.5962/p.295135>

Permalink: <https://www.biodiversitylibrary.org/partpdf/295135>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.