LONCHIURUS BARBATUS.

CHARACTER GENERICUS.

Pinnæ Ventræles discretæ : *Caudalis* lanceolata.

Bloch. ichth. 10. p. 119.

CHARACTER SPECIFICUS.

LONCHIURUS cirris duobus mentalibus.

Bloch. ichth. 10. p. 120. t. 360.

Pisces Thoracici.

Surinamiam incolit *Lonchiurus barbatus*, in fluviis præcipue degens. Longior raro est decem unciis, colore plerumque fusco, quem plus minus saturatum habent diversa specimina.

Pubd by Nodder March 1799.

THE
BEARDED LONCHIURUS.

GENERIC CHARACTER.

Ventral Fins separate: *Caudal Fin* lanceolate.

SPECIFIC CHARACTER, &c.

Brown LONCHIURUS with two cirri or beards
under the chin.

The Lancet-Tail, or Bearded LONCHIURUS.

The bearded Lonchiurus is a native of Surinam, where it is generally found in rivers. It seldom exceeds the length of about ten inches, and is of a brown color, deeper or lighter in different individuals.

Shaw, George. 1799. "The Bearded Lonchurus, *Lonchurus barbatus* [Pl. 375]." *The Naturalist's Miscellany* 10(CXV), <https://doi.org/10.5962/p.304557>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276641>

DOI: <https://doi.org/10.5962/p.304557>

Permalink: <https://www.biodiversitylibrary.org/partpdf/304557>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.