Eight additional new records of Philippine Mosses

Virgilio C. Linis¹ & Benito C. Tan^{2,3}

¹*Philippine National Herbarium, Botany Division, National Museum, Ermita 1000, Manila, Philippines*

²*Herbarium, Singapore Botanic Gardens, 1 Cluny Road, Singapore 259569*

³*Department of Biological Sciences, National University of Singapore, Science Drive 4, Singapore 117546*

Abstract

Eight new taxa are added to the Philippine moss flora: *Acroporium rigens* (Broth. ex Dixon) Dixon, *Braunfelsia plicata* (Sande Lac.) Broth., *Brotherella fauriei* (Cardot) Broth., *Bryoerythrophyllum recurvirostrum* (Hedw.) P.C.Chen, *Gammiella pterogonoides* (Griff.) Broth., *Oxyrrhynchium hians* (Hedw.) Loeske, *Pterogonidium pulchellum* Hook. and *Vesicularia ferriei* (Cardot & Thér.) Broth. The genus *Pterogonidium* is new to the Philippine flora.

New records

Linis and Tan (2008), as well as subsequent publications by Linis (2010) and Linis and Tan (2010), estimate the total moss flora for the Philippines as 755 species in 238 genera and 56 families. However, continued study of the many unidentified bryological collections kept at the Philippine National Herbarium (PNH), including those of the senior author, has added eight new species and one generic record. These additions bring the total moss taxa known to occur in the Philippines to 763 species and 239 genera.

All voucher specimens of these Philippine mosses are held in PNH with some duplicates distributed to the Herbarium of Singapore Botanical Gardens (SING) for reference purposes.

1. *Acroporium rigens* (Broth. ex Dixon) Dixon

Sematophyllaceae

In the field typical plants of *Acroporium rigens* are tufted, with erect, acrocarpous-like branches which can often be mistaken for small plants of *Pyrrhobryum spiniforme* (Hedw.) Mitt. *Acroporium rigens* is most similar to *A. johannis-winkleri* Broth., with which it shares the bristly plant habit and similar erect-divergent, narrowly lanceolate-linear and tubulose leaves (Tan 1994, Tan *et al.* 2007). *Acroporium rigens*, however, is much larger with stems and branches ascending to 2 cm in height. Moreover, the rigid and setaceous leaves of *A. rigens* are longer than those of *A. johannis-winkleri* with their length reaching to 5 mm as compared to the latter with leaf length only to 2 mm.

Distribution: Malay Peninsula, Philippines, and Borneo (West Malesia) and Papua New Guinea (Western Melanesia) (Tan 1994, Tan *et al.* 2007).

Specimens examined: **PHILIPPINES:** Panay Island, Antique Province, Mount Baloy, on moist stone; 1,650 m, 9 Oct 1989, J.R. Cabalquinto 1229 & E. Sagcal, Jr. (PNH). **MALAYSIA:** State of Penang, Penang Hill, base of frond of *Cyathea*; 700 m, 13 Apr 1971, H.N. Ridley 5605 (SING). State of Johor, Gunung Belumut, on lower part of tree trunk in mossy forest; 914 m, May 1923, R.E. Holttum 10998 (SING).

2. *Braunfelsia plicata* (Sande Lac.) Broth.

Dicranaceae

Apart from its deeply plicate curved leaves, *B. plicata* is similar in size, habit, color and other morphological details to *B. dicranoides* (Dozy & Molk.) Broth. *Braunfelsia plicata* tends, however, to have a more sharply defined costa and blunter leaf apices.

Distribution: a West Malesian endemic recorded in Indonesia (Java, Sumatra) and the Malay Peninsula (Eddy 1988), now found in the Philippines.

Specimens examined: **PHILIPPINES:** Mindanao Island, Bukidnon Province, Malaybalay City, Barangay Busdi, Sitio Magantol, on tree trunk; transitional lowland forest, 1,220 m, 17 May 2008, V.C. Linis 3617-08 (PNH); *ibid.*, V.C. Linis 3624-08 (PNH). **MALAYSIA:** State of Kedah, Kedah Peak, in very mossy place, on ground, just below summit; 1158 m, 1 Apr 1925, R.E. Holttum 14869 (SING). Borneo, State of Sarawak, Mount Dulit, Dulit Range, on tree trunk in moss forest; 1300 m, 6 Oct 1932, P.W. Richards M2167 (SING).

3. *Brotherella fauriei* (Cardot) Broth.

Pylaisiadelphaceae

Brotherella was at one time combined with *Pylaisiadelpha* retaining the latter as the name of the merged genus (Buck 1984), but the move did not meet full acceptance (Jia & He 2006; Afonina *et al.* 2007). Ando *et al.* (1989) had clarified the distinction between *Brotherella* and *Pylaisiadelpha* and their differences include poorly differentiated leaf alar cells and the presence of filiform propagules in *Pylaisiadelpha*. Furthermore, Tan and Jia (1999) observed that the exothelial cell walls of *Pylaisiadelpha yokohamae* and *P. tenuirostris* exhibit weak collenchymatous thickening, quite unlike the uniformly, thick vertical walls with thin lateral walls in exothelial cells in species of *Brotherella*. Although recent phylogenetic studies using a limited number of gene sequences have shown the species of the two genera to be inseparable molecularly, we are keeping them apart based on their morphological distinction.

Recent molecular studies (Buck *et al.* 2000; Tsubota *et al.* 1999, 2000, 2001a, b, 2002) have shown that Sematophyllaceae *s.l.* includes two sister clades: the core sematophyllous taxa (e.g. *Acanthorrhynchium* Fleisch., *Acroporium* Mitt., *Meiothecium* Mitt., *Papillidiopsis* Buck & Tan, *Sematophyllum* Mitt. and *Trichosteleum* Mitt.), and a clade that includes *Brotherella* Loeske ex Fleisch., *Heterophyllum* (Schimp.) Kindb. in Fleisch., some species of *Hypnum* Hedw., *Isocradiella* Dix., *Platygyrium* Schimp., *Pylaisiadelpha* Cardot, *Taxithelium* Spruce ex Mitt. and *Wijkia* Crum. Tsubota *et al.* (2001a) called the latter clade the ‘*Brotherella* lineage’, while Goffinet and Buck (2004) described a new family Pylaisiadelphaceae for it. Although this clade lacks any obvious morphological synapomorphy, we follow Goffinet and Buck (2004) in placing *Brotherella* in the family Pylaisiadelphaceae.

Brotherella fauriei is another addition to the growing number of *Brotherella* species reported from the Philippines. Unlike other Philippine species of *Brotherella*, however, *B. fauriei* is a slender plant with distantly spaced branches. The mostly erect-patent stem and branch leaves with more or less falcate leaf acumina are distinctive.

Distribution: Temperate East Asia mainly in China and Japan (Tan & Jia 1999).

Specimens examined: **PHILIPPINES:** Luzon Island, Benguet Province, Mount Santo Tomas, on exposed roots of pine (*Pinus kesiya*) tree, 1,900 m, 29 Nov 1988, J.R. Cabalquinto 332, J.F. Barcelona & M.Y. Bangis (PNH). **CHINA:** Jiangxi Province, Zixi County, Mount Ma-Tou, on decaying log, 890 m, 26 May 2001, M.C. Ji 12668 (SING).

4. *Bryoerythrophyllum recurvirostrum* (Hedw.) P.C.Chen

Pottiaceae

Recent phylogenetic analysis of eight Russian *Bryoerythrophyllum* suggested the genus is monophyletic (Fedosov & Ignatova, 2008). In the Philippines, these *Barbula*-like plants can normally be identified in the field by the markedly rufous coloration of their older parts.

Among its congeners, *Bryoerythrophyllum recurvirostrum* is closest to *B. ferruginascens* (Stirt.) Giac. However, unlike the latter which is dioicous, *B. recurvirostrum* is synoicous and/or paroicous. *Bryoerythrophyllum recurvirostrum* also differs from *B. ferruginascens* by being bigger with stems reaching 3 cm compared to 17 mm. Furthermore, *B. recurvirostrum* has flexuose leaves up to 3 mm long with margins recurved nearly to apices, with few teeth near the tips vs leaves reaching 1.3 mm in length with entire margins recurved only up to mid-leaf. The lack of rhizoidal gemmae, which are present in *B. ferruginascens*, is also helpful in identifying *B. recurvirostrum*.

Distribution: almost cosmopolitan in both Northern and Southern Hemispheres, reaching Malesia. Recorded from the island of New Guinea and Indonesia (Sulawesi) (Eddy 1990).

Specimen examined: **PHILIPPINES:** Luzon Island, Benguet Province, Kabayan, Mount Pulag, on soil; 1,600 m, 18 Oct 1988, J.R. Cabalquinto 211 & E. Sagcal, Jr. (PNH).

5. *Gammiella pterogonioides* (Griff.) Broth.

Sematophyllaceae

This species is distinguished from its congeners by its relatively large size. The imbricate, cordate and ovate to oblong-lanceolate leaves with abruptly cuspidate leaf apices are also distinctive. This is the fourth species of *Gammiella* reported from the Philippines.

Distribution: Eastern India, Himalayas, Indochina and southwest China (Gangulee 1969–1980; Tan & Jia 1999) to the Philippines.

Specimen examined: PHILIPPINES: Luzon Island, Benguet Province, Baguio City, Wright Park, on bark of pine (*Pinus kesiya*) tree; 1,400 m, 27 Jan 1989, J.R. Cabalquinto 474 & B.O. van Zanten (PNH)

6. *Oxyrrhynchium hians* (Hedw.) Loeske

Brachytheciaceae

This is the second species of the genus reported in the Philippines and it is superficially very similar to *O. vagans* (Jaeg.) Ignatov & Huttunen in the field with both taxa possessing subjulaceous branch foliage. However, *O. hians* has shorter median laminal cells (35–105 µm) in its well-developed branch leaves compared to those of *O. vagans* measuring 70–145 µm (Ignatov *et al.* 2005). When fertile, *O. hians* has rough setae compared to those of *O. vagans* being smooth (Ignatov *et al.* 2005).

Distribution: widespread in warm temperate areas of North America, Europe, East Asia (China, Japan, the Philippines) and the Himalayas (India, Nepal and Bhutan) and also in Africa (Ignatov *et al.*, 2005).

Specimens examined: PHILIPPINES: Luzon Island, Benguet Province, Mount Santo Tomas, on rocky trail; 7,300 ft, 17 Apr 1956, N. Velaira 1 (PNH); Luzon Island, Nueva Vizcaya, 1 km. east of Dalton Pass, on fallen, rotting log near creek; 1,500 m, 4 Mar 1967, R.M. del Rosario 587 (PNH); Luzon Island, Camarines Sur Province, north slope of Mount Isarog, on stone; 16 May 1991, J.R. Cabalquinto 2192 (PNH). ROMANIA: Moldavia, Iasi, in opp. Iasi in loco nominato Copou, ad terram uliginosam; 80 m, 16 Mar 1916, C. Oescu s.n. (SING).

7. *Pterogonidium pulchellum* Hook.

Hypnaceae

This is the second report of this invasive moss species from the Neotropics in Asia. The first report was made by Tan & Buck (2002), when the authors discovered two populations on a tree trunk (*Melaleuca* sp.) inside the National Orchid Garden of the Singapore Botanic Garden. Re-examination of *P. pulchellum* using data from *rbcL* gene sequences showed that it has a close relationship with members of Hypnaceae (Chang *et al.* 2006).

Distribution: common in West Indies and Central America reaching northern South America (Sharp *et al.* 1994; Tan & Buck 2002), reported now in SE Asia from Singapore and the Philippines.

Specimens examined: PHILIPPINES: Luzon Island, Benguet Province, Baguio City, Barangay Pacdal, Baguio Botanical Garden, on lower trunk of an ornamental tree, near a human settlement area; 1,476 m, 22 Jan 2011, V.C. Linis 4754-11 (PNH). SINGAPORE: Botanical Gardens of Singapore, on *Melaleuca* tree trunk; 6 Nov 1998, B.C. Tan 98-472 (SINU).

8. *Vesicularia ferriei* (Cardot & Thér.) Broth.

Hypnaceae

Tan and Iwatsuki (1991) reported five taxa of genus *Vesicularia* (Müll.) Müll. for the Philippines. *Vesicularia ferriei* is somewhat similar to *V. montagnei* (Bél.) Broth. in having ovate dorsal leaves and suborbicular lateral leaves with short leaf apices. However, *V. ferriei* is readily distinguished by the broadly acute or obtuse leaf apices rather than shortly acuminate as in *V. montagnei*. Furthermore, the median laminal cells of both dorsal and lateral leaves in *V. ferriei* are linear-hexagonal or elongate-rectangular compared to oblong-hexagonal in *V. montagnei*.

Distribution: in temperate East Asian regions of Japan and China (Noguchi *et al.* 1994), also in the Philippines.

Specimens examined: PHILIPPINES: Mindanao Island, Cotabato Province, Kidapawan City, Barangay Ilomavis, Sitio Agco, on tree fern trunk along trail leading to waterfalls near small lake; 1,600 m, 29 Mar 2011, V.C. Linis 4900-11 (PNH). JAPAN: Honshu Island, Hiroshima-ken, Hiroshima-shi, Asa-kita-ku, Uga George, along Takayama River, on wet boulder by stream; 200 m, 26 Mar 1998, H. Deguchi s.n. (SINU). CHINA: in confinibus provinces of Kianghsi and Fukien, Mt. Dangwa-schan, inter Schitsheng and Ninghwa, in lacuna quadam in cacumine cum Glossadelpho; 1400 m, May 1921, T. H. Wang s.n. (SING).

Discussion

The distributional patterns exhibited by the eight newly reported Philippine moss taxa are relevant and useful to phytogeographical studies of the Philippine moss flora. The additions of *Brotherella fauriei*, *Oxyrrhynchium hians* and *Vesicularia ferriei* for the Philippine moss flora further fortify its predominantly East Asiatic

composition (Tan 1992, Linis & Tan 2008) while the discovery in the Philippines of *Acroporium rigens* and *Braunfelsia plicata* from Western Malesia and *Gammiella pterogonoides* from the Himalayan/ Indochina/ Hainan areas maintains the current relationships of moss floras between Philippines and these regions.

Bryoerythrophyllum recurvirostrum adds to the growing list of cosmopolitan taxa already recorded in the country. *Pterogonidium pulchellum* is the first record of an introduced moss taxon in the country.

Acknowledgment

The first author wishes to thank the Singapore Botanic Garden's Herbarium for a research fellowship to use its herbarium and library facilities.

References

- Afonina OM, Tsubota H & Ignatova EA (2007) The genus *Pylaisiadelphina* (Pylaisiadelphaceae, Musci) in Russia. *Arctoa* 16: 127–132.
- Ando H, Seki T & Schofield WB (1989) Generic distinctness of *Brotherella* from *Pylaisiadelphina* (Musci). *Bryologist* 92: 209–215.
- Buck WR (1984) *Pylaisiadelphina* replaces *Brotherella* (Sematophyllaceae). *Yushania* 1(2): 11–13.
- Buck WR, Goffinet B & Shaw AJ (2000) Testing morphological concepts of orders of pleurocarpous mosses (Bryophyta) using phylogenetic reconstructions based on *trnL-trnF* and *rps4* sequences. *Molecular Phylogenetics and Evolution* 16: 180–198.
- Chang Y, Khiong CW, Graham SW & Tan BC (2006) Molecular evidence for the systematic positions of two enigmatic mosses: *Pterogonidium pulchellum* (Sematophyllaceae, Musci) and *Piloecium pseudorufescens* (Myuriaceae, Musci). *Canadian Journal of Botany* 84: 501–507.
- Eddy A (1988) *A Handbook of Malesian Mosses. 1. Sphagnales to Dicranales*. (British Museum (Nat. Hist.); London)
- Eddy A (1990) *A Handbook of Malesian Mosses. 2. Leucobryaceae to Buxbaumiaceae*. British Museum (Nat. Hist.). London.
- Fedosov VE & Ignatova EA (2008) The genus *Bryoerythrophyllum* (Pottiaceae, Bryophyta) in Russia. *Arctoa* 17: 19–38.
- Gangulee A (1969–1980) *Mosses of Eastern India Fascicle 8 (Hypnobryales–Hypnaceae: Floristic trends: Index: Errata)*: 1753–2145. Calcutta.
- Goffinet B & Buck WR (2004) Systematics of the Bryophyta (mosses): from molecules to a revised classification. *Monographs in Systematic Botany from the Missouri Botanical Garden* 98: 205–239.
- Ignatov MS, Huttunen S & Koponen T (2005) Bryophyte flora of Hunan Province, China. 5. Brachytheciaceae (Musci), with an overview of *Eurhynchiadelphus* and *Rhynchostegiella* in SE Asia. *Acta Botanica Fennica* 178: 1–56.
- Jia Y & He X-J (2006) Notes on two species of *Brotherella* (Bryopsida: Sematophyllaceae) from Asia. *Journal of Bryology* 28: 268–277.
- Linis VC (2010) The mossflora of Camiguin island, Philippines and their floristic relations to some adjacent islands in the archipelago. *Telopea* 12: 525–542.
- Linis VC & Tan BC (2008) Progress of studies on phytogeography and biodiversity of the Philippine moss flora 1991 to 2006. Pp. 13–22 in Mohamed H, Baki BB, Nasrullag-Boyce A & Lee PKY (eds) *Bryology in the New Millennium*. Proceedings of World Conference of Bryology (University of Malaya: Kuala Lumpur)
- Linis VC & Tan BC (2010) Eleven new records of Philippine mosses. *Acta Bryolichenologica Asiatica* 3: 95–100.
- Noguchi A, Iwatsuki Z & Yamaguchi T (1994) *Illustrated Moss Flora of Japan, part 5*: 1013–1253 (Hattori Botanical Laboratory, Nichigan)
- Sharp AJ, Crum AP & Eckel M eds. (1994) *The Moss Flora of Mexico, part 2. Orthotrichales to Polytrichales*. Memoirs of the New York Botanical Garden 69.
- Tan BC (1994) The bryophytes of Sabah (North Borneo) with special reference to the BRYOTROP transect of Mount Kinabalu. XIX. The genus *Acroporium* (Sematophyllaceae, Musci) in Borneo, with notes on species of Java and the Philippines. *Willdenowia* 24: 255–294.
- Tan BC & Iwatsuki Z (1991) A new annotated Philippine moss checklist. *Harvard Papers in Botany* 3: 1–64.
- Tan BC & Buck WR (2002) *Pterogonidium pulchellum*, a new moss record for Asia from Singapore. *Folia Malaysiana* 3(3 & 4): 166–170.
- Tan BC & Jia Y (1999) A preliminary revision of Chinese Sematophyllaceae. *Journal of the Hattori Botanical Laboratory* 86: 1–70.

- Tan BC, Koponen T & Norris DH (2007) Bryophyte flora of the Huon Peninsula, Papua New Guinea. LXX. Sematophyllaceae (Musci) 1. *Acanthorrhynchium*, *Acroporium*, *Clastobryophilum*, *Pseudopiloecium*, *Radulina*, and *Trichosteleum*. *Annales Botanici Fennici* 44: 35–78.
- Tsubota H, Nakao N, Arikawa T, Yamaguchi T, Higuchi M, Deguchi H & Seki T (1999) A preliminary phylogeny of Hypnales (Musci) as inferred from chloroplast *rbcL* sequence data. *Bryological Research* 7: 233–248.
- Tsubota H, Nakao N, Yamaguchi T, Seki T & Deguchi H (2000) Preliminary phylogenetic relationships of the genus *Brotherella* and its allied genera (Hypnales, Musci) based on chloroplast *rbcL* sequence data. *Journal of the Hattori Botanical Laboratory* 88: 79–99.
- Tsubota H, Akiyama H, Yamaguchi T & Deguchi H (2001a) Molecular phylogeny of the Sematophyllaceae (Hypnales, Musci) based on chloroplast *rbcL* sequences. *Journal of the Hattori Botanical Laboratory* 90: 221–240.
- Tsubota H, Akiyama H, Yamaguchi T & Deguchi H (2001b) Molecular phylogeny of the genus *Trismegistia* and related genera (Sematophyllaceae, Musci) based on chloroplast *rbcL* sequences. *Hikobia* 13: 529–549.
- Tsubota H, Arikawa T, Akiyama H, Luna ED, Gonzalez D, Higuchi M & Deguchi H (2002) Molecular phylogeny of Hypnobryalean mosses as inferred from a large-scale dataset of chloroplast *rbcL*, with special reference to the Hypnaceae and possibly related families. *Hikobia* 13: 645–665.

Manuscript received 08 June 2012, accepted 12 April 2013

Linis, Virgilio C and Tan, Benito Ching. 2013. "Eight additional new records of Philippine mosses." *Telopea: Journal of plant systematics* 15, 45–49.

<https://doi.org/10.7751/telopea2013006>.

View This Item Online: <https://www.biodiversitylibrary.org/item/282073>

DOI: <https://doi.org/10.7751/telopea2013006>

Permalink: <https://www.biodiversitylibrary.org/partpdf/304706>

Holding Institution

The Royal Botanic Gardens and Domain Trust, New South Wales, Australia

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: The Royal Botanic Gardens and Domain Trust, New South Wales, Australia

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.