

**Case 2827**

***Gebia major capensis* Krauss, 1843 (currently *Upogebia capensis*; Crustacea, Decapoda): proposed replacement of neotype, so conserving the usage of *capensis* and also that of *G. africana* Ortmann, 1894 (currently *Upogebia africana*)**

N. Ngoc-Ho

*Laboratoire de Zoologie (Arthropodes), Muséum National d'Histoire Naturelle, 61 rue de Buffon, 75231 Paris, France*

Gary C.B. Poore

*Department of Crustacea, Museum of Victoria, Swanston Street, Melbourne, Victoria 3000, Australia*

**Abstract.** The purpose of this application is to conserve the accustomed usage of the specific names of two South African species of prawns: *Upogebia capensis* (Krauss, 1843) and *U. africana* (Ortmann, 1894). The latter species is commonly known as the mud-prawn or mud-shrimp. It is proposed to designate a replacement neotype for *capensis* from material of the species as presently understood; the previously designated neotype is a specimen of *africana*.

---

1. Three species of *Gebia* Leach, 1815 (p. 342; family UPOGEBIIDAE) were described from South Africa. *Gebia major* var. *capensis* Krauss, 1843 (p. 54) was originally described as a variety of *Gebia major* de Haan, [1841] (pl. 35, fig. 7; text (p. 165) published in [1849]; see Sherborn & Jentink (1895, p. 150) and Holthuis (1953, p. 37) for the dates of publication). The type material from Table Bay is now lost. The original description was short and by modern standards very incomplete and cannot be definitely reconciled with any single species known today. *G. subspinosus* Stimpson, 1860 (p. 22) was described from Simon's Bay; the fate of its type material is unknown. *G. africana* Ortmann, 1894 (p. 22, pl. 2, fig. 4) was described from Port Elizabeth. The holotype of this species is in the Zoological Museum, Strasbourg; it is a male without its abdomen (cephalothorax length 19.5 mm). Although in rather poor condition, it still shows the main characteristics of the species.

2. Since 1910 all three species have been referred to the genus *Upogebia* Leach, [1814] (pp. 386, 400; see Rathbun, 1897, p. 154, footnote for the date of publication). Until 1947 there was confusion between the three taxa and usually only one nominal species, *U. capensis*, was recognised (see, for example, Stebbing, 1900, p. 45; Stebbing, 1910; Balss, 1916, p. 34; Lenz & Strunck, 1914, p. 291; de Man, 1927, pp. 32–34; de Man, 1928, pp. 37, 41, 51). Barnard (1947, pp. 380, 381; 1950, pp. 514–520, fig. 96) revised the South African species of *Upogebia* and concluded that two species were involved: *U. capensis* (Krauss), characterised by a subdistal spine on the upper border of the merus of pereopod 1 and coxal spines on pereopods 1–3, and *U. africana* (Ortmann),



characterised by the absence of these spines. Stimpson's nominal species *Gebia subspinosa* was considered to be a synonym of *U. capensis* as the presence of coxal spines was mentioned in its original description.

3. Barnard's taxonomic arrangement has been generally adopted and at least 15 papers have been published since 1950 using his nomenclature. Besides agreeing on Barnard's morphological definition of the species, several authors have agreed on their ecological and geographical separation which is consistent with their type localities. There are no river outlets in Table Bay, and Krauss's material of *U. capensis* was therefore almost certainly from a marine rather than an estuarine habitat; the species is currently regarded as mainly marine to 80 metres depth, from southwestern and southern Africa between Lüderitz and Mossel Bay (Hill, 1981; Branch & Branch, 1981; Kensley, 1981). *U. africana* is estuarine to 18 metres depth mostly in eastern South Africa between Olifants River and Natal (Siegfried, 1962; Hill, 1977; Branch & Branch, 1981; Kensley, 1981; Hanekom, 1982; Martin & Baird, 1987; Hanekom & Erasmus, 1988; Zoutendyk & Bickerton, 1988). A further six references demonstrate this usage (Schaefer, 1970; Hill & Allanson, 1971; Ngoc-Ho, 1979, 1991; Emmerson, 1983; Atkinson & Taylor, 1988) and this is the usage in general marine biology texts in South Africa.

4. Sakai (1982, p. 44, fig. 9c, pls. A6, D5–6) selected a neotype for *Upogebia capensis* (Krauss, 1843) from material collected from Knysna, eastern South Africa by Hartmann in 1967, and originally identified as *africana* Ortmann, 1894 (see Hartmann-Schröder & Hartmann, 1974, p. 49). The specimen is a male, 55 mm in total length, housed in the Zoologisches Museum, Hamburg (catalogue no. ZMH 30877, selected from material originally registered as ZMH 29852). Sakai considered that *U. africana* was a junior synonym of *U. capensis*, and his selection of a neotype for *capensis* from material commonly assigned to *U. africana* in effect sank the latter name. The name *U. subspinosa* was revived by Sakai for what has been commonly called *U. capensis*, and the latter name was applied to *U. africana* (as defined by the holotype and as generally understood).

5. Sakai's (1982) selection of a neotype for *Upogebia capensis* has been ignored, probably not deliberately, by 10 authors in seven ecological papers since 1982 (see para. 3 above) and followed by only one (Holthuis, 1991, p. 233), who noted the unfortunate consequence that the name *capensis* has been transferred from one species to the other. The neotype upsets the nomenclature generally adopted since Barnard's (1947), and (1950) papers, and it came from material collected at the Knysna estuary (G. Hartmann, personal communication) which is not only far from Table Bay but is ecologically different and outside the geographical range of *Upogebia capensis* as generally understood (see Kensley, 1981, p. 31). In order to preserve the current usage of *Upogebia capensis* we propose that Sakai's (1982) neotype should be set aside and a replacement selected from material corresponding to the *U. capensis* of authors, collected in a marine environment, within the accepted geographical range and as close as practicable to the original type locality. The proposed replacement neotype, specimen no. 14895 in the South African Museum, Cape Town, was determined by K.H. Barnard as an ovigerous female with carapace length 22 mm and total length 65 mm; it is from Saldanha Bay, South Africa.

6. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary powers to set aside the neotype designation of Sakai (1982) for *Gebia major capensis* Krauss, 1843 and to designate in its place specimen no.



14895 in the South African Museum, for which the data are given in para. 5 above;

- (2) to place the following names on the Official List of Specific Names in Zoology:
  - (a) *capensis* Krauss, 1843, as published in the trinomen *Gebia major* var. *capensis* and as defined by the neotype designated in (1) above;
  - (b) *africana* Ortmann, 1894, as published in the binomen *Gebia africana*.

### Acknowledgements

We thank Prof L.B. Holthuis for comments on this proposal, and Dr G. Hartmann (Zoologische Museum, Hamburg) and Ms M. van der Merve (South African Museum) for the loan of material.

### References

- Atkinson, R.J.A. & Taylor, A.C. 1988. Physiological ecology of burrowing decapods. *Symposia of the Zoological Society of London*, **59**: 201–226.
- Balss, H. 1916. Crustacea II: Decapoda Macrura und Anomura (ausser Fam. Paguridae). *Beiträge zur Kenntniss der Meeresfauna Westafrikas*, **2**(1): 11–46.
- Barnard, K.H. 1947. Descriptions of new species of South African decapod Crustacea, with notes on synonymy and new records. *Annals and Magazine of Natural History*, (11)**13**(102): 361–392.
- Barnard, K.H. 1950. Descriptive catalogue of South African decapod Crustacea (crabs and shrimps). *Annals of the South African Museum*, **38**: 1–837.
- Branch, G. & Branch, M. 1981. *The living shore of South Africa*. 272 pp., 60 pls., 388 figs. Struik, Cape Town.
- Emmerson, W.D. 1983. Tidal exchange of two decapod larvae *Palaemon pacificus* (Caridea) and *Upogebia africana* (Thalassinidea) between the Swartkops River estuary and adjacent coastal waters. *South African Journal of Zoology*, **18**(4): 326–330.
- Haan, W. de. [1841], [1849]. Crustacea. In Siebold, P.F. de, *Fauna Japonica, sive descriptio animalium, quae in itinere per Japoniam... annis 1823–1830 collegit*. Pls. 33–37, 39–42, 47 [1841]; pp. 165–243 [1849]. Lugduni-Batavorum.
- Hanekom, N.M. 1982. A study of two thalassinid prawns in the non-*Spartina* regions of the Swartkops estuary. *Occasional Bulletin of the Zoological Society of Southern Africa*, **2**: 99–100.
- Hanekom, N.M. & Erasmus, T. 1988. Variations in size compositions of populations of *Upogebia africana* (Ortmann) (Decapoda, Crustacea) within the Swartkops Estuary and possible influencing factors. *South African Journal of Zoology*, **23**(4): 259–265.
- Hartmann-Schröder, G. & Hartmann, G. 1974. Zur Kenntnis des Eulitorals der afrikanischen Westküste zwischen Angola und Kap der Guten Hoffnung und der afrikanischen Ostküste von Südafrika und Moçambique unter besonderer Berücksichtigung der Polychaeten und Ostracoden. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, **69** (Ergänzungsband): 1–514.
- Hill, B.J. 1977. The effect of heated effluent on egg production in the estuarine prawn *Upogebia africana* (Ortmann). *Journal of Experimental Marine Biology and Ecology*, **29**(3): 291–302.
- Hill, B.J. 1981. Respiratory adaptations of three species of *Upogebia* (Thalassinidea, Crustacea) with special reference to low tide periods. *Biological Bulletin, Marine Biological Laboratory Woods Hole*, **160**(2): 272–279.
- Hill, B.J. & Allanson, B.R. 1971. Temperature tolerance of the estuarine prawn *Upogebia africana*. *Marine Biology*, **11**(4): 337–343.
- Holthuis, L.B. 1953. On the dates of publication of W. de Haan's volume on the Crustacea of P.F. von Siebold's 'Fauna Japonica'. *Journal of the Society for the Bibliography of Natural History*, **3**(1): 36–47.



- Holthuis, L.B.** 1991. FAO species catalogue. Vol. 13. Marine lobsters of the world. An annotated and illustrated catalogue of species of interest to fisheries known to date. *FAO Fisheries Synopsis*, **125**(13): 1–292.
- Kensley, B.** 1981. On the zoogeography of southern African decapod Crustacea, with a distributional checklist of the species. *Smithsonian Contributions to Zoology*, **338**: 1–64.
- Krauss, F.** 1843. *Die Südafrikanischen Crustaceen. Eine Zusammenstellung aller bekannten Malacostraca...* 68 pp., 4 pls. Schweizerbart, Stuttgart.
- Leach, W.E.** [1813]–[1814]. Crustaceology. In: Brewster, D. (Ed.), *The Edinburgh Encyclopedia*, vol. 7. Part 1, pp. 383–384 [1813]; part 2, pp. 385–437 [1814].
- Leach, W.E.** 1815. A tabular view of the external characters of four classes of animals, which Linné arranged under Insecta; with the distribution of genera composing three of these classes into orders, & c, and descriptions of several new genera and species. *Transactions of the Linnean Society of London*, **11**(2): 306–400.
- Lenz, H. & Strunk, K.** 1914. Die Dekapoden der Deutschen Südpolar-Expedition 1901–1903. 1. Brachyuren und Macruren mit Ausschluss der Sergestiden. *Deutsche Südpolar-Expedition, Zoologie*, **15**(7,3): 257–345.
- Man, J.G. de.** 1927. A contribution to the knowledge of twenty-one species of *Upogebia* Leach. *Capita Zoologica*, **2**(5): 1–58.
- Man, J.G. de.** 1928. The Decapoda of the Siboga Expedition. Part 7. The Thalassinidae and Callianassidae collected by the Siboga-Expedition with some remarks on the Laomediidae. *Siboga-Expedition*, **39a**(6): 1–187.
- Martin, A.P. & Baird, D.** 1987. Seasonal abundance and distribution of birds on the Swartkops estuary, Port Elisabeth. *Ostrich*, **58**(3): 122–134.
- Ngoc-Ho, N.** 1979. A taxonomic study of six species of *Upogebia* Leach (Crustacea, Decapoda, Thalassinidea) in the collections of the British Museum (Natural History), London. *Bulletin of the British Museum (Natural History)*, Zoology, **35**(2): 127–200.
- Ngoc-Ho, N.** 1991. Sur quelques Callianassidae et Upogebiidae de la Nouvelle Calédonie (Crustacea, Thalassinidea). Pp. 281–311 in Richer de Forges, B. (Ed.), *Le benthos des fonds meubles de Nouvelle-Calédonie*, vol. 1. ORSTOM Editions, Paris.
- Ortmann, A.** 1894. Crustaceen. Zoologische Forschungsreisen in Australien und dem Malayischen Archipel. *Denkschriften der Medicinisch-Naturwissenschaftlichen Gesellschaft zu Jena*, **8**: 3–80.
- Rathbun, M.J.** 1897. Revision of nomenclature of the Brachyura. *Proceedings of the Biological Society of Washington*, **11**: 153–167.
- Sakai, K.** 1982. Revision of Upogebiidae (Decapoda, Thalassinidea) in the Indo-West Pacific region. *Researches on Crustacea. The Carcinological Society of Japan*, Special Number, **1**: 1–106.
- Schaefer, N.** 1970. The functional morphology of the foregut of three species of decapod crustacea: *Cyclograpsus punctatus* (Milne-Edwards), *Diogenes brevisrostris* Stimpson, and *Upogebia africana* (Ortmann). *Zoologica Africana*, **5**(2): 309–326.
- Sherborn, C.D. & Jentink, F.A.** 1895. On the dates of the parts of Siebold's 'Fauna Japonica' and Giebel's 'Allgemeine Zoologie' (first edition). *Proceedings of the Zoological Society of London*, **1895**: 149–150.
- Siegfried, W.R.** 1962. A preliminary report on the biology of the mud-prawn *Upogebia africana* (Ortmann). *Investigational Report. Department of Nature Conservation*, **1**: 24.
- Stebbing, T.R.R.** 1900. South African Crustacea. *Marine Investigations in South Africa*, **1**: 14–66.
- Stebbing, T.R.R.** 1910. General catalogue of South African Crustacea. *Annals of the South African Museum*, **6**: 281–593.
- Stimpson, W.** 1860. Prodromus descriptionis animalium evertibratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit. Pars VIII: Crustacea Macrura. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **12**: 22–47.
- Zoutendyk, P. & Bickerton, I.** 1988. Burrow identification of some estuarine organisms. *South African Journal of Zoology*, **23**(3): 235–238.



Ngoc-Ho, Nguyen and Poore, Gary C. B. 1992. "Gebia major capensis Krauss, 1843 (currently Upogebia capensis; Crustacea, Decapoda): proposed replacement of neotype, so conserving the usage of capensis and also that of G. africana Ortmann, 1894 (currently Upogebia africana)." *The Bulletin of zoological nomenclature* 49, 187–190.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/44490>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/30725>

#### **Holding Institution**

Natural History Museum Library, London

#### **Sponsored by**

Natural History Museum Library, London

#### **Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: International Commission on Zoological Nomenclature

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.