PROCEEDINGS OF THE UNITED STATES NATIONAL MUSEUM

SMITHSONIAN INSTITUTION
U. S. NATIONAL MUSEUM

Vol. 101

Washington: 1951

No. 3272

THE OCEANIC CRABS OF THE GENERA *PLANES* AND *PACHYGRAPSUS*

By FENNER A. CHACE, Jr.

ON September 17, 1492, at latitude approximately 28° N. and longitude 37° W., Columbus and his crew, during their first voyage to the New World, "saw much more weed appearing, like herbs from rivers, in which they found a live crab, which the Admiral kept. He says that these crabs are certain signs of land . . ." (Markham, 1893, p. 25). This is possibly the first recorded reference to oceanic crabs. Whether it refers to *Planes* or to the larger swimming crab, *Portunus* (*Portunus*) *sayi* (Gibbes), which is seldom found this far to the east, may be open to question, but the smaller and commoner *Planes* is frequently called Columbus's crab after this item in the discoverer's diary.

Although these crabs must have been a source of wonder to mariners on the high seas in the past as they are today, the first adequate description of them did not appear until more than two centuries after Columbus's voyage when Sloane (1725, p. 270, pl. 245, fig. 1) recorded specimens from seaweed north of Jamaica. A short time later Linnaeus (1747, p. 137, pl. 1, figs. 1, *a-b*) described a similar form, which he had received from a Göteborg druggist and which was reputed to have come from Canton. This specimen, which Linnaeus named *Cancer cantonensis*, may be the first record of the Pacific *Planes cyaneus*. The Atlantic species, found "in Palgi *Fuco natante*," was finally described under the name *Cancer minutus* by Linnaeus in the tenth edition of "Systema Naturae" (1758, p. 625). Bowdich (1825, p. 15, pl. 12, figs. 2, *a-b*) briefly described and figured a crab found on a floating log northeast of Madeira as *Planes clypeatus*.

There is little doubt that Bowdich's species is conspecific with that described earlier by Linnaeus, and the combination *Planes minutus* is now generally accepted for the Atlantic species, despite repeated attempts to substitute *Nautilograpsus* H. Milne-Edwards (1837, p. 90) for *Planes*.

As specimens of *Planes* trickled into collections from all parts of the world in the early part of the nineteenth century, there was a natural tendency to propose new names for varieties from widely separated localities. A dozen different specific names were applied to these crabs between 1775 and 1858. Thereafter the acquisition of larger series of specimens served to illustrate the variability of these forms. All the post-Linnaean names were subsequently synonymized with *P. minutus*, and no additional ones were proposed except *Planes marinus* Rathbun (1914, p. 120). The validity of even that species became more and more questionable as time passed without additional specimens being found.

It was the rediscovery of that form that led to the present study. In December 1947 four crabs were found among marine organisms attached to a derelict Japanese mine that drifted ashore at Lincoln Beach, Oreg. These proved to be the species described by Dr. Rathbun and not reliably reported since her record of the type series taken in the Pacific Ocean west of Baja California. Search of the uncataloged collections in the U. S. National Museum revealed two more specimens of this species, these from the Hawaiian Islands. A subsequent query addressed to Dr. C. H. Edmondson at the Bernice P. Bishop Museum in Honolulu led to the discovery of several lots in that institution. A comparison of this form with specimens of *Planes minutus* indicates that the two are generically as well as specifically distinct and that there are at least two species of *Planes* s. s., one found in the Atlantic and one in the Pacific.

No attempt is made here to delimit these oceanic forms completely either taxonomically or geographically. Additional material from critical areas and "hosts" must be studied before this can be done. It has been my aim to bring together as many of the scattered references to these species as possible in the hope that workers elsewhere may be encouraged to re-examine material at their disposal or to collect specimens from other areas with a view to verifying or modifying the conclusions outlined here. A species believed to be identical and common in all the warmer seas of the world soon loses its interest; collectors fail to take the trouble to preserve specimens when they are found, and museum workers neglect to examine critically those that are received. If this trend can be reversed as far as these crabs are concerned, it is not unlikely that interesting findings related to speciation and zoogeography may be forthcoming.

Practically none of the references to *Planes* in the literature are determinable to species except on a geographic basis. The synonymies listed below have been compiled more or less arbitrarily, all the records from the Atlantic and Indian Oceans being credited to *P. minutus* and those from the Pacific to *P. cyaneus*. It is very possible that this disposition, particularly as it concerns the Indian Ocean forms, may have to be altered when additional data are obtained.

This opportunity is taken to thank those persons whose gracious assistance has made this study possible: Dr. Louis W. Hutchins, of the Woods Hole Oceanographic Institution, for furnishing the specimens of *Pachygrapsus marinus* which prompted the survey; John C. Armstrong, of the American Museum of Natural History, Dr. Elisabeth Deichmann, of the Museum of Comparative Zoology, and Dr. C. H. Edmondson, of the Bernice P. Bishop Museum, for providing loans of critical specimens; Mr. and Mrs. Boonyong Nikrodhananda, of Bangkok, Siam, for making a cursory survey of drifting objects that might harbor crabs, during an extended transpacific voyage; and finally my colleagues in the U. S. National Museum for their assistance and encouragement during the preparation of this paper.

MORPHOLOGICAL CHARACTERS

The species of *Planes* are extremely variable. This fact has led to the general abandonment of all previous attempts to recognize more than one species in the genus. When Stimpson (1860, p. 231) made the statement that "we are unable to distinguish the specimens [from Baja California] from those collected from Gulfweed in the North Atlantic," the conspecific status of the various forms of *Planes* seemed to be finally established, and few subsequent attempts to delimit other than the genotype were attempted; even the validity of the very distinct form described as *Planes marinus* by Dr. Rathbun has been questioned.

The present study indicates, however, that at least some of the variable characters involved show a discontinuity between the Atlantic and Pacific forms of the genus (table 1). Probably the most useful of these characters is based on the relative lengths of the walking legs. In *P. cyaneus* from the Pacific the legs are noticeably shorter than in *P. minutus* from the Atlantic and Indian Oceans. This difference in relative leg length is not so apparent in figure 1 as it would be if specimens of *P. minutus* and *P. cyaneus* of similar carapace lengths had been available for figuring. The discrepancy is more obvious in figure 2, *d*, *e*, where legs from specimens of similar size are compared. It was found that the simplest way to demonstrate this difference is to compare the combined lengths of the three distal segments of the second walking leg with the carapace length (figs. 6, 7). The slight overlap in this character between the Atlantic and Pacific forms may

be a real one, or it may be a result of contraction or expansion of the leg in preservative or of the possibility that regenerating and not fully developed legs of *P. minutus* have been included among the material measured.

TABLE 1.—*Synopsis of characters*

Characters	<i>Planes minutus</i>	<i>Planes cyaneus</i>	<i>Pachygrapsus marinus</i>
CARAPACE:			
Length.....	3.7 to 19 mm.....	5 to 25 mm.....	5 to 19 mm.
Proportions (see fig. 4)...	About as wide as long, length to width as 1:0.91 to 1.12.	Usually slightly longer than wide, length to width as 1:0.86 to 1.06.	Distinctly wider than long, length to width as 1:1.07 to 1.16.
Shape (see fig. 3).....	Subquadrate in young (3.7-6 mm.); trapezoidal in medium sized (6-12 mm.); laterally convex in old specimens (12-19 mm.).	Laterally convex at all ages.	Subquadrate at all ages.
Surface of branchial regions.	Faintly striate laterally..	Faintly striate laterally....	Distinctly striate laterally.
MALE ABDOMEN:			
Shape (see fig. 2, <i>g-i</i>)..	Rather broadly triangular, length of four distal segments about 1.24 times basal width of fourth segment.	Narrowed beyond third segment, length of four distal segments about 1.26 times basal width of fourth segment.	Broadly triangular, length of 4 distal segments about 1.08 times basal width of fourth segment.
Terminal segment.....	Rather narrowly triangular, about as long as basal width.	Laterally convex, about 0.9 as long as basal width.	Broadly triangular, about 0.9 as long as basal width.
CHELAE:			
Fixed finger (see fig. 2, <i>a-c</i>).	Bent sharply downward, especially in males.	Bent sharply downward, especially in males.	Not noticeably bent downward.
Surface of hand.....	With prominent sharp granules near lower margin.	With prominent sharp granules near lower margin.	Few inconspicuous granules near lower margin.
WALKING LEGS:			
Form (see fig. 2, <i>d-f</i>)..	Long, slender, and flattened.	Short, slender, and flattened.	Rather short, stout, and not noticeably flattened.
Hairs on upper anterior margin of propodus of three anterior pairs.	Forming a natatory fringe.	Forming a natatory fringe.	Stout and forming a dense growth without a natatory fringe.
Proportion of length of three distal segments of second pair to carapace length (see fig. 6).	0.83 to 1.07.....	0.68 to 0.89.....	0.77 to 0.99.

Planes cyaneus appears to be a somewhat larger form than *P. minutus*. The smallest specimens of the latter examined measure 3.7 mm. in carapace length, and the largest 19 mm. Comparable measurements for *P. cyaneus* are 5 and 25 mm. The mean carapace width is usually about the same as the mean length in *P. minutus*, whereas it is somewhat less in *P. cyaneus* (figs. 4, 5); this character is demonstrable only in long series of specimens, however, and is of far less importance than the length of the walking legs.

FIGURE 1.—*a*, *Planes minutus*, male (U. S. N. M. No. 4558), carapace 13.4 mm. long
b, *Planes cyaneus*, male (U.S.N.M. No. 20695), carapace 16.6 mm. long; *c*, *Pachygrapsus marinus*, male paratype (U.S.N.M. No. 22833), carapace 13.8 mm. long.

The shape of the carapace is much more variable in *P. minutus* than in *P. cyaneus*, but much of the variation in the former species is correlated with size (fig. 3). In the smallest specimens of *P. minutus* examined (carapace length 3.8 to 4.5 mm.) the carapace is nearly square. At a carapace length from about 4.5 to 5.5 mm., the carapace becomes more or less convex laterally, as in *P. cyaneus*; this stage is smaller than any specimens of the Pacific species examined, however, and so there is little difficulty in separating the young stages of the two species. Following this stage the carapace becomes narrowed posteriorly and assumes a more or less trapezoidal shape; this shape persists in most specimens up to those with carapace lengths of 11 or 12 mm. and is the form commonly found living on *Sargassum*. As the crab becomes larger than this, the carapace becomes more convex laterally and less narrowed posteriorly and thus closely resembles *P. cyaneus* once more. No such polymorphism has been noted in the latter species; in that form the carapace is laterally convex at all sizes. The front is often more deeply excavate in *P. cyaneus* than in *P. minutus*, but this character is extremely variable.

The shape of the male abdomen, one of the characters upon which Dana based *P. cyaneus*, is usually distinctive in the two species (fig. 2, *g, h*). The lateral margins are less sharply convergent from the third to the sixth segment in *P. cyaneus* than in *P. minutus*, thereby giving the abdomen of the former a narrower appearance. The terminal segment is in the form of a nearly equilateral triangle in *P. minutus*, whereas its lateral margins are noticeably convex in *P. cyaneus*.

No consistent differences in the form of the male abdominal appendages could be found. The apparent differences indicated in figure 2, *j-r*, are probably no greater than may be found by examining first pleopods from several individuals of the same species. Even the form of that appendage in *Pachygrapsus marinus* is of questionable value as a taxonomic character.

From the material available the two forms of *Planes* appear to be sufficiently well marked to be recognized as distinct species. Additional material, particularly from the South Atlantic, Indian, and western Pacific regions should be examined, however; if intermediate forms are found in those areas it may be advisable to reduce *P. cyaneus* to subspecific status. As mentioned in the remarks on relative growth (p. 73), there is some indication that two forms may be represented in the Atlantic: a small trapezoidal one living on floating weed and a larger one, approaching *P. cyaneus* in form, found on turtles, floating logs, and other flotsam. Additional specimens of various sizes from the latter habitat should be examined to determine whether this form is distinct or not.

FIGURE 2.—a, Right chela of *Planes minutus*, male (U.S.N.M. No. 4558), carapace 13.4 mm. long; b, right chela of *Planes cyaneus*, male (U.S.N.M. No. 17449), carapace 15.9 mm. long; c, right chela of *Pachygrapsus marinus*, male paratype (U.S.N.M. No. 22833), carapace 13.8 mm. long; d, right second walking leg of *Planes minutus*, male (U.S.N.M. No. 17712), carapace 15.5 mm. long; e, right second walking leg of *Planes cyaneus*, male (U.S.N.M. No. 20700), carapace 15.3 mm. long; f, right second walking leg of *Pachygrapsus marinus*, male (U.S.N.M. No. 88069), carapace 15.1 mm. long; g, Abdomen of *Planes minutus*, male (U.S.N.M. No. 4558), carapace 13.4 mm. long; h, abdomen of *Planes cyaneus*, male (U.S.N.M. No. 20695), carapace 16.6 mm. long; i, abdomen of *Pachygrapsus marinus*, male paratype (U.S.N.M. No. 22833), carapace 13.8 mm. long; j, first right pleopod, mesiocaudal view, of *Planes minutus*, male (U.S.N.M. No. 4558), carapace 13.4 mm. long; k, denuded tip of first right pleopod, mesiocaudal view, of *Planes minutus*, male (U.S.N.M. No. 17712), carapace 15.5 mm. long; l, same, laterocephalic view; m, first right pleopod, mesiocaudal view of *Planes cyaneus*, male (U.S.N.M. No. 20695), carapace 16.6 mm. long; n, denuded tip of same, mesiocaudal view; o, same, laterocephalic view; p, first right pleopod, mesiocaudal view, of *Pachygrapsus marinus*, male paratype (U.S.N.M. No. 22833), carapace 13.8 mm. long; q, denuded tip of same, mesiocaudal view; r, same, laterocephalic view.

The species that Dr. Rathbun (1914, p. 120, pl. 3) described as *Planes marinus* differs so markedly from the other species of *Planes* in its wider, subquadrate, and distinctly striate carapace, differently formed chelae, and the absence of a natatory fringe of hairs on the propodi of the first three pairs of walking legs (table 1) that it is here transferred to the genus *Pachygrapsus*. Except for its much smaller size, this species is very similar to *Pachygrapsus crassipes* Randall from the Pacific coast of North America and the shores of the Japanese islands and Korea. The only obvious character, apart from size, in which *P. crassipes* and *P. marinus* differ is the absence of lateral lobules on the front in the latter species. Unless this species is removed from *Planes* on the basis of the characters mentioned above, the distinction between *Planes* and *Pachygrapsus* would cease to exist and *Pachygrapsus* would have to be synonymized with *Planes*.

Examination of one of the two specimens recorded as *Planes marinus* by Ward (1939, p. 14) indicates that these specimens, like the others mentioned in that paper, are *P. cyaneus*.

RELATIVE GROWTH

The examination and measurement of the large number of specimens of *Planes minutus* used in this study suggest a problem involving the relative growth of that species which might prove to be of considerable interest if it could be followed up with the analysis of more material from drift logs, turtles, and so forth.

The carapace length-width relationship remains fairly constant, with a slight tendency toward narrowing, from the smallest immature specimens examined to a carapace length of about 11 mm. At this stage the carapace seems to become somewhat narrower rather abruptly and continues to become narrower at a slightly more rapid rate than during the younger stages (figs. 4, 5). A similar, but even more striking, trend is noticed in the relative shortening of the walking legs (figs. 6, 7). In other words, there is a tendency for the larger specimens of *P. minutus* to approach the laterally convex, narrow form (fig. 3) and short walking legs of *P. cyaneus*. These similarities are so striking that I was at first led toward the belief that the larger Atlantic specimens are conspecific with those from the Pacific. It will be seen from figures 4 through 7, however, that the mean width of the carapace in the Atlantic form never becomes quite as narrow relative to the length as in the Pacific form, and the legs are always distinctly longer in the Atlantic species (tables 2, 3).

Although the material at hand has not been sufficiently well documented to verify the hypothesis, it is not improbable that most of the specimens of *P. minutus* having a carapace length of more than about 11 mm. are not found on *Sargassum*, but on flotsam and turtles. The

TABLE 2.—Mean values of carapace proportions

<i>Planes minutus</i>			<i>Planes cyaneus</i>			<i>Pachygrapsus marinus</i>		
Number of specimens	Mean carapace length	Mean carapace width	Number of specimens	Mean carapace length	Mean carapace width	Number of specimens	Mean carapace length	Mean carapace width
	<i>Milli-meters</i>	<i>Milli-meters</i>		<i>Milli-meters</i>	<i>Milli-meters</i>		<i>Milli-meters</i>	<i>Milli-meters</i>
57	4.2	4.2	-----	-----	-----	-----	-----	-----
89	5.0	5.1	-----	-----	-----	-----	-----	-----
78	6.0	6.2	15	6.2	6.0	-----	-----	-----
76	7.0	7.3	-----	-----	-----	6	6.9	7.7
76	8.0	8.3	8	8.3	8.3	-----	-----	-----
53	9.0	9.4	-----	-----	-----	8	8.7	9.8
46	10.0	10.4	-----	-----	-----	-----	-----	-----
22	10.9	11.2	-----	-----	-----	-----	-----	-----
9	12.4	12.7	11	11.8	11.6	6	12.3	13.9
10	14.3	14.7	12	14.1	14.2	16	14.0	15.8
-----	-----	-----	-----	-----	-----	7	15.5	17.4
9	17.0	17.3	10	17.3	17.2	5	17.6	19.9
-----	-----	-----	2	22.6	22.5	-----	-----	-----

TABLE 3.—Mean values of total length of three distal segments of second walking leg

<i>Planes minutus</i>			<i>Planes cyaneus</i>			<i>Pachygrapsus marinus</i>		
Number of specimens	Mean carapace length	Mean length of leg segments	Number of specimens	Mean carapace length	Mean length of leg segments	Number of specimens	Mean carapace length	Mean length of leg segments
	<i>Milli-meters</i>	<i>Milli-meters</i>		<i>Milli-meters</i>	<i>Milli-meters</i>		<i>Milli-meters</i>	<i>Milli-meters</i>
16	4.2	4.1	-----	-----	-----	-----	-----	-----
46	5.0	5.0	-----	-----	-----	-----	-----	-----
46	6.0	6.0	15	6.3	5.1	-----	-----	-----
52	7.0	7.0	-----	-----	-----	4	7.3	6.8
54	7.9	7.8	8	8.2	6.6	-----	-----	-----
45	8.9	8.7	-----	-----	-----	7	8.7	7.8
41	9.9	9.6	-----	-----	-----	-----	-----	-----
21	10.9	10.3	12	11.6	9.0	-----	-----	-----
15	12.6	11.6	-----	-----	-----	6	12.3	10.4
12	14.5	13.0	12	14.1	10.5	16	14.0	11.8
12	16.3	14.4	-----	-----	-----	7	15.5	12.8
4	18.3	16.0	10	17.0	13.1	5	17.6	14.2
-----	-----	-----	5	22.5	16.9	-----	-----	-----

shorter legs may be better adapted to this more sessile existence or, conversely, the longer legs of the smaller individuals may be advantageous in scurrying and swimming about among the floating weed. Whether there is a gradual reduction in the relative lengths of the legs with the transfer of the animal to a different means of transport or whether there are two forms of *Planes* in the North Atlantic—a small, trapezoidal form that spends its entire life on floating weed and a larger, laterally convex, and short-legged form that lives only on flotsam and turtles—can be determined only when more material,

particularly of immature specimens, from flotsam and turtles is available for study.

Similar trends are not apparent in the specimens of *Planes cyaneus* or *Pachygrapsus marinus*, but the relative growth of these two species cannot be reliably demonstrated from the material studied.

FIGURE 3.—Carapace growth in males of *Planes minutus* (a-h), *Planes cyaneus* (i-n), and *Pachygrapsus marinus* (o-t). The figures denote the U.S.N.M. catalog numbers and, in parentheses, the carapace lengths in mm.

FIGURE 4.—Scatter diagram showing the variation in proportions of the carapace in *Planes minutus*, *Planes cyaneus*, and *Pachygrapsus marinus*.

COLOR

The variable and protective coloration of *Planes minutus* has been a cause of speculation for more than a century. Some of the early workers believed that those specimens which were greenish or brownish yellow mottled with darker shades of the same color sought out *Sargassum* clumps to live on in order to be less conspicuous to their enemies. These *Sargassum* inhabitants also often have a large white or yellowish patch on the dorsal surface, which seems to duplicate the patches of bryozoan colonies with which the plant fronds are almost invariably encrusted. Those specimens found on turtles are usually yellowish, those on logs and drifting trees often dark reddish brown, and those on the pelagic gastropod *Janthina* lilac blue. Light reddish shades have also been recorded on several occasions. Examples of some of these colors are shown in Murray and Hjort (1912, pl. 6).

Müller (1881, p. 473) was apparently the first to suggest that these crabs change their color according to the object to which they cling.

Not until a few years ago, however, was the nature of this change demonstrated. Dr. Hitchcock (1941, pp. 26–30) found that the chromatophores of specimens of *P. minutus* he investigated at Bermuda were responsive to white, black, red, blue, yellow, and green backgrounds, but that there was no immediate change in the general color of the crab because of the extracellular pigment in the hypodermis and exoskeleton. These findings seem to indicate that the apparent color of any particular specimen can be changed only at the time of molting, and any individual that moves to a differently colored background subsequent to hardening of the shell would cease to be protectively colored. Interesting results might be obtained by confining molting specimens to backgrounds of various colors and patterns.

The color of *P. cyaneus* has been recorded as cerulean blue, bluish gray mottled with brown, yellowish clouded with brown, and reddish brown. It is probable that a type of color change comparable to that in *P. minutus* occurs in this species as well, although no similar experiments on it have been attempted.

FIGURE 5.—Relative growth of the carapace in *Planes minutus*, *Planes cyaneus*, and *Pachygrapsus marinus*, based on mean values (table 2) plotted logarithmically.

The color in life of *Pachygrapsus marinus* has not been published. The specimens from a Japanese mine which prompted this survey were rather dark reddish brown a short time after preservation in alcohol.

FIGURE 6.—Scatter diagram showing the variation in the combined length of the three terminal segments of the second walking leg in *Planes minutus*, *Planes cyaneus*, and *Pachygrapsus marinus*.

FOOD

The only published record of the stomach contents of any of these species is that of Miss Crane (1937, p. 78) based on a large ovigerous female of *Planes cyaneus* taken from the tail of a green turtle. The stomach of this specimen contained finely digested animal matter. Miss Crane suggests the possibility that this food material might represent the excrement of the turtle. Some support is given to this theory by the fact that larger specimens of *Planes* are commonly found around the tail and hind legs of turtles. Specimens living on other objects, however, must be able to assimilate less digested food.

DISTRIBUTION

The relative abundance of these oceanic crabs is more or less dependent upon the prevalence of flotsam or of floating or swimming organisms to which the crabs may cling. They are therefore found far more commonly on *Sargassum* in the Sargasso Sea area of the North Atlantic than in any other part of the world. Records of their occur-

rence in other areas are still too scattered and frequently too unreliable to justify the usual distribution note that *Planes* is "found in all tropical and temperate seas."

FIGURE 7.—Relative growth of the three terminal segments of the second walking leg in *Planes minutus*, *Planes cyaneus*, and *Pachygrapsus marinus*, based on mean values (table 3) plotted logarithmically.

The more reliable locality records for the species of *Planes* are indicated in figure 8. Not included on this chart are several references in the literature to Australian specimens without specific locality data. Even some of the localities indicated, particularly those from older records, require verification. It is obvious that in some cases specimens picked up at sea have been referred to the last port of call of the vessel involved. The large lot of specimens recorded by Dr. Rathbun (1918, p. 257) from the Galápagos Islands (U.S.N.M. No. 5046) prove to belong to the Atlantic *Planes minutus*. This determination is substantiated by the fact that the shrimp from the same collection belong to the *Sargassum* fauna of the North Atlantic. Similar circumstances may explain the record of the single specimen of *P. cyaneus* from Mauritius (U.S.N.M. No. 17667) if the presence of *P. minutus* alone in the Indian Ocean is borne out by other material. Even one of the lots of *P. minutus* from that region, that from Kerguelen Island (U.S.N.M. No. 15054), is questionable; in the collections of

the U. S. National Museum is a specimen of *Portunus* (*Portunus*) *sayi* (Gibbes) that bears the same locality data, and this crab is otherwise reliably reported only from the Atlantic. Examination of additional material from the Indian Ocean will be necessary before it can be definitely ascertained whether one or both species occur there. The fact that Alcock (1900) recorded no specimens of the genus indicates that it is not a common component of the Indian fauna.

One of the most interesting aspects of the distribution of *Planes* is its apparent absence from the Gulf of Mexico west of the Dry Tortugas despite its relative abundance from the Straits of Florida eastward and its occurrence in the Caribbean area. Joel W. Hedgpeth, of the Institute of Marine Science at Port Aransas, Tex., and Dr. Allan F. Archer, of the Alabama Museum of Natural History, inform me that they have never encountered the crab although they have searched through great masses of *Sargassum* from the Gulf. It would be interesting to know what factor or factors preclude the dispersion of *Planes* into an area such as this where the presence of floating weed and particularly of sea turtles would seem to afford ideal sources of attachment for the crabs.

There are likewise very few records of the presence of *Planes* in the central South Atlantic, possibly because of the comparatively small amount of exploratory work done in that extensive region. If the occurrence of *P. minutus* in the Indian Ocean is finally confirmed, it would be of considerable interest to know more of the distribution of the genus in the South Atlantic, particularly whether the Pacific *P. cyaneus* extends into the western part of the South Atlantic and, if it does, whether the ranges of the two species overlap there. Inasmuch as *Planes* is not infrequently found among the under-water growth on ships' hulls and as the crabs are said to be common in the Straits of Magellan (Jacquinot and Lucas, 1853, p. 78), through which much of the steamship traffic passed in the early part of this century, it would be remarkable if intermingling of the two forms had not occurred commonly in the past.

The known distribution of *Pachygrapsus marinus* is limited to the records from the coast of Oregon and the Hawaiian Islands, in addition to the type locality west of Baja California; further collecting will probably show its range to be far more extensive in the eastern North Pacific and possibly in the western portion as well. Dr. C. H. Edmondson informs me that, of 18 lots and 150 or more specimens identified as *Planes minutus* in the Bernice P. Bishop Museum, 3 lots containing a total of 22 specimens are apparently *Pachygrapsus marinus* and 3 of the remaining 15 lots contain both *Planes* and *Pachygrapsus*, including 30 or more specimens of the latter. All the lots containing *Pachygrapsus marinus* were taken off Oahu.

FIGURE 8.—Known distribution of the species of *Planes* based only on locality records that are apparently valid and reasonably exact.

Genus PLANES Bowdich

PLANES MINUTUS (Linnaeus)

- Cancellus marinus minimus quadratus* SLOANE, 1725, p. 270, fig. 1 ("on the Sargasso and other Submarine Sea-Plants, on the Northside of Jamaica").
- Cancer minutus* LINNAEUS, 1758, p. 625 ("Habitat in Palgi *Fuco natante*, supra aquam saepius cursitans"); 1767, p. 1040 [2040] (west Europe and "Pelagi *Fuco natante*").—OSBECK, 1771, p. 116 (Sargasso Sea).—FABRICIUS, 1775, p. 402.—MÜLLER, O. F., 1776, p. 195.—PENNANT, 1777, p. 3, pl. 1, fig. 2 (British shores, among algae).—FABRICIUS, 1781, p. 497.—HERBST, 1783, p. 110, pl. 2, fig. 32.—FABRICIUS, 1787, p. 316.—OLIVIER, 1791, pp. 143, 154.—FABRICIUS, 1793, p. 443 ("In Pelago frequens").
- Cancer pusillus* FABRICIUS, 1775, p. 402 ("in Oceano boreali").—HERBST, 1783, p. 112.—FABRICIUS, 1787, p. 316.—OLIVIER, 1791, pp. 143, 155.—FABRICIUS, 1793, p. 443.
- Turtle-Crab BROWNE, 1789, p. 421, pl. 42, fig. 1 ("on the back of a turtle, near the western islands").
- Pinnotheres glaberimus* Bosc, 1801-02, p. 244 ("dans la haute mer sur les fucus").
Not *Cancer glaberrimus* Herbst, 1790.
- Pinnotheres minutus* Bosc, 1801-02, p. 244.
- Pinnotheres pusillus* Bosc, 1801-02, p. 244.
- Grapsus minutus* LATREILLE, 1802-03, p. 68.—LEACH, 1814, p. 430.
- Grapsus cinereus* SAY, 1817, p. 99 (Gulf Stream, common). Not *G. cinereus* Bosc, 1802.
- Grapsus pelagicus* SAY, 1818, p. 442.—ROUX, 1828, p. 55, pl. 6, figs. 7-9 (off Sardinia, on turtles).—HITCHCOCK, E., 1835, p. 29 (Gulf Stream, on seaweed).—THOMPSON, J. V., 1836, p. 374 (off America, on gulfweed, abundant).—MILNE-EDWARDS, H., 1852, p. 174.
- Planes clypeatus* BOWDICH, 1825, p. 15, pl. 12, figs. 2a-b (northeast of Madeira, on water-logged logs of American pine, in great numbers among *Lepas anatifera*).
- Grapsus testudinum* ROUX, 1828, p. 52, pl. 6, figs. 1-6 (under folds of skin of legs and anus of turtles).
- Nautilograpsus minutus* MILNE-EDWARDS, H., 1837, p. 90.—MACLEAY, 1838, p. 66 (Atlantic, on gulfweed).—KRAUSS, 1843, p. 44 (Gulf Stream).—DE KAY, 1844, p. 15 (off New York harbor, on seaweed).—GOODSIR, 1845, p. 73, pl. 7, fig. 1 (lat. 25° to 36° N., long. 40° W.; lat. 40° N.).—GIBBES, 1850, p. 182 [18].—MILNE-EDWARDS, H., 1852, p. 174.—STIMPSON, 1858, p. 103 [49] (North Atlantic, common in *Sargassum*).—MILNE-EDWARDS, A., 1862, p. F-6 (Réunion Island, Indian Ocean).—HELLER, 1863, p. 114 (African coast of Mediterranean; Cyprus; Genoa; Lesina); 1865, p. 50 (Sargasso Sea).—SMITH and HARGER, 1874, p. 26 (off Georges Bank; lat. 41°25' N., long. 65°05'-30' W.).—BROCCHI, 1875, p. 80, pl. 17, fig. 133.—MAN, 1879, p. 69 (part: Atlantic Ocean).—MOSELEY, 1879, p. 568 (Sargasso Sea, on *Sargassum*).—SMITH, 1879, p. 120 (Bermuda; lat. 41°20'-30' N., long. 65° to 65°30' W.); 1880, p. 263 (Woods Hole, Mass.).—STOSSICH, 1880, p. 192 (Lesina).—WAGNER, 1880, p. 96.—KINGSLEY, 1881, p. 202 (part: Gulf Stream; West Indies; Surinam; Falkland Islands [?]; Natal; Rio Gambia; France).—CZERNIAVSKY, 1884, p. 135 (Odessa).—MILNE-EDWARDS, A., 1884, p. 25 (Sargasso Sea, on *Sargassum*).—CARUS, 1885, p. 524.—VERRILL, 1885, p. 591 (Gulf Stream, common, usually clinging to clusters of floating seaweeds).—MIERS, 1886, p. 254 (part: Gomera, Canary Islands; between Bermuda and Azores; off Sombrero Island, West Indies; northwest Atlantic,

- on gulfweed; North Atlantic, on *Fucus*).—POUCHET and GUERNE, 1886, p. 878 (Azores, from stomach of turtle, *Thalassochelys caretta*).—SMITH, 1886, p. 618.—BARROIS, 1888, p. 15 (Ponta-Delgada, Sao Miguel, Azores, on timber).—GOURRET, 1888, p. 10 (Marseilles, on ships from the Atlantic and the Indies).—HEILPRIN, 1888, p. 320 (off Shelly Bay, Bermudas).—CHEVREUX and GUERNE, 1893, pp. 443-444 (lat. 37°55' N., long. 0°40' E., clinging to tail of turtle, *Thalassochelys caretta*; lat. 37°26' N., long. 0°50' E., clinging to tail of *Thalassochelys*).—ORTMANN, 1893, p. 58 (Florida Stream, on *Sargassum*; Sargasso Sea, on *Sargassum*; South Equatorial Current).—MILNE-EDWARDS, A., and BOUVIER, 1894, p. 49 (lat. 39°08' N., long. 28°34' W., on floating barrel; lat. 42°21'29'' N., long. 18°33'45'' W.; lat. 42°31'21'' N., long. 19°38'08'' W.; lat. 43°29'30'' N., long. 21°33'40'' W., on flotsam; Fayal, Horta, Azores, on rocks at low tide; lat. 42°02'26'' N., long. 44°05'30'' W., on *Sargassum*; lat. 39°20'30'' N., long. 33°29'40'' W., on flotsam).—ORTMANN, 1894, p. 710 (Oran; Africa; Atlantic Ocean; "Sargasso").—NUTTING, 1895, p. 29 (Gulf Stream, in *Sargassum*).—HODGSON, 1896, p. 178 (Plymouth Sound, England, on bottom of ship from Iquique, Peru; English Channel, 2-3 miles off Plymouth, on fragment of cork structure).—MILNE-EDWARDS, A., and BOUVIER, 1899, p. 37 (lat. 37°16'-39' N., long. 27°05'-58' W., on turtles, *Thalassochelys caretta*; lat. 39°12' N., long. 33°09' W., on flotsam; lat. 38°08' N., long. 25°55' W., on flotsam; lat. 38°26' N., long. 28°51' W.); 1900, p. 108 (off Ponte-Delgado, on *Sargassum*).—RANKIN, 1900, p. 526 (Bermuda, in tide-pools with *Pachygrapsus* among *Sargassum*).—YOUNG, 1900, p. 286 (part: not all distribution records).—HOWE, 1900, p. 240 (approx. lat. 40° N., long. 70° W.).—NORMAN and SCOTT, 1906, p. 5.—BOUVIER, 1907, p. 37 (variable coloration).—COULON, 1907, p. 123 [17] (Sargasso Sea).—NORMAN, 1907, p. 357 (Channel Islands).—STIMPSON, 1907, p. 121 (North Atlantic, between lat. 20-36° N., common on floating logs, seaweed (*Sargassum*), etc.).—NORMAN and BRADY, 1909, p. 9.—PESTA, 1918, p. 446, fig. 147 (Lesina).—BORASCHI, 1921, p. 10, pl. 1, fig. 9 (larvae: Quarto dei Mille; Palermo).—MIRANDA Y RIVERA, 1921, p. 189 (Playa de la Malagueta, Spain).—MONOD, 1932, p. 219 (Fedhala, Morocco, common on marine plants and other objects).
- Nautilograpsus major* MACLEAY, 1838, p. 66 (Cape of Good Hope).—KRAUSS, 1843, p. 44.—MILNE-EDWARDS, H., 1852, p. 175.
- Nautilograpsus smithii* MACLEAY, 1838, p. 67 (Cape of Good Hope).—KRAUSS, 1843, p. 44.—MILNE-EDWARDS, H., 1852, p. 175.
- Grapsus diris* COSTA, 1838-53, [p. 2, pl. 4, fig. 1 (color)] (Naples, "among rocks above water").—MILNE-EDWARDS, H., 1852, p. 175.
- Planes linnaeana* LEACH, 1815, pl. 27, figs. 1-3, and accompanying text (coasts of Devon and Cornwall).—BELL, 1844-53, p. 135, figs. (Devonshire; Cornwall).—WHITE, 1847, p. 41 (Devonshire; south Atlantic).—COCKS, 1849, p. 80 (Falmouth, England, from stomach of fish).—WHITE, 1850, p. 19 (Devonshire).—GOSSE, 1855, p. 159, fig. 302.—MCANDREW, 1860, p. 21.—LUKIS and LE LIEVRE, 1862, p. 232 (Guernsey, Channel Islands).—MARCUSEN, 1867, p. 359 (Black Sea).—CARRINGTON and LOVETT, 1882, p. 181 (Looe, Cornwall, among stalked barnacles on timber; Penzance, from soil and seaweed on derelict cask of paraffin).—CORNISH, 1882, p. 118 (off Penzance, England, from soil and seaweed growing on derelict cask of paraffin); 1884, p. 116 (near Penzance, England, from derelict water-logged barrel of paraffin drifted ashore).
- Planes minutus* WHITE, 1847, p. 42 ("Gulf of Florida;" Atlantic Ocean, on gulfweed).—DANA, 1852b, p. 346; 1854, p. 27.—CUNNINGHAM, 1870, p. 493

(south of Azores, on *Sargassum*); 1871, p. 501.—RATHBUN, 1897b, p. 29 (Kingston Harbor, Jamaica); 1898a, p. 288 (Dry Tortugas, in gulfweed; lat. 31°30' N., long. 75°11' W.); 1898b, p. 604, (part: lat. 31°16' N., long. 71°50' W.).—ORTMANN, 1899, p. 1188 (part).—RATHBUN, 1900a, p. 279; 1900b, p. 587 (part).—THOMPSON, D'A. W., 1901, p. 6 (Atlantic).—BORRADAILE, 1903, p. 432 (Male, Maldives Islands, on floating cuttlebone).—DOFLEIN, 1904, p. 130 (lat. 30°6.7' S., long. 87°50.4' E.).—MAYER, 1905, p. 102.—STEBBING, 1905, p. 43 (False Bay, South Africa).—RATHBUN, 1905, p. 4 (Woods Hole and Wareham, Mass.; Sakonnet Point, R. I., under rocks).—NOBILI, 1906, p. 321 (Djibouti).—VERRILL, 1908, p. 325, fig. 7, pl. 13, figs. a-j', pl. 27, fig. 6 (off Bermuda).—LENZ, 1910, pp. 560, 574 (Ste. Marie, Madagascar).—STEBBING, 1910, p. 320 (South Africa: False Bay; 38 miles northwest of Rable Bay).—DOFLEIN and BALSS, 1912, p. 39 (part?: lat. 44°32' S., long. 61°25' W.).—FOWLER, 1912, p. 443, pls. 140-141 (Cape May, N. J.).—MURRAY and HJORT, 1912, pp. 103, 633, 671, pl. 6 (color) (approx. lat. 36°05' N., long. 43°58' W.).—PESTA, 1912, p. 122 (Lesina).—SUMNER, OSBURN, and COLE, 1913, p. 675 (western part of Vineyard Sound, Mass., commonly on gulfweed).—LENZ and STRUNCK, 1914, p. 284 (off Madeira, on floating weed; off New Amsterdam Island, on *Macrocystis*; southeast of Madagascar; south of Azores, on *Sargassum*).—STEBBING, 1914a, p. 345 (Falkland Islands); 1914b, p. 266, pl. 24 (Saldanha Bay, South Africa; lat. 29°54' N., long. 34°10' W. to lat. 33°53' N., 32°27' W.).—BOUVIER, 1915, p. 127 (Port Louis, Mauritius).—HAY and SHORE, 1918, p. 448, pl. 36, fig. 6 (Beaufort, N. C.).—RATHBUN, 1918, p. 253 (part: off Woods Hole, Mass., to Florida and Bahamas; Azores; Indian Ocean; Kerguelen Island).—TESCH, 1918, p. 83 [footnote] (part: not all distribution records).—CROZIER, 1918, pp. 262-263 (Bermuda, on floating tree; coloration).—LUEDERWALDT, 1919, p. 435 (Iguape, São Paulo, Brazil).—BALSS, 1922a, p. 83 (Monrovia, West Africa).—BOUVIER, 1922, p. 74, pl. 2, fig. 2 (color) (east-central North Atlantic between lat. 28° and 39° N., and long. 23° and 43° W., on turtles, *Sargassum*, floating wood, etc.).—HYMAN, 1924, p. 5, pl. 1, figs. 1-10 (Beaufort Harbor, N. C.).—RATHBUN, 1924, p. 157 (off eastern Cuba, on *Sargassum*).—MONOD, 1927, p. 621 (Souelaba, Douala Bay, Cameroon, on floating tree trunk).—PERKINS, 1928, p. 53.—RATHBUN, 1929, p. 35, fig. 48 (off Nova Scotia).—NOBRE, 1931, p. 107, figs. 60-63 (Douro River, Portugal, on root of tree washed in by storm).—MONOD, 1933, p. 80.—RATHBUN, 1933, p. 89, fig. 83 (Puerto Rico; St. Thomas).—EKMAN, 1935, pp. 9, 425 (part, not fig.).—NOBRE, 1936, p. 63, pl. 24, fig. 45, pl. 46, fig. 45a.—MONOD, 1939, p. 567 (lat. 30°10' N., long. 47° W., on *Sargassum*).—BOUVIER, 1940, p. 291, fig. 182.—HITCHCOCK, H. B., 1941, pp. 26-30 (Bermuda; color change).—LEBOUR, 1944, p. 114, fig. 4 (Bermuda, in *Sargassum*; larvae).—ZARIQUIEY ALVAREZ, 1946, p. 163, pl. 18, fig. b.

Nautilograpsus diris HELLER, 1863, p. 115.

Nautilograpsus pelagicus HELLER, 1863, p. 115.

Nautilograpsus cyaneus TARGIONI TOZZETTI, 1872, p. 8 (Sargasso Sea); 1877, p. 125, pl. 8, figs. 5a-f (Atlantic Ocean). Not *Planes cyaneus* Dana, 1852.

Planes linneana COUCH and BATE, 1878, p. 469 (English Channel off French coast, under tail of hawksbill turtle; Devon or Cornwall).

Nautilus grapsus minutus MORISON, 1942, p. 269.

Planes sp. WHITE, 1847, p. 42 (St. Kitts Island; Falkland Islands; Brazil; Swan River).—BEEBE, 1932, pp. 188-196 (Bermuda).

Nautilograpsus sp. MÜLLER, F., 1881, p. 473 (color change).

TABLE 4.—Material examined of *Planes minutus*¹

Locality	Latitude N.	Longitude W.	Sur- face temp.	Date	Sta- tion	Collector	Specimens	Catalog No.²	Remarks
Off eastern North America south of Newfoundland.	44 29 --	52 53 --	° F.	Sept. 21, 1935		A. H. Leim	1♂	72195	
Maine: New Harbor, Muscongus Bay.	-- --	-- --		Aug. 15, 1944		L. W. Scattergood	1♂	88063	From loggerhead turtle.
Massachusetts: Georges Bank	41 52 --	66 24 --		Sept. 16, 1927		W. C. Schroeder	1 ovig. ♀	8720 (MCZ)	
Vicinity of Woods Hole	-- --	-- --		1899		B. A. Bean	1♂	44672	
Wareham	-- --	-- --		Aug. 12, 1898		Outram Bangs	1♂	5976 (MCZ)	Found alive on beach.
Rhode Island: Sakonnet Point	-- --	-- --		Sept. 13, 1904		W. Nye, Jr.	1 ovig. ♀	31476	Under rocks.
5-6 miles off Block Island	-- --	-- --		July 30, 1928		A. K. Fisher	1 ♂ 1 ovig. ♀	62510	From hawksbill turtle.
New Jersey: East of	40 50 --	60 53 --		July 2, 1915		C. W. Furlong, <i>Kitty A.</i>	7♂ 2♀	53351	From <i>Sargassum</i> .
Do	40 34 30	66 48 00	67	July 14, 1885	2332	<i>Albatross</i>	1♀	11038	
Do	40 10 15	70 26 00	64	Sept. 26, 1884	2243	<i>Albatross</i>	3♂ 2♀ (1 ovig.)	7285	
Do	39 59 40	70 41 10	68	Sept. 20, 1883	2090	<i>Albatross</i>	1 ovig. ♀	5422	
Do	{39 45 00	69 44 45	70	Aug. 4, 1881	935	<i>Fish Hawk</i>	2♂ 1♀	4558	1 male figured.
Do	{39 49 25	69 49 00	72	do	937				
Do	39 40 --	71 30 --	68	Sept. 8, 1881	994	<i>Fish Hawk</i>	1♀	40519	
Do	39 34 15	71 41 15	74	Aug. 19, 1884	2203	<i>Albatross</i>	4♂ 2 ovig. ♀	8220	
Do	39 29 00	70 58 40	69.5	Sept. 30, 1883	2095	<i>Albatross</i>	1♂	15052	
Do	39 22 50	68 25 00	72	July 30, 1883	2041	<i>Albatross</i>	4♂ 3 ovig. ♀	6515	
Cape May Point	-- --	-- --		Sept. 24, 1928		H. G. Richards	1♀	74887	
Maryland: East of	38 19 26	68 20 20	81	July 28, 1883	2039	<i>Albatross</i>	2♂	40520	Carapace of smaller speci- men figured.
North Carolina: East of	36 20 24	74 46 30	69	June 4, 1885	2425	<i>Albatross</i>	1♂ 2♀	17895	
Edge of Gulf Stream off Cape Hatteras.	-- --	-- --		June 6, 1903		B. A. Bean	5♂ 6 ovig. ♀	31080	From <i>Sargassum</i> .
200 miles off Cape Hatteras	-- --	-- --		Oct. 1921		<i>Carnegie</i>	16♂ 15♀ (2 ovig.)	60268	Carapaces of three males figured.

Beaufort	July 16, 1912	W. P. Hay, Fish Hawk	1♂ 1♀	51019	From log with <i>Lepas</i> on shore.
Do		Kinglsey (?)	2♂ 1♀	56798	
South Carolina:					
East of	Jan. 5, 1885	Albatross	7♂ 7♀ (1 ovig.) 1 megalops.	15053	
Do	Jan. 30, 1914	Bache	1♀	49957	
Do	Feb. 1, 1914	Bache	5♂ 4♀	49942	
Do	Feb. 2, 1914	Bache	9♂ 6♀	49946	
Georgia:					
East of	May 9, 1893	University of Iowa Bahama Expedition.	1♂ 1♀	74517	From <i>Sargassum</i> .
Do	Nov. 23, 1887	Albatross	19♂ 19♀	22098	
Florida:					
St. Augustine		E. Samuels	1♂ 1 ovig. ♀	71132	Do.
Florida Keys			2♂ 6♀ (4 ovig.)	5977 (MCZ)	
Key West	Apr. 15-27, 1884	Albatross	1♀	18448	
Dry Tortugas		University of Iowa	1♂	68857	
Bahamas and West Indies					
Bahamas:					
Northeast of Abaco Island	Feb. 28, 1914	Bache	1♀	49940	
Do	June 15, 1903	Bean and Riley	3♂ 2 ovig. ♀	31061	
About 120 miles north by east of Abaco Island.	do	do	6♂ 18♀ (10 ovig.)	31081	Do.
60 miles north of Abaco Id.	June 16, 1903	B. A. Bean	5♀ (4 ovig.)	31079	Do.
North of Abaco Island	May 2, 1886	Albatross	5♂ 4♀ (2 ovig.)	11399	
Green Turtle Cay		E. A. Andrews	1♂ 2♀	20714	
Between Nassau and Elbow Cay.	July 21, 1903	B. A. Bean	2♂ 1 ovig. ♀	31062	
Eleuthera Island: 1 mile northeast of Governor's Harbor.	May 14, 1936	W. J. Clench	2♂ 2 ovig. ♀	9486 (MCZ)	Along open beach.
Between Nassau and Cat Island.	Mar. 18, 1937	W. L. Schmitt	2♂	75155	

See footnotes at end of table.

TABLE 4.—Material examined of *Planes minutus* 1—Continued

Locality	Latitude N.	Longitude W.	Surface temp.	Date	Station	Collector	Specimens	Catalog No.	Remarks
Bahamas—Continued									
Exuma Sound	24 30 43	76 23 45	73	Mar. 13, 1886	2632	Albatross	1♂	11409	
Rum Cay				Feb. 13, 1934		T. Barbour and J. Greenway.	1♂ 3♀ (2 ovig.)	8974 (MCZ)	
Off eastern Cuba				1923	12h	Williams Galapagos Expedition.	2♀ (1 ovig.)	57739	From <i>Sargassum</i> .
Do				1923	12j	do	2♂	57738	
Haiti						Dr. Weinland	1♂	5982 (MCZ)	
North of St. Thomas	20 --	63 --				Dr. Kershner	7♂ 5♀ (2 ovig.)	6714	
St. Thomas				Jan. 17-24, 1884		Albatross	1♀	18563	
Do						Hassler	23♂ 22♀ 169y	5981 (MCZ)	
Jamaica: Kingston Harbor				May-July 1896		F. S. Conant	1♂ 1 ovig. ♀	19600	
Swan Island, Caribbean Sea				1916		George Nelson	1 ovig. ♀	9077 (MCZ)	
Bermudas				1876-77		F. V. Hamlin	6♂ 8♀ (1 ovig.)	4027	
Do						G. Brown Goode	12♂ 14♀ (4 ovig.)	43047	
Do						Walter Faxon	1♂ 1♀	5980 (MCZ)	
Do						Bermuda Biological Station.	9♂ 5♀	7692 (MCZ)	
Do				Jan. 8, 1903		do	6♂ 13♀ (7 ovig.)	8420 (MCZ)	Do.
Do				July 2, 1903			12♂ 9♀ (6 ovig.)	9089 (MCZ)	Do.
Do				Mar. 10, 1916			1y.	9109 (MCZ)	In floating seaweed.
North shore				Jan. 12, 1882		G. Howes	2♂ 8♀	5170	From <i>Sargassum</i> thrown up on shore. Carapaces of 2 males figured.
Agers Island				Apr. 1928		E. Deichmann	30♂ 15♀ (6 ovig.)	65645	
Hamilton Island				Mar. 3-7, 1910		T. Barbour	17♂ 12♀ (7 ovig.)	9114 (MCZ)	On beach.
South shore of Long Bird Island.				June 13-18, 1936		F. A. Chace, Jr.	1 ovig. ♀	9229 (MCZ)	
Northwest of	32 30 --	65 48 --	66.6	Feb. 5-6, 1914	10176	Bache	7♂ 3♀	49944	
Southwest of	30 27 --	66 05 --	68.2	Feb. 19, 1914	10182	Bache	3♂ 4♀ (3 ovig.)	49945	
Do	30 27 --	66 05 --	68.2	do	10182	Bache	1♀	49948	
Do	29 17 --	67 07 --	68.1	Feb. 21, 1914	10184	Bache	14♂ 8♀ (3 ovig.)	49943	

Do	28	51	--	70	08	--	67	Feb. 24, 1914	10188	Bache	2♂ 2♀	49941
Do	28	51	--	70	08	--	67	do	10188	Bache	1♂ 1♀	49947
Do	31	15	42	67	39	10	--	Jan. 13, 1884	Hyd. 37	Albatross	44♂ 50♀ (4 ovig.)	7744
400 miles southeast of										Crosby	2 ovig. ♀	74916
Mid-Atlantic	38	50	--	42	40	--	--	Apr. 27, 1859	--	O. Dabney	8♂ 13♀	1593 (MCZ)
Do	38	50	--	42	40	--	--	--	--	do	1 ovig. ♀	8790 (MCZ)
Do	32	--	--	36	--	--	--	July 5, 1860	--	D. D. Raulet	13♂ 16♀ (10 ovig.) 4y.	8787 (MCZ)
Do	29	23	--	37	08	--	--	Feb. 22, 1861	--	do	20♂ 21♀ (3 ovig.) 47y.	5990 (MCZ)
Do	21	--	--	51	--	--	--	--	--	C. C. Craft	1♂ 2♀	44671
Do	11	05	--	50	01	--	--	--	--	C. W. Furlong, <i>Kitty A.</i>	7♂ 13♀ (3 ovig.)	53350
Azores:										O. Dabney	1♂ 1 ovig. ♀	5983 (MCZ)
Fayal								Jan. 1893	--	Lewis Dexter	3♂ 3♀	17712
3 miles off Fayal Harbor												From log. Parts of largest and smallest males figured.
Madeira Islands:												From floating timber.
Bay of Funchal								Aug. 1921	--	Adolfo Cesar de Noronha.	1♀	55365
Do								Sept. 1921	--	do	1♂	55366
Canary Islands: Tenerife								Feb. 1884	--	Capt. J. R. Lyon, <i>Cashmere</i> .	3♂ 4♀ (1 ovig.)	72521
Indian Ocean.	90° E. on								--		5♂ 9♀ (2 ovig.)	13914
Kerguelen Island (?)								Sept. 28, 1874	--	J. H. Kidder	1♀	15054

¹ Including only lots for which the locality data are apparently valid and reasonably exact.

² Unless otherwise noted, catalog numbers refer to specimens in the U. S. National Museum; (MCZ) stands for the Museum of Comparative Zoology, Cambridge, Mass.

PLANES CYANEUS Dana

- Cancer cantonensis* LINNAEUS, 1747, p. 137, pl. 1, figs. 1a-b (Canton [?]).
- Grapsus pusillus* HAAN, 1835, p. 59, pl. 16, fig. 2.—MILNE-EDWARDS, H., 1852, p. 175. Not *Cancer pusillus* Fabricius, 1775.
- Nautilograpsus minutus* MILNE-EDWARDS, H., and LUCAS, 1843, p. 28 (record from off Valparaiso credited to Eydoux).—NICOLET, 1849, p. 168 (Valparaiso Bay).—JACQUINOT and LUCAS, 1853, p. 78 (Port Famine, Straits of Magellan, very abundant).—STIMPSON, 1860, p. 231 [103] (Cape San Lucas, Baja California).—MAN, 1879, p. 69 (part: Japan; New Guinea).—KINGSLEY, 1881, p. 202 (part: Peru; west coast of Mexico; Alaska; China; New Zealand; Falkland Ids. [?]).—MIERS, 1886, p. 254 (part: South Pacific, near Kermadec Islands, among seaweed; North Pacific, off Volcano Island; off Japan).—LENZ, 1901, p. 472 (between Honolulu and San Francisco). Not *Cancer minutus* Linnaeus, 1758.
- Planes cyaneus* DANA, 1852a, p. 250 (lat. 28° N., long. 174° E.); 1852b, p. 347 (lat. 15°50' S., long. 105° W.); 1853, p. 1593; 1855, pl. 22, fig. 1.—WARD, 1939, p. 14 (lat. 29°20' N., long. 141°25' W.).
- Nautilograpsus angustatus* STIMPSON, 1858, p. 103 [49] (lat. 34° N., long. 155° E.); 1907, p. 121, pl. 16, fig. 1 (lat. 34° N., long. 151° W.).
- Planes minutus* MIERS, 1876, p. 39 (New Zealand).—HASWELL, 1882, p. 99.—FILHOL, 1885, p. 390.—FAXON, 1895, p. 30 (lat. 00°13'00'' S., long. 84°52'00'' W.; lat. 2°34'00'' N., long. 92°06'00'' W.; off Acapulco, Mexico; all on green turtles).—RATHBUN, 1898b, p. 604 (part: lat. 1°03' N., long. 80°15' W.; Gulf of California).—ORTMANN, 1899, p. 1188 (part).—RATHBUN, 1900b, p. 587 (part); 1902, p. 278 (about 200 miles north of Wenman Island, Galápagos, at base of tail of green turtle); 1904, p. 189 (lat. 41° N., long. 141° W.; Point Loma, Calif.; off San Diego, Calif.).—FULTON and GRANT, 1906, p. 19.—RATHBUN, 1906, p. 840 (between Erben Bank and Kaiwi Channel, on *Veleva*: south coast of Oahu Island; south coast of Molokai Island, on floating stick).—RATHBUN, 1907, p. 69 (south of Gulf of California; off Guatemala; off Peru); 1910, pp. 589, 610-614, 616.—WEYMOUTH, 1910, p. 63, pl. 14, fig. 44.—CHILTON, 1911, p. 561 (Kermadec Islands).—DOFLEIN and BALSS, 1912, p. 39 (part: Smith Channel, Straits of Magellan).—RATHBUN, 1918, p. 253, pl. 63 (part: Humboldt Bay, Calif., to Peru; Galápagos Islands; Hawaiian Islands; Marquesas Islands; Japan; Mauritius).—TESCH, 1918, p. 83 [footnote] (part).—SCHMITT, 1921, p. 272, pl. 46, figs. 1-2 (La Jolla, Calif.; off Point Loma, Calif.).—BALSS, 1922b, p. 149 (Ogasawara Island).—BALSS, 1924, p. 336 (Juan Fernández Island).—URITA, 1926, p. 23 (off Ohsima, Ohsumi, Japan).—HALE, 1927, p. 181, fig. 182.—CHILTON and BENNETT, 1929, p. 768 (Moko Hinau, New Zealand, washed ashore on piece of pumice).—SIVERTSEN, 1933, p. 21.—GLASSELL, 1934, p. 302 (Gulf of California).—SAKAI, 1934, p. 323 (Japan).—EKMAN, 1935, p. 9, fig. 3 (not p. 425).—SAKAI, 1936, p. 226, fig. 120 (Japan).—AIKAWA, 1937, p. 152 (larvae).—CRANE, 1937, p. 77 (Santa Inez Bay, Mexico, on tail of green turtle; food).—SAKAI, 1939, p. 664, pl. 108, fig. 2 (Misaki, from turtle; Odawara, Sagami Bay; Simoda, on *Sargassum*; Ise Bay; Seto; Tosa Bay; Nagasaki; northern Daitozima; Tansui, Formosa).—STEINBECK and RICKETTS, 1941, pp. 45, 474 (south of Point Abrojos, Baja Calif., in folds of skin beside tail of tortoise-shell turtle, *Eretmochelys imbricata* [on flipper, p. 474]).—EDMONDSON, 1946, p. 306, fig. 183f (Hanauma Bay, Oahu Island).—POPE, 1948, pp. 296-297, fig.—RICHARDSON, 1949, p. 34, pl. 2, fig. 16. Not *Cancer minutus* Linnaeus, 1758.
- Nautilograpsus pusillus* MAN, 1879, p. 69. Not *Cancer pusillus* Fabricius, 1775.
- Planes marinus* WARD, 1939, p. 14 (lat. 32° 00' N., long. 135°20' W.; lat. 29°20' N., long. 141°25' W.). Not *P. marinus* Rathbun, 1914.

TABLE 5.—*Material examined of Planes cyaneus*¹

Locality	Latitude (N.)	Longitude (W.)	Surface temp.	Date	Station	Collector	Specimens	Catalog No. ²	Remarks
Off southern California; San Pedro.	° ' "	° ' "	° F.			H. N. Lowe.	1 ♀	32975	
Santa Catalina Island: Avalon Bay.				Oct. 22, 1910.		P. S. Barnhart, Anton Dohrn.	1 ♀	50269	
WSW. of San Nicholas Island.	33 02 15	120 36 30	61-62	Mar. 28, 1904.	4391	Albatross.	1 ♀	44674	
Off San Diego.	32 47 30	118 10 00	59	Jan. 23, 1889.	2928	Albatross.	1 ♂	17450	
Point Loma.				Jan. 28, 1889.		Albatross.	1 ♂ 1 ♀	17449	
Off Pacific coast of Mexico: Baja California: Turtle Bay.				Aug. 1, 1896.		A. W. Anthony.	1 ♂ 1 ovig. ♀	19517	
San José del Cabo.						do.	1 ♀	20694	
Gulf of California.				Apr. 29, 1888.		Albatross.	2 ♂ 1 ♀	22101	
Escuinapa, Sinaloa.						Dirac. Forest. y de Casa y Pesa.	1 ovig. ♀	60238	
Off Cabo Corrientes, Jalisco.	20 00 --	106 12 --	81-83	Oct. 12, 1904.	4587	Albatross.	1 ♀	33214	
San Benedicto Island.						A. W. Anthony.	1 ♂ 1 ♀	20695	Male figured.
Clarion Island.						do.	2 ♂ 3 ♀ (1 ovig.)	20700	Walking leg figured.
Off Acapulco.				Feb. 10, 1891.		Albatross.	1 ♂	20624	From Green Turtle.
Salina Cruz, Oaxaca.				Dec. 25, 1898.		Chas. C. Deam.	1 ♂ 1 ♀	32262	On sand beach.
Off Guatemala.	12 20 --	92 13 --	85	Oct. 17, 1904.	4605	Albatross.	1 ♂	33215	Carapace figured.
Off Panama: Taboga Island.				May 12-15, 1911.		Meek and Hildebrand.	1 ♂ 1 ♀	44173	
Between Panama and Galápagos Islands.	4°-5°	83°-85°		Apr. 1, 1888.		Albatross.	1 ♂	22100	
Northwest of Galápagos Islands.	2 34 00	92 06 00	82	Apr. 5, 1891.	3413	Albatross.	2 ♂ 2 ovig. ♀	4493 (MCZ)	From Green Turtle.
Off Ecuador.	1 03 --	80 15 --				Albatross.	1 ♀	22097	
Off Peru: Paífa.				Aug. 12, 1884.		W. H. Jones.	1 ♂ 1 ♀	14063	Carapace of male figured.

See footnotes at end of table.

TABLE 5.—*Material examined of Planes cyaneus*¹—Continued

Locality	Latitude (S.)	Longitude (W.)	Surface temp.	Date	Station	Collector	Specimens	Catalog No. ²	Remarks
Off Peru—Continued									
West of Palta	5 17	85 20	° F. 70-71	Nov. 10, 1904	4649	<i>Albatross</i>	2 ♀	33213	
Off Juan Fernández Islands, Chile.				Dec. 18, 1926		W. L. Schmitt	1 ♀	70886	
Do.				Dec. 10, 1926		do	1 ♂	70992	
40 miles east of				Sept. 13, 1944	U-42	U. S. Fish. Miss. in Chile.	1 ♂	81058	From drifting life raft. Carapace figured.
					-264				
West of California	Latitude (N.)			Sept. 7, 1871		W. H. Dall	1 ♂	15051	Carapace figured.
Do.	41 --	141 --		Aug. 25, 1945		M. F. Groves	8 ♀ (5 ovig)	82012	From floating wooden box.
Between California and Hawaiian Islands.	32 00	135 20				<i>Zaca</i>	1 ♂	7485 (AMNH)	Identified as <i>Planes martinus</i> by M. Ward.
Do.	30 56	139 50		Apr. 26, 1884		Lt. G. M. Stoney	3 ♂ 1 ♀	13617	
Do.	28 23 00	141 41 05	66	Mar. 19, 1902	3800	<i>Albatross</i>	1 ♂	29347	
Do.	28 23 00	141 41 05	66	do	do	<i>Albatross</i>	1 ♂	29348	From <i>Velilla</i> .
Do.	27 55	144 15				<i>Zaca</i>	1 ♂	7484 (AMNH)	Identified as <i>P. cyaneus</i> by M. Ward.
Hawaiian Islands:									
Molokai Island: Off south coast.			76	Apr. 2, 1902	3833	<i>Albatross</i>	1 ♂	29345	From floating stick. Carapace figured.
Oahu.									
Off south coast.			74	Mar. 28, 1902	3813	W. J. Thompson	1 ♂	88066	
Malae Kahana.						<i>Albatross</i>	1 ♂	29346	Carapace figured.
Midway Island.				April 1941		C. M. Cooke, Jr.	1 ♀	88068	
Baker Island.						V. D. P. Spicer	1 ♀	88065	
						Capt. J. Hammond	1 ♂ 3 ♀	5992 (MCZ)	One female appears to be <i>P. minutus</i> .
Mauritius.				1891 (?)		H. A. Ward	1 ♀	17667	

¹ Including only lots for which the locality data are apparently valid and reasonably exact.² Unless otherwise noted, catalog numbers refer to specimens in the U. S. National Museum; (MCZ) stands for the Museum of Comparative Zoology, Cambridge, Mass., and AMNH for the American Museum of Natural History, New York City.

TABLE 6.—Material examined of *Pachygrapsus marinus*

Locality	Latitude (N.)	Longitude (W.)	Surface temper- ature	Date	Sta- tion	Collector	Specimens	Catalog No. ¹	Remarks
Oregon: Lincoln Beach	° ' "	° ' "	° F.	Dec. 29, 1947		Ordnance Officer 13th Naval District.	3♂	88069	From Japanese mine. Walking leg figured.
Do				do		do	1♂	(WHOI) 22833	From Japanese mine.
West of Baja California	23 49 --	127 50 --				D. D. Raulet	1♂ 1♀	87988	Paratypes. Male fig- ured.
Do	23 49 --	127 50 --				do	27♂ 14 ovig. ♀		Paratypes. Carapaces of several males figured.
Hawaiian Islands:									
Oahu						C. M. Cooke, Jr.	2♂	88070	
Do				July 8, 1916		C. M. Cooke, III	1♂ 1♀	88064	
Mokapu				July 1, 1932		C. H. Edmondson	1♂ 1♀	88067	Found beside drift log.

¹ Catalog numbers refer to specimens in the U. S. National Museum; (WHOI) stands for the Woods Hole Oceanographic Institution, Woods Hole, Mass.

Genus **PACHYGRAPSUS** Randall**PACHYGRAPSUS MARINUS** (Rathbun)

Planes marinus RATHBUN, 1914, p. 120, pl. 3 (lat. 23°49' N., long. 127°50' W.); 1918, p. 258, pl. 64.—TESCH, 1918, p. 83 [footnote].—GLASSELL, 1934, p. 302.

LITERATURE CITED

AIKAWA, HIROAKI.

1937. Further notes on brachyuran larvae. Rec. Oceanogr. Works in Japan, vol. 9, No. 1, pp. 87-162, 36 figs.

ALCOCK, ALFRED WILLIAM.

1900. Materials for a carcinological fauna of India, No. 6: The Brachyura Catometopa or Grapsoidea. Journ. Asiatic Soc. Bengal, vol. 69, pt. 2 No. 3, pp. 279-456.

BALSS, HEINRICH.

1922a. Crustacea VII: Decapoda Brachyura (Oxyrhyncha und Brachyrhyncha) und geographische Übersicht über Crustacea Decapoda. In Michaelsen's "Beiträge zur Kenntnis der Meeresfauna Westafrikas," vol. 3, Lief. 3, pp. 71-110, 1 fig.

1922b. Ostasiatische Decapoden, IV: Die Brachyrhynchen (Canceridea). Arch. Naturg., Jahrg. 88, Abt. A, Heft 11, pp. 94-166, 2 figs., 2 pls.

1924. Decapoden von Juan Fernandez. In Skottsberg's "The Natural History of Juan Fernandez and Easter Island," vol. 3, pp. 329-340, 3 figs.

BARROIS, THÉODORE.

1888. Catalogue des Crustacés marins recueillis aux Açores durant les mois d'Août et de Septembre 1887, 110 pp., 8 figs., 4 pls. Lille.

BEEBE, WILLIAM.

1932. Nonsuch: Land of water, xv+259 pp., 46 figs., frontis. New York.

BELL, THOMAS.

1844-53. A history of the British stalk-eyed Crustacea, lxx+386 pp. London.

BORASCHI, LILLIA.

1921. Osservazioni sulle larve dei crostacei decapodi brachiuri e anomuri. R. Comitato Talassografico Italiano, Mem. 87, 32 pp., 2 pls.

BORRADAILE, LANCELOT ALEXANDER.

1903. Marine crustaceans, pt. 5: The crabs of the catometope families. In Gardiner's "The Fauna and Geography of the Maldives and Laccadive Archipelagoes," pp. 429-433, 4 figs. Cambridge.

BOSC, LOUIS A. G.

1801-02. Histoire naturelle des crustacés, contenant leur description et leurs mœurs; avec figures dessinées d'après nature, vol. 1, 258 pp. Paris.

BOUVIER, EUGÈNE LOUIS.

1907. Quelques impressions d'un naturaliste au cours d'une campagne scientifique de S. A. S. le Prince de Monaco (1905). Bull. Inst. Océanogr. Monaco, No. 93, 103 pp., 69 figs.

1915. Décapodes marcheurs (Reptantia) et stomatopodes recueillis à l'île Maurice par M. Paul Carié. Bull. Sci. France et Belgique, ser. 7, vol. 48, fasc. 3, 141 pp., 42 figs., 4 pls.

1922. Observations complémentaires sur les crustacés décapodes (abstraction faite des carides) provenant des campagnes de S. A. S. le Prince de Monaco. Rés. Camp. Sci. Monaco, fasc. 62, 106 pp., 6 pls.

1940. Décapodes marcheurs. Faune de France, vol. 37, 404 pp., 222 figs., 14 pls. Paris.

BOWDICH, THOMAS EDWARD.

1825. Excursions in Madeira and Porto Santo during the autumn of 1823, while on his third voyage to Africa, xii+278 pp., 22 pls. London.

BROCCHI, PAUL.

1875. Recherches sur les organes génitaux males des crustacés décapodes. Ann. Sci. Nat., Zool., ser. 6, vol. 2, art. 2, 131 pp., 7 pls.

BROWNE, PATRICK.

1789. The civil and natural history of Jamaica, viii+503 pp., 49 pls., 1 map. London.

CANO, G.

1891. Sviluppo postembrionale dei dorippidei, leucosiadi, corystoidei e grapsidi. Mem. Soc. Ital. Sci., vol. 8, ser. 3, No. 4, 14 pp., 3 pls.

CARRINGTON, JOHN THOMAS, and LOVETT, EDWARD.

1882. Notes and observations on British stalk-eyed Crustacea. Zoologist, ser. 3, vol. 6, pp. 178-185.

CARUS, JULIUS VICTOR.

1885. Coelenterata, Echinodermata, Vermes, Arthropoda. Prodromus faunae Mediterraneae sive descriptio animalium Maris Mediterranei incolarum quam comparata silva rerum quatenus innotuit adiectis locis et nominibus vulgaribus eorumque auctoribus in commodum zoologorum, vol. 1, xi+524 pp. Stuttgart.

CHEVREUX, EDOUARD and GUERNE, JULES DE.

1893. Crustacés et cirrhipèdes commensaux des tortues marines de la Méditerranée. Comptes Rendus Acad. Sci. Paris, vol. 116, pp. 443-446.

CHILTON, CHARLES.

1911. The Crustacea of the Kermadec Islands. Trans. New Zealand Inst., vol. 43, art. 49, pp. 544-573, 4 figs.

CHILTON, CHARLES, and BENNETT, E. W.

1929. Contributions for a revision of the Crustacea Brachyura of New Zealand. Trans. New Zealand Inst., vol. 59, pp. 731-778.

COCKS, W. P.

1849. Contributions to the fauna of Falmouth. Ann. Rep. Royal Cornwall Polytechnic Soc., pp. 38-102.

CORNISH, THOMAS.

1882. Floating crab at Penzance. Zoologist, ser. 3, vol. 6, No. 63, p. 118.

1884. Floating crabs at Penzance. Zoologist, ser. 3, vol. 8, p. 116.

COSTA, ORONZIO GABRIELE.

- 1838-53? Crostacei ed aracnedi. Fauna del Regno di Napoli ossia enumerazione di tutti gli animali che abitano le diverse regioni di questo regno e le acque che le bagnano contenente la descrizione de nuovi o poco esattamente conosciuti, 290 pp., 29 pls.

COUCH, JONATHON, and BATE, C. SPENCE.

1878. Crustacea: A Cornish fauna. Journ. Roy. Inst. Cornwall, No. 19, pt. 2, pp. 451-522.

COULON, LOUIS.

1907. Les crustacés du Musée d'Histoire naturelle d'Elbeuf, pt. 2: Les malacostracés. I. Thoracostracés (podophthalmaires). Bull. Soc. Étud. Sci. Nat. Elbeuf, vol. 26, pp. 119-194, 1 pl.

CRANE, JOCELYN.

1937. The Templeton Crocker Expedition, III: Brachygnathous crabs from the Gulf of California and the west coast of Lower California. Zoologica, New York, vol. 22, pt. 1, No. 3, pp. 47-78, 8 pls.

CROZIER, WILLIAM JOHN.

1918. On the coloration of *Planes minutus*. Amer. Nat., vol. 52, No. 616, pp. 262-263.

CUNNINGHAM, ROBERT OLIVER.

1870. Notes on the Reptiles, Amphibia, Fishes, Mollusca, and Crustacea obtained during the voyage of H. M. S. *Nassau* in the years 1866-69. Trans. Linn. Soc. London, vol. 27, pp. 465-502, 2 pls.

1871. Notes on the natural history of the Strait of Magellan and west coast of Patagonia, made during the voyage of H. M. S. *Nassau* in the years 1866, 67, 68, & 69, xvii+517 pp., 20 pls., 1 map. Edinburgh.

CZERNIAVSKY, VOLDEMAR.

1884. Materialia ad zoographiam Ponticam comparatam, fasc. 2: Crustacea decapoda Pontica littoralia. Trav. Soc. Univ. Kharkov, vol. 13, suppl., 268 pp., 7 pls.

DANA, JAMES DWIGHT:

1851. On the classification of the Crustacea Grapsoidea. Amer. Journ. Sci. and Arts, vol. 12, ser. 2, pp. 283-291.

- 1852a. Conspectus crustaceorum quae in orbis terrarum circumnavigatione, Carolo Wilkes e classe Reipublicae Faederatae duce. Proc. Acad. Nat. Sci. Philadelphia, vol. 5, 1850-51, pp. 247-254.

- 1852b. Crustacea (pt. 1). U. S. Exploring Expedition, vol. 13, pp. i-viii, 1-685.

1853. Crustacea (pt. 2). U. S. Exploring Expedition, vol. 14, pp. 686-1618.

1854. On the geographical distribution of Crustacea. Amer. Journ. Sci. and Arts, ser. 2, vol. 18, pp. 1-45, 1 map.

1855. Crustacea. U. S. Exploring Expedition, atlas, 27 pp., 96 pls.

DE KAY, JAMES ELSWORTH.

1844. Zoology of New-York or the New-York fauna, pt. 6: Crustacea, 70 pp, 13 pls. Albany.

DOFLEIN, FRANZ.

1904. Brachyura, Wiss. Ergebn. Deutsch. Tiefsee-Exped. *Valdivia*, vol. 6, xiv+314 pp., 68 figs., 58 pls.

DOFLEIN, FRANZ, and BALSS, HEINRICH.

1912. Die dekapoden und stomatopoden der Hamburger Magalhaensischen Sammelreise 1892/93. Jahrb. Hamburg. Wiss. Anst., vol. 29, pp. 25-44, 4 figs.

EDMONDSON, CHARLES HOWARD.

1946. Reef and shore fauna of Hawaii. Bernice P. Bishop Museum Special Publ. 22, iii+381 pp., 223 figs.

EKMAN, SVEN.

1935. Tiergeographie des meeres, xii+542 pp., 244 figs. Leipzig.

FABRICIUS, JOHANNES CHRISTIAN.

1775. Systema entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus, 832 pp. Flensburg and Leipzig.

1781. Species insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosin adjectis observationibus, descriptionibus, vol. 1, viii+552 pp. Hamburg and Kiel.

1787. Mantissa insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus, vol. 1, xviii+348 pp. Copenhagen.

1793. Entomologia systematica, emendata et aucta. Classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus, vol. 2, viii+519 pp. Copenhagen.

FAXON, WALTER.

1895. Reports on an exploration off the west coasts of Mexico, Central and South America, and off the Galapagos Islands, in charge of Alexander Agassiz, by the U. S. Fish Commission steamer, *Albatross*, during 1891, Lieut.-Commander Z. L. Tanner, U. S. N., commanding, 15: The stalk-eyed Crustacea. Mem. Mus. Comp. Zool., vol. 18, 292 pp., 6 figs., 66 pls., 1 chart.

FILHOL, HENRI.

1885. Catalogue des crustacés de la Nouvelle-Zélande, des Iles Auckland et Campbell. Recueil de mémoires, rapports et documents relatifs à l'observation du passage de Vénus sur le soleil. Mission de l'Ile Campbell. Zoologie, vol. 3, pt. 2, No. 1, pp. 349-510, 18 pls. Paris.

FOWLER, HENRY WEED.

1912. The Crustacea of New Jersey. Ann. Rep. New Jersey State Mus., 1911, pt. 2, pp. 29-650, 149 pls.

FULTON, S. W., and GRANT, F. E.

1906. Census of the Victorian decapod Crustacea, pt. 1, (Brachyura). Proc. Roy. Soc. Victoria, new ser., vol. 19, pt. 1, art. 3, pp. 16-20.

GIBBES, LEWIS R.

1850. On the carcinological collections of the cabinets of natural history in the United States, with an enumeration of the species contained therein, and descriptions of new species, 37 pp. Charleston.

GLASSELL, STEVE ANDREW.

1934. Affinities of the brachyuran fauna of the Gulf of California. Journ. Washington Acad. Sci., vol. 24, No. 7, pp. 296-302.

GOODSIR, HARRY D. S.

1845. Description of some animals found amongst the gulf-weed. Ann. Mag. Nat. Hist., vol. 15, No. 96, pp. 73-76, 1 pl.

GOSSE, PHILIP HENRY.

1855. A manual of marine zoology for the British Isles, xi+203 pp., 335 figs. London.

GOURRET, PAUL.

1888. Révision des crustacés podophthalmes de Golfe de Marseille, suivie d'un essai de classification de la class de crustacés. Ann. Mus. Hist. Nat. Marseille, Zool., vol. 3, pp. 1-212, 18 pls.

HAAN, W. DE.

1835. Crustacea. In Siebold's "Fauna Japonica," decas 2, pp. 25-64, 9 pls. Leiden.

HALE, HERBERT M.

1927. The crustaceans of South Australia, 201 pp., 202 figs. Adelaide.

HASWELL, WILLIAM AITCHESON.

1882. Catalogue of the Australian stalk- and sessile-eyed Crustacea, xxiv+324 pp., 8 figs., 4 pls. Sydney.

HAY, WILLIAM PERRY, and SHORE, C. A.

1918. The decapod crustaceans of Beaufort, N. C., and the surrounding region. Bull. U. S. Bur. Fish., vol. 35, pp. 371-475, 20 figs., 15 pls.

HEILPRIN, ANGELO.

1889. Contributions to the natural history of the Bermuda Islands. Proc. Acad. Nat. Sci. Philadelphia, 1888, pp. 302-328, 3 pls.

HELLER, CAMIL.

1863. Die Crustaceen des südlichen Europa, xi+336 pp., 10 pls. Vienna.

1865. Crustaceen. Reise der österreichischen Fregatte *Novara* um die Erde in den Jahren 1857, 1858, 1859 unter den Befehlen des Commodore B. von Wüllerstorff-Urbair: Zool. Theil, vol. 2, pt. 3, 280 pp., 25 pls. Vienna.

HERBST, JOHANN FRIEDRICH WILHELM.

1783. Versuch einer Naturgeschichte der Krabben und Krebse nebst einer systematischen Beschreibung ihrer verschiedenen Arten, vol. 1, Heft 2-5, pp. 87-182, 8 pls. Berlin and Stralsund.

HITCHCOCK, EDWARD.

1835. Catalogues of the animals and plants of Massachusetts, with a copious index, 142 pp. Amherst.

HITCHCOCK, HAROLD BRADFORD.

1941. The coloration and color changes of the gulf-weed crab, *Planes minutus*. Biol. Bull. Woods Hole, vol. 80, No. 1, pp. 26-30.

HODGSON, T. V.

1896. *Nautilograpsus minutus*, Milne Edwards: Faunistic notes. Journ. Mar. Biol. Assoc. United Kingdom, new ser., vol. 4, No. 2, p. 178.

HOWE, FREELAND, JR.

1900. Report of a dredging expedition off the southern coast of New England, September, 1899. Bull. U. S. Fish Comm., vol. 19, 1899, pp. 237-240.

HYMAN, ORREN WILLIAMS.

1924. Studies on larvae of crabs of the family Grapsidae. Proc. U. S. Nat. Mus., vol. 65, art. 10, 8 pp., 3 pls.

JACQUINOT, HONORÉ and LUCAS, HIPPOLYTE.

1853. Crustacés. In Hombron and Jacquinot, "Voyage au pôle sud et dans l'océanie sur les corvettes l'*Astrolabe* et la *Zélée*": Zool., vol. 3, 107 pp. Paris.

KINGSLEY, JOHN STERLING.

1881. Carcinological notes, No. IV: Synopsis of the Grapsidae. Proc. Acad. Nat. Sci. Philadelphia, 1880, pp. 187-224.

KRAUSS, FERDINAND.

1843. Die Südafrikanischen Crustaceen, 68 pp., 4 pls. Stuttgart.

LATREILLE, PIERRE ANDRÉ.

1802-03. Histoire naturelle, générale et particulière, des crustacés et des insectes, vol. 6, 391 pp., 14 pls. Paris.

LEACH, WILLIAM ELFORD.

1814. Crustaceology. In Brewster's "Edinburgh Encyclopedia," vol. 7, pp. 383-437.

1815. Malacostraca podophthalmata Britanniae, Nos. 18-19, pls. 9A, 24A, 27, 34, 35, 37C, 45. London.

LEBOUR, MARIE VICTORIA.

1944. Larval crabs from Bermuda. Zoologica, New York, vol. 29, pt. 3, pp. 113-128, 19 figs.

LENZ, HEINRICH.

1901. Ergebnisse einer Reise nach dem Pacific (Schauinsland 1896-1897): Crustaceen. Zool. Jahrb., Abt. Syst., vol. 14, pp. 429-482, 1 pl.

1910. Crustaceen von Madagaskar, Ostafrika und Ceylon. In Alfred Voeltzkow's "Reise in Ostafrika in den Jahren 1903-1905 mit Mitteln der Hermann und Elise geb. Heckmann Wentzel-Stiftung ausgeführt," Wiss. Ergebn., vol. 2, Syst. Arb., pp. 539-576, 4 figs.

LENZ, HEINRICH, and STRUNCK, K.

1914. Die Dekapoden der Deutschen Südpolar-Expedition 1901-1903, I: Brachyuren und Macruren mit ausschluß der Sergestiden. Deutsche Südpolar-Expedition 1901-1903, vol. 15, Zool. 7, pp. 261-345, 5 figs., 11 pls.

LINNAEUS, CARL.

1747. Wästgöta-Resa, På Riksens Högloflige Ständers Befallning Förrättad år 1746, 284 pp., 5 pls. Stockholm.
1758. Systema naturae, ed. 10, vol. 1, 824 pp. Stockholm.
1767. Systema naturae, ed. 12, vol. 1, pt. 2, pp. 533-1327. Stockholm.

LUEDERWALDT, HERMANN.

1919. Lista dos crustaceos superiores (Thoracostraca) do Museu Paulista que foram encontrados no Estado de S. Paulo. Rev. Mus. Paulista, vol. 11, pp. 429-435.

LUKIS and LE LIEVRE.

1862. Stalk-eyed crustaceans. In Ansted and Latham, "The Channel Islands," pp. 232-234. London.

MACLEAY, WILLIAM SHARP.

1838. Illustrations of the Annulosa of South Africa, 75 pp., 4 pls. London.

MAN, JOHANNES GOVERTUS DE.

1879. On some new or imperfectly known podophthalmous Crustacea of the Leyden Museum. Notes Leyden Mus., vol. 19, pp. 53-73.

MARCUSEN, JOHANN.

1867. Zur Fauna des schwarzen Meeres. Vorläufige Mittheilung. Arch. Naturg., vol. 1, No. 4, pp. 357-363.

MARKHAM, CLEMENTS ROBERT.

1893. The journal of Christopher Columbus (during his first voyage, 1492-93), and documents relating to the voyages of John Cabot and Gaspar Corte Real, liv+240 pp. Hakluyt Society, London.

MAYER, ALFRED GOLDSBOROUGH.

1905. Sea-shore life: The invertebrates of the New York coast. New York Aquarium Nature Series, No. 1, 181 pp., 119 figs.

MCANDREW, ROBERT.

1860. List of the British marine invertebrate fauna, 71 pp. London.

MCINTOSH, WILLIAM CARMICHAEL.

1901. The coloration of marine animals. Ann. Mag. Nat. Hist., ser. 7, vol. 7, pp. 221-240.

MIERS, EDWARD J.

1876. Catalogue of the stalk- and sessile-eyed Crustacea of New Zealand, xii+136 pp., 3 pls. London.
1886. Report on the Brachyura: Report on the Scientific Results of the Voyage of H. M. S. *Challenger*, Zoology, vol. 17, pt. 49, 1+362 pp., 29 pls.

MILNE-EDWARDS, ALPHONSE.

1862. Faune carcinologique. In Maillard's "Notes sur l'Île de la Réunion (Bourbon)," pp. F1-F16, 3 pls. Paris.
1884. L'expédition du *Talisman* faite dans l'océan Atlantique sous les auspices des ministres de la marine et de l'instruction publique. Bull. Hebdomadaire Assoc. Sci. France, 1884, pp. 1-31.

MILNE-EDWARDS, ALPHONSE, and BOUVIER, EUGÈNE LOUIS.

1894. Crustacés décapodes provenant des campagnes du yacht *l'Hirondelle* (1886, 1887, 1888), pt. li Brachyures et anomures. Rés. Camp. Sci. Monaco, fasc. 7, 112 pp., 4 figs., 11 pls.

1899. Crustacés décapodes provenant des campagnes de l'*Hirondelle* (supplément) et de la *Princesse-Alice* (1891-1897). Rés. Camp. Sci. Monaco, fasc. 13, 106 pp., 4 pls.
1900. Crustacés décapodes, pt. 1: Brachyures et anomures. Expéd. Sci. Travailleur et Talisman, 396 pp., 32 pls. Paris.
- MILNE-EDWARDS, HENRI.
1837. Histoire naturelle des crustacés, vol. 2, 532 pp. Paris.
1852. Observations sur les affinités zoologiques et la classification naturelle des crustacés. Premier mémoire: Remarques sur la manière de caractériser les groupes zoologiques, et sur les principaux types secondaires qui dérivent du type classique de l'animal crustacé. Ann. Sci. Nat., ser. 3, vol. 18, pp. 109-196, 9 pls.
- MILNE-EDWARDS, HENRI, and LUCAS, HIPPOLYTE.
1843. Crustacés. In d'Orbigny's "Voyage dans l'Amérique Méridionale," vol. 6, pt. 1, pp. 1-39, 19 pls. Paris.
- MIRANDA Y RIVERA, ALVARO DE.
1921. Algunos crustáceos de la colección del Laboratorio Biológico de Málaga. Bol. Pescas, Madrid, pp. 179-205, 3 figs.
- MONOD, THÉODORE.
1927. Contribution à la faune du Cameroun: Crustacea, IV: Decapoda (excl. Palaemonidae, Atyidae et Potamonidae). Faun. Colon. Française, vol. 1, pp. 593-624, 3 figs.
1932. Brachyura maroccana, pt. 2: Dromiidea, Oxystomata, Oxyrhyncha, Brachyrhyncha (excl. Pinnoteridae). Bull. Soc. Sci. Nat. Maroc, vol. 12, Nos. 7-8, pp. 199-220, 7 figs.
1933. Sur quelques crustacés de l'Afrique Occidentale. (Liste des décapodes mauritaniens et des xanthidés ouestafricains.) Bull. Com. Étud. Hist. Sci. Afr. Occ. Française, vol. 15, Nos. 2-3, 1932, pp. 1-93, 26 figs.
1939. Sur quelques crustacés de la Guadeloupe (Mission P. Allorge, 1936). Bull. Mus. Hist. Nat. Paris, ser. 2, vol. 11, No. 6, pp. 557-568, 11 figs.
- MORISON, SAMUEL ELIOT.
1942. Admiral of the ocean sea: A life of Christopher Columbus, vol. 1, xlv+448 pp. Boston.
- MOSELEY, HENRY NOTTIGE.
1879. Notes by a naturalist on the *Challenger*, being an account of various observations made during the voyage of H. M. S. *Challenger* round the world, xvi+620 pp., figs. London.
- MÜLLER, FRITZ.
1881. Farbenwechsel bei Krabben und Garneelen. Kosmos, Leipzig, vol. 8, fasc. 12, pp. 472-473.
- MÜLLER, OTTO FREDERIK.
1776. Zoologiae Danicae Prodrumus, seu Animalium Daniae et Norvegiae indigenarum characteres, nominam et synonyma imprimis popularium, xxxii+282 pp. Copenhagen.
- MURRAY, JOHN, and HJORT, JOHAN.
1912. Depths of the ocean, xx+821 pp., 575 figs., 9 pls., 1 map. London.
- NICOLET, HERCULE.
1849. Crustaceos. In Gay's "Historia Fisica y Politica de Chile," Zool., vol. 3, pp. 115-318. Paris.
- NOBILI, GIUSEPPE.
1906. Faune carcinologique de la Mer Rouge: Décapodes et Stomatopodes. Ann. Sci. Nat., Zool., ser. 9, vol. 4, pp. 1-347, 12 figs., 11 pls.

NOBRE, AUGUSTO.

1931. Crustáceos decápodes e stomatópodes marinhos de Portugal, iv+307 pp., 144 figs., 2 pls. Pôrto.

1936. Fauna marinha de Portugal: IV, Crustáceos decápodes e stomatópodes marinhos de Portugal, viii+213 pp., 61 pls. Pôrto.

NORMAN, ALFRED MERLE.

1907. Notes on the Crustacea of the Channel Islands. *Ann. Mag. Nat. Hist.*, ser. 7, vol. 20, pp. 356-371, 2 pls.

NORMAN, ALFRED MERLE, and BRADY, GEORGE STEWARDSON.

1909. The Crustacea of Northumberland and Durham. *Trans. Nat. Hist. Soc. Northumberland, Durham, and Newcastle-upon-Tyne*, new ser., vol. 3, pt. 2, 168 pp., 2 pls.

NORMAN, A. M., and SCOTT, THOMAS.

1906. The Crustacea of Devon and Cornwall, xv+232 pp., 24 pls. London.

NUTTING, CHARLES CLEVELAND.

1895. Narrative and preliminary report of Bahama Expedition. *Bull. Lab. Nat. Hist. State Univ. Iowa*, vol. 3, Nos. 1-2, vii+251 pp., 20 pls.

OLIVIER, GUILLAUME ANTOINE.

1791. *Encyclopedie methodique: Histoire naturelle: Insectes*, vol. 6, 704 pp. Paris.

ORTMANN, ARNOLD EDWARD.

1893. Decapoden und schizopoden. *Ergebn. Plankton-Exped.*, vol. 2, 120 pp., 1 fig., 10 pls.

1894. Die Decapoden-Krebse des Strassburger Museums. VIII. Theil. Abtheilung: Brachyura (*Brachyura genuina* Boas). III. Unterabtheilung: Cancroidea, 2. Section: Cancrinea, 2. Gruppe: Catometopa. *Zool. Jahrb., Abt. Syst.*, vol. 7, pp. 683-772, 1 pl.

1899. Gliederfüssler: Arthropoda. In Bronn's "Klassen und ordnungen des Thier-reichs," vol. 5, Abt. 2, Lief. 53-56, pp. 1169-1232. Leipzig.

OSBECK, PETER.

1771. *A voyage to China and the East Indies*, vol. 2, 152 pp. London. [Translated by J. R. Forster.]

PENNANT, THOMAS.

1777. Crustacea, Mollusca, Testacea. *British Zoology*, vol. 4, xx+154 pp., 93 pls.

PERKINS, MICHAEL.

1928. A simple key to the crabs (*Brachyura*) of Britain and the north-east Atlantic. *Scottish Nat.*, Mar.-Apr., 1928, pp. 53-59; May-June, 1928, pp. 87-94.

PESTA, OTTO.

1912. Die Decapoden-Krebse der Adria in Bestimmungstabellen zusammengestellt. *Arch. Naturg.*, Jahrg. 78, Abt. A, Heft 1, pp. 93-126.

1918. Die Decapodenfauna der Adria. Versuch einer Monographie, x+500 pp., 150 figs., 1 chart. Leipzig and Vienna.

POPE, ELIZABETH C.

1948. Hitch-hikers of the sea. *Australian Mag.*, vol. 9, No. 9, pp. 293-297.

POUCHET, GEORGES, and GUERNE, JULES DE.

1886. Sur l'alimentation des tortues marines. *Comptes Rendus Acad. Sci. Paris*, vol. 102, No. 15, pp. 877-879.

RANKIN, WALTER MEAD.

1900. The Crustacea of the Bermuda Islands, with notes on collections made by the New York University Expeditions in 1897 and 1898. *Ann. New York Acad. Sci.*, vol. 12, No. 12, pp. 521-548, 1 pl.

RATHBUN, MARY JANE.

- 1897a. A revision of the nomenclature of the Brachyura. *Proc. Biol. Soc. Washington*, vol. 11, pp. 153-167.
- 1897b. List of the decapod Crustacea of Jamaica. *Ann. Inst. Jamaica*, vol. 1, No. 1, 46 pp.
- 1898a. The Brachyura of the biological expedition to the Florida Keys and the Bahamas in 1893. *Bull. Lab. Nat. Hist. State Univ. Iowa*, vol. 4, pp. 250-294, 9 pls.
- 1898b. The Brachyura collected by the U. S. Fish Commission Steamer *Albatross* on the voyage from Norfolk, Virginia, to San Francisco, California, 1887-1888. *Proc. U. S. Nat. Mus.*, vol. 21, pp. 567-616, 4 pls.
- 1900a. The decapod crustaceans of West Africa. *Proc. U. S. Nat. Mus.*, vol. 22, pp. 271-316, 2 figs.
- 1900b. Synopsis of North-American invertebrates, XI: The catometopus or grapsoid crabs of North America. *Amer. Nat.*, vol. 34, No. 403, pp. 583-592, 15 figs.
1902. Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899, VIII: Brachyura and Macrura. *Proc. Washington Acad. Sci.*, vol. 4, pp. 275-292, 4 figs., 1 pl.
1904. Decapod crustaceans of the northwest coast of North America. *Harri-man Alaska Exped.*, vol. 10, pp. 1-190, 95 figs., 10 pls.
1905. Fauna of New England, 5: List of Crustacea. *Occas. Pap. Boston Soc. Nat. Hist.*, vol. 7, pp. 1-117.
1906. The Brachyura and Macrura of the Hawaiian Islands. *Bull. U. S. Fish Comm.*, 1903, pt. 3, pp. 827-930, i-viii, 79 figs., 24 pls.
1907. Reports on the scientific results of the expedition to the tropical Pacific, in charge of Alexander Agassiz, by the U. S. Fish Commission Steamer *Albatross*. etc., IX: Reports on the scientific results of the expedition to the eastern tropical Pacific, in charge of Alexander Agassiz, by the U. S. Fish Commission Steamer *Albatross*, etc. X: The Brachyura. *Mem. Mus. Comp. Zool.*, vol. 35, No. 2, pp. 23-74, 9 pls.
1910. The stalk-eyed Crustacea of Peru and the adjacent coast. *Proc. U. S. Nat. Mus.*, vol. 38, pp. 531-620, 21 pls.
1914. New genera and species of American brachyrhynchous crabs. *Proc. U. S. Nat. Mus.*, vol. 47, pp. 117-129, 5 figs., 10 pls.
1918. The grapsoid crabs of America. *U. S. Nat. Mus. Bull.* 97, xxii+461 pp., 172 figs., 161 pls.
1924. Brachyuran crabs collected by the Williams Galápagos Expedition, 1923. *Zoologica*, New York, vol. 5, No. 14, pp. 153-159, 1 fig., 1 pl.
1929. Canadian Atlantic fauna, 10: Arthropoda; 10m: Decapoda, 38 pp., 53 figs. St. Andrews.
1933. Brachyuran crabs of Porto Rico and the Virgin Islands. *Sci. Surv. Porto Rico and Virgin Ids.*, New York Acad. Sci., vol. 15, pt. 1, pp. 1-121, 107 figs.

RICHARDSON, L. R.

1949. A guide to the brachyrhynchous crabs. *Tuatara*, New Zealand, vol. 2, No. 1, pp. 29-36, 2 pls.

ROUX, POLYDORE.

1828. *Crustacés de la Méditerranée et de son littoral*, 176 pp., 45 pls. Paris.

SAKAI, TUNE.

1934. *Brachyura* from the coast of Kyusyu, Japan. *Sci. Rep. Tokyo Bunrika Daigaku*, sec. B, vol. 1, No. 25, pp. 281-330, 26 figs., 2 pls.
1936. *Crabs of Japan, 1935*, 239 pp., 122 figs., 66 pls. Tokyo.
1939. *Studies on the crabs of Japan, IV: Brachygnatha, Brachyrhyncha*, pp. 365-741, 1-23, 129 figs., 70 pls. Tokyo.

SAY, THOMAS.

1817. *An account of the Crustacea of the United States*. *Journ. Acad. Nat. Sci. Philadelphia*, vol. 1, No. 6, pp. 97-102.
1818. *Observations on some of the animals described in the account of the Crustacea of the United States*. *Journ. Acad. Nat. Sci. Philadelphia*, vol. 1, pp. 442-444.

SCHMITT, WALDO LASALLE.

1921. *The marine decapod Crustacea of California*. *Univ. California Publ. Zool.*, vol. 23, 470 pp., 165 figs., 50 pls.

SIVERTSEN, ERLING.

1933. *The Norwegian Zoological Expedition to the Galápagos Islands 1925, conducted by Alf Wollebaek, VII: Littoral Crustacea Decapoda from the Galápagos Islands*. *Medd. Zool. Mus. Oslo*, No. 38, 23 pp., 4 pls.

SLOANE, HANS.

1725. *A voyage to the islands of Madera, Barbados, Nieves, St. Christophers, and Jamaica*, vol. 2, xviii+499 pp., 125 pls. London.

SMITH, SIDNEY IRVING.

1879. *The stalk-eyed crustaceans of the Atlantic coast of North America north of Cape Cod*. *Trans. Connecticut Acad. Arts and Sci.*, vol. 5, pt. 1, pp. 27-136, 5 pls.
1880. *Occasional occurrence of tropical and sub-tropical species of decapod Crustacea on the coast of New England*. *Trans. Connecticut Acad. Arts and Sci.*, vol. 4, pp. 254-267.
1886. *Report on the decapod Crustacea of the Albatross dredgings off the east coast of the United States during the summer and autumn of 1884*. *Ann. Rep. U. S. Fish Comm.*, 1885, pp. 605-705, 20 pls.

SMITH, S. I., and HARGER, OSCAR.

1874. *Report on the dredgings in the region of St. George's Banks, in 1872*. *Trans. Connecticut Acad. Arts and Sci.*, vol. 3, pt. 1, pp. 1-57, 8 pls.

STEBBING, THOMAS ROSCOE REDE.

1893. *A history of Crustacea. Recent Malacostraca*, xvii+466 pp., 32 figs., 19 pls. New York.
1905. *South African Crustacea, part 3: Marine Investigations in South Africa*, vol. 4, pp. 21-123, 10 pls.
1910. *General catalogue of South African Crustacea*. [Pt. 5: of South African Crustacea, for the Marine Investigations in South Africa.] *Ann. South African Mus.*, vol. 6, pt. 4, pp. 281-593, 8 pls.
- 1914a. *Crustacea from the Falkland Islands collected by Mr. Rupert Vallentin, F. L. S.*, pt. 2. *Proc. Zool. Soc. London*, 1914, pp. 341-378, 9 pls.
- 1914b. *Stalk-eyed Crustacea Malacostraca of the Scottish National Antarctic Expedition*. *Trans. Roy. Soc. Edinburgh*, vol. 50, pt. 2, No. 9, pp. 253-307, 10 pls.

STEINBECK, JOHN, and RICKETTS, EDWARD FLANDERS.

1941. *Sea of Cortez*, x+598 pp., 40 pls., 2 charts. New York.

STIMPSON, WILLIAM.

1858. Prodromus descriptionis animalium evertibratorum, quae in Expeditione ad Oceanum Pacificum Septentrionalem, a Republica Federata missa, Cadwaladaro Ringgold et Johanne Rodgers Ducibus, observavit et descripsit, pt. 5: Crustacea Ocypodoidea. Proc. Acad. Nat. Sci. Philadelphia, vol. 10, pp. 93-111 [39-56].

1860. Notes on North American Crustacea, in the Museum of the Smithsonian Institution, No. II: Ann. Lyc. Nat. Hist. New York, vol. 7, pp. 177-246 [49-118], 2 pls.

1907. Report on the Crustacea (Brachyura and Anomura) collected by the North Pacific Exploring Expedition, 1853-1856. Smithsonian Misc. Coll., vol. 49, pp. 1-240, 26 pls.

STOSSICH, MICHELE.

1880. Prospetto della fauna del mare Adriatico, pt. 3. Boll. Soc. Adriatica Sci. Nat. Trieste, vol. 6, pp. 178-271.

SUMNER, FRANCIS B.; OSBURN, RAYMOND C.; and COLE, LEON J.

1913. A biological survey of the waters of Woods Hole and vicinity, pt. 2, sec. 3: A catalogue of the marine fauna. Bull. U. S. Bur. Fish., vol. 31, pt. 2, pp. 547-794.

TARGIONI TOZZETTI, ADOLFO.

1872. Catalogo di crostacei podottalmi brachiuri e anomouri raccolti nel viaggio di circumnavigazione della Fregata italiana *Magenta*. Atti. Soc. Ital., Milan, vol. 15, pp. 1-12.

1877. Zoologia del viaggio intorno al globo della R. Pirocorvetta *Magenta* durante gli anni 1865-68. Crostacei brachiuri e anomouri. Pubblicazioni del R. Istituto di Studi Superiori pratici e di Perfezionamento in Firenze, Sezione di Scienze Fisiche e Naturali, vol. 1, xxix+257 pp., 3 figs., 13 pls.

TESCH, JOHAN JACOB.

1918. The Decapoda Brachyura of the Siboga Expedition, I: Hymenosomidae, Retroplumidae, Ocypodidae, Grapsidae and Gecarcinidae. *Siboga-Exped.*, monogr. 39^c, 148 pp., 6 pls.

THOMPSON, D'ARCY WENTWORTH.

1901. A catalogue of Crustacea and of Pycnogonida contained in the Museum of University College, Dundee, v+56 pp. Dundee.

THOMPSON, J. V.

1836. Of the double metamorphosis in *Macropodia phalangium*, or spider-crab, with proofs of the larvae being zoëa in *Gegarcinus hydrodromus*, *Thelphusa erythropus*, *Eriphia carribaea*, and *Grapsus pelagicus*. Ent. Mag., vol. 3, art. 33, pp. 370-375, 6 figs.

URITA, TOMOYE.

1926. A check list of Brachyura found in Kagoshima Prefecture, Japan, iii+41 pp. Tsingtao.

VERRILL, ADDISON EMORY.

1885. Results of the explorations made by the steamer *Albatross* off the northern coast of the United States, in 1883. Ann. Rep. U. S. Fish. Comm., 1883, pp. 503-699 [1-197], 44 pls.

1908. Decapod Crustacea of Bermuda I: Brachyura and Anomura, their distribution, variations, and habits. Trans. Connecticut Acad. Arts and Sci., vol. 13, pp. 299-474, 68 figs., 20 pls.

WAGNER, MORITZ.

1880. Über die Entstehung der Arten durch Absonderung, II: Die Mimicry. Kosmos, Leipzig, vol. 7, Heft 2, pp. 89-99.

WARD, MELBOURNE.

1939. The Brachyura of the Second Templeton Crocker-American Museum Expedition to the Pacific Ocean. Amer. Mus. Nov., No. 1049, 15 pp., 17 figs.

WEYMOUTH, FRANK WALTER.

1910. Synopsis of the true crabs (Brachyura) of Monterey Bay, California. Leland Stanford Junior Univ. Publ., univ. ser., No. 4, 64 pp., 9 figs., 14 pls.

WHITE, ADAM.

1847. List of the specimens of Crustacea in the collection of the British Museum, viii+143 pp. London.
1850. List of the specimens of British animals in the collection of the British Museum, pt. 4: Crustacea, iv+141 pp. London.

YOUNG CHARLES G.

1900. The stalk-eyed Crustacea of British Guiana, West Indies, and Bermuda, xix+514 pp., figs., 7 pls. London.

ZARIQUIEY ALVAREZ, RICARDO.

1946. Crustáceos Decápodos Mediterráneos. Manual para la clasificación de las especies que pueden capturarse en las costas mediterráneas españolas. Inst. Español Estudios Mediterráneos Publ. Biología Mediterránea, vol. 2, 181 pp., 174 figs., 26 pls.

Chace, Fenner A. 1951. "The oceanic crabs of the genera *Planes* and *Pachygrapsus*." *Proceedings of the United States National Museum* 101(3272), 65-103.

View This Item Online: <https://www.biodiversitylibrary.org/item/31795>

Permalink: <https://www.biodiversitylibrary.org/partpdf/30772>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.