PIPRA CAUDATA.

CHARACTER GENERICUS.

Rostrum capite brevius, basi subtrigonum, integerrimum, apice incurvum.

Pedes gressorii.

Lin. Syst. Nat. p. 338.

CHARACTER SPECIFICUS.

PIPRA CÆRULEA, vertice subcristato coccineo, alis nigris, rectricibus duabus intermediis elongato-acuminatis.

Colli caudæque latera viridi-splendentia.

Aviculam in suo genere proculdubio pulcherrimam et coloribus eximiis superbientem, simulque novam nec antea descriptam, magnitudine naturali depictam, sistit tabula. In America Australi et calidiori, sicut aliæ pleræque congeneres, præcipue invenitur.

THE
LONG-TAILED MANAKIN.

GÉNÉRIC CHARACTER.

Bill shorter than the head, somewhat triangular at the base, bent at the tip.

Feet gressorial.

SPECIFIC CHARACTER.

BLUE MANAKIN with scarlet crown, black wings, and the two middle tail-feathers elongated and acuminate.

The sides of the neck and tail have a green lustre.

The bird here represented, which is unquestionably the most brilliant of its tribe, and is besides a new and hitherto undescribed species, is figured in its natural size. Like the rest of its genus it is principally found in the warmer parts of South America.

Shaw, George. 1793. "The Long-Tailed Manakin, *Pipra caudata* [Pl. 153]." *The Naturalist's Miscellany* 5(L), <https://doi.org/10.5962/p.310732>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276340>

DOI: <https://doi.org/10.5962/p.310732>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310732>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.