MEDUSA CEPHEA.

CHARACTER GENERICUS.

Corpus gelatinosum, orbiculatum, depressum.

Os subtus, centrale.

Lin. Syst. Nat. p. 1096.

CHARACTER SPECIFICUS, &c.

MEDUSA hemisphærico-depressa fusco-rufescens,
subtus brachiis octo apice villosis, tentacu-
lisque novem filiformibus longissimis.

MEDUSA hemisphærica tuberculata fusco-rufes-
cens, margine crenato, brachiis subtus octo
lanatis.

Lin. Syst. Nat. Gmel. p. 3158.

Forsk. Fn. Aegypt. Arab. p. 108. n. 22.

Medusa Cephea, quæ inter majores numeratur,
pellucida admodum est, et gelata, palletque fusco-
rufescens, in nonnullis partibus obscurior et fere
subcærulea. Circa litora maris rubri, non infre-
quens reperitur.

THE
CEPHEAN MEDUSA.

GENERIC CHARACTER.

Body gelatinous, orbicular, commonly depressed.
Mouth central, beneath.

SPECIFIC CHARACTER, &c.

HEMISPHERIC-DEPRESSED REDDISH-
BROWN MEDUSA; furnished with eight
arms, villous at their extremities, and with
nine very long filiform tentacula.

This is one of the larger Medusæ: it is extremely pellucid, of a gelatinous consistence, and of a pale reddish-brown color, tinged in some parts with a darker hue, approaching nearly to blueish. It is not uncommon towards the coasts of the Red Sea.

Shaw, George. 1795. "The Cephean Medusa, Medusa cephea [Pl. 224]." *The Naturalist's Miscellany* 7(LXXIV), <https://doi.org/10.5962/p.310767>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276356>

DOI: <https://doi.org/10.5962/p.310767>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310767>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.