TANAGRA BRASILIA.

CHARACTER GENERICUS.

Rostrum conicum, acuminatum, emarginatum,
basi subtrigonum, apice declive.

Lin. Syst. Nat. p. 314.

CHARACTER SPECIFICUS, &c.

TANAGRA COCCINEA, alis caudaque nigris.

Lin. Syst. Nat. p. 314.

MERULA BRASILICA.

Bell. av. 319.

TIJEPIRANGA.

Raj. syn. p. 87. 13.

Will. orn. p. 184. 185.

CARDINALIS.

Briss. av. 3. p. 42. n. 24. t. 3. f. 1.

Pulcherrima hæc avis, cujus magnitudinem naturalem ostendit tabula, amat omnem Americam Australem, præcipue autem Brasiliam.

A. N.

London, Published Feb. 1846, by T. D. Nodder & Co. N. 15, Brewer Street.

THE
BRASILIAN TANAGER.

GENERIC CHARACTER.

Bill conic, sharp-pointed, flogging at the tip and slightly emarginated; somewhat trigonal towards the base.

SPECIFIC CHARACTER, &c.

SCARLET TANAGER with black wings and tail.

Le CARDINAL.

Briff. orn. 3. p. 42. 24. pl. 3. f. 1.

BRASILIAN TANAGER.

Lath. Syn. 1. p. 215.

This highly elegant species is a native of South America, and is principally found in Brasil. The plate represents it in its natural size.

Shaw, George. 1796. "The Brasilian Tanager, Tanagra brasilia [Pl. 234]." *The Naturalist's Miscellany* 7(LXXIX), <https://doi.org/10.5962/p.310777>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276356>

DOI: <https://doi.org/10.5962/p.310777>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310777>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.