LACERTA PALUSTRIS.

CHARACTER GENERICUS.

Corpus tetrapodum, caudatum, nudum.

Lin. Syst. Nat. p. 359.

CHARACTER SPECIFICUS, &c.

LACERTA nigricans, lateribus albido-punctatis,
abdomine croceo, maculis inæqualibus nigris.

SALAMANDRA alepidota verrucosa.

Gronov. Mus. 2. p. 77. n. 51.

SALAMANDRA aquatica.

Houttuyn. Aët. Ulyss. 9. p. 328.

Triton PALUSTRIS.

Laur. amphib. p. 39. n. 43. t. 4. f. 2.

Superat interdum magnitudine Lacerta palustris
specimina quæ in tabula ostenduntur; rarior et am-
plior lacerta aquatica, quam alunt fere omnes aquæ
stagnantes. Color corporis superioris fusco-nigricat
faturatior; infra aurantius, maculis magnis, inæqua-
libus, nigris, variatus. Latera verruculis plurimis
seu atomis convexis, albis, asperguntur; caudaque
utrinque fasciam plerumque habet latam, subargen-
team,

team, cum mistura quadam subcærulei. Affurgit
magis minusve per dorsum maris quasi crista conti-
nuata et compressa inæqualiter ferrata.

Amat ut plurimum, hæc species aquas puriores
et frigidiores quam quas incolit lacerta aquatica;
et ineunte vere, conspici possit vespertino tempore
in locis udis juxta margines aquarum, nec non sub
sepibus, aliisque latibulis, quæ umbram præbent et
humorem.

THE
WARTED NEWT.

GENERIC CHARACTER.

Body four-footed, tailed ; naked ; or unfurnished with any distinct integument.

SPECIFIC CHARACTER, &c.

Blackish NEWT with the sides speckled with white ; the abdomen orange with irregular black spots.

The Warty LIZARD.

Pennt. Brit. Zool. 3. p. 22. t. 3. fig. 1.

The black-and-orange WATER-NEWT.

The greater WATER-NEWT.

The *Lacerta palustris* is a species somewhat less common than the *lacerta aquatica*, which occurs in almost every stagnant water : it is also of a superior size ; being sometimes found considerably larger than the specimens here represented. Its color is an extremely deep blackish brown above, and a bright orange beneath, variegated with large irregular black spots. The sides of the body are sprinkled
over

over with numerous white convex specks or tubercles, and on each side of the tail there is commonly a broad streak of a silvery hue, with a cast of blueish. The back of the male is more or less distinguished by a kind of flattened process, divided into irregular serratures. This species is commonly met with in waters of a clearer and colder nature than those in which the *lacerta aquatica* is found; and in the beginning of spring may occasionally be observed, towards the evening, in damp situations near the margins of ponds, as well as under hedges, and other shady situations.

Shaw, George. 1797. "The Warted Newt, *Lacerta palustris* [Pl. 279]." *The Naturalist's Miscellany* 8(XCI), <https://doi.org/10.5962/p.310822>.

View This Item Online: <https://www.biodiversitylibrary.org/item/276320>

DOI: <https://doi.org/10.5962/p.310822>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310822>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.