CHÆTODON TRICOLOR.

CHARACTER GENERICUS.

Dentes setacei : corpus pictum.

Bloch. ichth. 6. p. 35.

CHARACTER SPECIFICUS, &c.

CHÆTODON antice flavus, postice niger, cauda
marginibusque pinnarum rubris.

CHÆTODON TRICOLOR.

Bloch. ichth. 12. p. 97. t. 425.

Americana est mirabilis hæc species, primusque
eam descripsisse videtur celeberrimus Blochius.
Crescere solet in longitudinem plus pedalem.

TRICOLOR CHÆTODON.

GENERIC CHARACTER.

Teeth setaceous. *Body* varied or banded.

SPECIFIC CHARACTER, &c.

Yellow CHÆTODON, with the hind part black;
the tail and fins bordered with red.

The TRICOLOR or triple-coloured CHÆTO-
DON.

This highly singular species, which grows to the length of a foot or more, is a native of the American seas, and seems to have been first described in the work of Dr. Bloch.

Shaw, George. 1800. "Tricolor Chætodon, Chætodon tricolor [Pl. 474]." *The Naturalist's Miscellany* 12(CXXXIII), <https://doi.org/10.5962/p.310920>.

View This Item Online: <https://www.biodiversitylibrary.org/item/278722>

DOI: <https://doi.org/10.5962/p.310920>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310920>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.