PAPILIO XUTHUS.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes, (volatu diurno.)

Lin. Syst. Nat. p. 774.

CHARACTER SPECIFICUS, &c.

PAPILIO alis nigris albo striato maculatis; posterioribus subtus ocellis cæruleis fulvisque subfasciatis.

Fab. sp. ins. 2. p. 19.

PAPILIO alis caudatis nigris albido striato-maculatis; posticis subtus cæruleo fulvoque subocellatis.

Lin. Syst. Nat. p. 751.

Inter Lepidoptera rariora numeratur Papilio Xuthus, Indiæ orientalis incola. Tabula magnitudine vera pictum exhibet.

21

484

XUTHUS.

GENERIC CHARACTER.

Antennæ thickening towards the upper part, and generally terminating in a clavated tip.

Wings (when sitting) erect, and meeting upwards. (Flight diurnal.)

SPECIFIC CHARACTER, &c.

Black subcaudated Butterfly, with yellowish white striated spots; the lower wings marked beneath with a band of blue and orange ocellated spots.

The *Papilio Xuthus* is numbered among the rarer species of *Lepidoptera*, and is a native of the East Indies. The plate represents it in its natural size.

Shaw, George. 1800. "Xuthus, Papilio xuthus [Pl. 484]." *The Naturalist's Miscellany* 12(CXXXV), <https://doi.org/10.5962/p.310930>.

View This Item Online: <https://www.biodiversitylibrary.org/item/278722>

DOI: <https://doi.org/10.5962/p.310930>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310930>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.