PAPILIO AGENOR.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes, (volatu diurno.)

Lin. Syst. Nat. p. 744.

CHARACTER SPECIFICUS, &c.

PAPILIO alis dentatis nigris basi sanguineis, primoribus striatis, posticis disco albo maculis nigris.

Lin. Syst. Nat. p. 747.

Clerk. ic. ins. rar. t. 15.

Cram. pap. 3. t. 32. f. A. B.

Jablonsky pap. 1. t. 8. f. 3.

In variis Asiæ regionibus generatur Papilio Agenor, magnitudine vera in tabula depictus.

London, Published Sept. 1st 1801, by Elizabeth & William, Sons of C. Newman Street.

AGENOR.

GENERIC CHARACTER.

Antennæ commonly thickening towards the end into a clavated tip.

Wings (when at rest) meeting upwards. (Flight diurnal.)

SPECIFIC CHARACTER.

Black Butterfly with dentated wings red at the base; the lower pair white on the disk, with black spots.

This elegant insect is found in several of the Asiatic regions, and is represented on the plate in its natural size.

Shaw, George. 1802. "Agenor, Papilio agenor [Pl. 500]." *The Naturalist's Miscellany* 13(CLIV), <https://doi.org/10.5962/p.310946>.

View This Item Online: <https://www.biodiversitylibrary.org/item/281546>

DOI: <https://doi.org/10.5962/p.310946>

Permalink: <https://www.biodiversitylibrary.org/partpdf/310946>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.