PAPILIO MEROPE.

CHARACTER GENERICUS.

Antennæ apicem versus crassiores, sæpius clavato-capitatae.

Alæ (sedentis) erectæ sursumque conniventes, (volatu-diurno.)

CHARACTER SPECIFICUS, &c.

PAPILIO alis flavis limbo nigro, posterioribus subtus fascia fusca.

PAPILIO Brutus.

Lin. Syst. Nat. Gmel. p. 2237.

PAPILIO MEROPE.

Cram. t. 378. f. D. E.

Africæ varias partes incolit Papilio Merope, cujus veram magnitudinem cernere est in tabula: coloribus interdum paulo variat.

MEROPE.

GENERIC CHARACTER.

Antennæ thickening towards the upper part, and generally terminating in a clavated tip.

Wings (when sitting) erect, and meeting upwards. (Flight diurnal.)

SPECIFIC CHARACTER, &c.

Butterfly with yellow wings bordered with black; the lower pair marked beneath by a transverse brown band.

P. MEROPE.

Cram. t. 378. f. D. E.

This species is a native of several parts of Africa, and is represented on the plate in its natural size. In colour it sometimes varies a little.

Shaw, George. 1803. "Merope, Papilio merope [Pl. 604]." *The Naturalist's Miscellany* 15(CLXXII), <https://doi.org/10.5962/p.311047>.

View This Item Online: <https://www.biodiversitylibrary.org/item/280634>

DOI: <https://doi.org/10.5962/p.311047>

Permalink: <https://www.biodiversitylibrary.org/partpdf/311047>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.