TANAGRA MISSISSIPENSIS.

CHARACTER GENERICUS.

Rostrum conicum, acuminatum, emarginatum,
basi subtrigonum, apice declive.

Lin. Syst. Nat. p. 313.

CHARACTER SPECIFICUS, &c.

TANAGRA tota rubra, rostro fusco.

TANAGRA tota rubra.

Lath. ind. orn. p. 421.

Cardinalis Mexicanus.

Briss. av. 3. p. 46.

Varias Americæ partes incolit hæc avis, magnitudine
Tanagram Brasiliensem æquans.

693

Red Bird. D. d. 2. June.

THE
MISSISSIPPI TANAGER.

GENERIC CHARACTER.

Bill conic, acuminated, a little inclining towards the point; the upper mandible slightly ridged, and notched near the end.

SPECIFIC CHARACTER, &c.

Red TANAGER, with brown beak.

MISSISSIPPI TANAGER.

Lath. Syn. 2. p. 218.

TANGARA du MISSISSIPPI.

Buff. 4. p. 252.

Pl. Enl. 741.

This species is found in several parts of America, and is equal in size to the Brazilian Tanager.

Shaw, George. 1806. "The Mississippi Tanager, *Tanagra mississippiensis* [Pl. 693]." *The Naturalist's Miscellany* 17(CC), <https://doi.org/10.5962/p.311131>.

View This Item Online: <https://www.biodiversitylibrary.org/item/281954>

DOI: <https://doi.org/10.5962/p.311131>

Permalink: <https://www.biodiversitylibrary.org/partpdf/311131>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.