SCYLLÆA PELAGICA.

CHARACTER GENERICUS.

Tentacula duo, foliacea.

Corpus subcompressum, sulco subtus longitudinali.

Lobi utrinque suberecti, foliacei, branchiis muricati.

CHARACTER SPECIFICUS, &c.

SCYLLÆA flavescens, lobis utrinque duobus suberectis foliaceis.

SCYLLÆA PELAGICA.

Lin. Syst. Nat. p. 1094.

Veram Scyllææ pelagicæ conformationem luculenter descripsit celeberrimus Cuvierus. In maribus generatur Indicis et Americanis, eadem fere magnitudine quam ostendit tabula. Color flavo-fusco pallescit.

R. P. Nodder. Sculpt.

THE
PELAGIC SCYLLÆA.

GENERIC CHARACTER.

Tentacula two, foliaceous.

Body subcompressed, with a longitudinal furrow beneath.

Foliaceous Lobes on each side, suberect, and mucated with branchiæ.

SPECIFIC CHARACTER, &c.

Yellowish SCYLLÆA, with two suberect foliaceous lobes on each side.

La Scyllée.

Cuvier, Ann. Mus. Hist. Nat. 36.

p. 416, &c.

This marine animal, of which the true structure has been so well ascertained and described by Mons. Cuvier, is found in the Atlantic and Indian seas, and is generally of the size represented in the plate. Its colour is a pale yellowish brown.

Shaw, George. 1805. "The Pelagic Scyllæa, Scyllæa pelagica [Pl. 728]." *The Naturalist's Miscellany* 17(CXCVI), <https://doi.org/10.5962/p.311165>.

View This Item Online: <https://www.biodiversitylibrary.org/item/281954>

DOI: <https://doi.org/10.5962/p.311165>

Permalink: <https://www.biodiversitylibrary.org/partpdf/311165>

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.