





## SULA FUSCA, *Briss.*

### Brown Gannet.

*Sula fusca*, Briss. Orn., tom. vi. p. 499. tab. 43. fig. 1.—Id. 8vo, tom. ii. p. 491.—Vieill. Gal. des Ois., tom. ii. pl. 277.—Bonap. Syn. Birds of Am., p. 408.—Nutt. Man., vol. ii. p. 500.—Aud. Birds of Am. vol. iii. pl. 207 ; Orn. Bio., vol. iii. p. 63 ; Syn. Birds of Am., p. 310.—Bonap. List of Eur. and Am. Birds, p. 60.

*Brown Booby*, Lath. Gen. Syn., vol. vi. p. 613.—Ib. Gen. Hist., vol. x. p. 441.

*Sula Brasiliensis*, Spix, Av. Sp. Nov., tom. ii. tab. cvii. p. 84.

*Sula fiber*, List of Birds in Brit. Mus. Coll., part iii. p. 183.

*Pelecanus fiber*, Linn. Syst. Nat., vol. i. p. 218.—Gmel. Edit. Linn. Syst. Nat., vol. i. p. 579.—Lath. Ind. Orn., vol. ii. p. 893.—Catesby's Car., vol. i. pl. 87. p. 87.—Bonn. et Vieill. Ency. Méth. Orn., part i. p. 48. pl. 16. fig. 2.

———— *Sula*, Linn. Syst. Nat. vol. i. p. 218.

———— *plotus*, Forst. Icon. Ined. 108 ?

*Mār-ga*, Aborigines of Port Essington.

*Booby*, of the Colonists.

---

THE Gannet figured in the accompanying Plate, which may perhaps be identical with the *Pelecanus fiber* of Linnaeus, is abundantly dispersed round the northern shores of the Australian continent ; I have specimens killed within the harbour at Port Essington, and from Raines Island in Torres' Straits, where it breeds in considerable numbers.

The plumage of the two sexes is so precisely similar that it is utterly impossible to distinguish them by external observation ; it is true that the colouring of the feet, face and other soft parts is not always alike, but this difference I believe to be the result of age, rather than of a difference in sex ; and if this opinion be correct, the bright yellow-coloured feet are indicative of the bird being fully adult, and the olive-brown of its being immature.

In its habits, manners, mode of life, and in the nature of its food, this species resembles the other members of the genus.

Head, neck, breast, all the upper surface, wings and tail dark chocolate-brown ; under surface pure white, separated from the brown of the breast by a sharply defined line ; irides very pale yellow ; bill and orbits primrose-yellow, blotched before and beneath the eye with bluish ; eyelash light ash-grey ; legs and feet pale yellow.

The figures represent adult birds rather less than the natural size.



Gould, John. 1848. "Sula fusca, Briss., Brown Gannet, [Pl. 78]." *The birds of Australia* 7, -. <https://doi.org/10.5962/p.313484>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/189241>

**DOI:** <https://doi.org/10.5962/p.313484>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/313484>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Biodiversity Heritage Library

#### **Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.