RAMPHOMICRON STANLEYI.

RAMPHOMICRON STANLEYI.

Stanley's Thorn-bill.

Trochilus Stanleyi, Bourc. et Muls. in Ann. de la Soc. d'Agr., &c. de Lyon, May 24, 1850.

So largely have our collections been enriched of late years with additional species of this lovely group of birds, that I believe I am right in affirming that only fifteen years prior to the present period (1852), not a single species of the restricted genus *Ramphomicron* was known to the Ornithologists of Europe. This may have arisen in a great measure from the scanty knowledge we then had of the productions of the great Andean range, particularly that portion bordering on the Equator, from the circumstance of some of the species being apparently confined to very limited areas, and to the fact of many of them inhabiting localities wherein no one could have imagined that such delicate creatures would be found, and where, consequently, they were never sought for. The species figured on the accompanying Plate is a remarkable illustration of the case in point, the bird being, as I have reason to believe, confined to the inner sides of the great crater of Pichincha in Ecuador, where it was discovered by the French Trochilidist, M. Bourcier, whose account of his ascent of the mountain and discovery of the bird is so admirably detailed, that it is only an act of justice to him to give the account as nearly as may be in his own words. Before doing so, however, I would remark that this bird, which is of no ordinary interest, possesses characters so peculiarly its own, that it cannot for a moment be confounded with any other species.

The dark and lugubrious colouring, relieved alone by the narrow metallic streak down the throat, indicates, on the one hand, the kind of situation in which it would probably be found; and on the other, its short and feeble bill points out that minute insects constitute its food, and that a corresponding diminutive flora doubtless clothes the situations which the bird frequents.

The present makes the fifth species of the well-defined genus *Ramphomicron*, and although not the most highly coloured, it is not the least interesting in the eyes of the true Naturalist. In the blue colouring of its back it is nearly allied to *R. microrhyncha*; while in the form of its beard or throat-mark it reminds one of the well-known *R. heteropogon*. As is the case with those species, the female differs from the male in wanting the brilliant throat-mark; and the young birds are mottled on the throat with white.

M. Bourcier has dedicated this species to the son of the Earl of Derby, as a tribute of respect to his Lordship, and to his noble grandfather, who devoted a long lifetime to the pursuit of Natural History, and contributed so largely and so munificently to the advancement of that branch of science.

The following account by M. Bourcier of his discovery of this beautiful species, as given in a letter to his friend M. Mulsant, will be read with interest:—

“A few days ago, accompanied by two Indians, I attempted an ascent of the mountain Pichincha, a volcano so called, situated about three leagues and a half to the west-north-west of the city of Quito, but which is not usually reached until after a seven or eight hours' march. Part of the journey may be easily performed on horseback, but on attaining a certain elevation you are compelled to quit your horses, and let them await your return: fortunately the uninhabited position permits of your tethering them without fear of robbers. From this point the ascent became very difficult, as we had continually to climb over the heaps of pumice-stones with which the steep sides of the mountain were strewn. But how can I describe to you the magnificent picture which, after having surmounted these obstacles, met our delighted gaze? Picture to yourself two crater-formed cavities, separated by a trachytic wall, from the bottom of which opens nearly forty mouths vomiting smoke. We were at a height above the level of the sea corresponding with the summit of Mont Blanc, without a trace of vegetation around us; it was the desert in its majesty and its silence. The Condor alone, the King of these elevated solitudes, hovered above these desert places, his eternal domain. At our feet were immense gullies, of which the distance disguised their vast depth. They had been formed without doubt by the dreadful eruptions of which history and tradition have preserved the most unhappy remembrances. Unfortunately we had proceeded in the direction of the most abruptly elevated point, and were obliged to make a detour of three quarters of a league to find a more gradual ascent; nevertheless, it was still from 400 to 500 metres. This delayed my arrival at the middle of the desert circle, and prevented me from approaching near to these breathing-holes, which perhaps preserve the country from fresh eruptions of the volcano, and from exploring with ornithological views a country too rarely visited. Then think of the troubles and dangers to be encountered before being repaid! To descend these steep declivities, one is obliged to support oneself on one's hands, and occasionally to slide over the surface. Often the calcined accumulations on which you tread pulverize under your feet, causing those around to lose their equilibrium and draw others after them, the falling masses frequently menacing the life of the traveller. After four hours of fatigue alleviated by the hope of pleasures to come, we arrived at the wished-for destination, and found ourselves in a crater of a circular or rather slightly oval form, surrounded by a kind of wall of trachyte of a uniform elevation, except on the western side, where it is cut down for the passage of the waters, which falling into the Esmeralda, carry their tribute to the Pacific

Ocean. In this sort of enclosure arises an elevated cone, from whence are emitted numerous jets of sulphurous acid, and from whence escapes, accompanied by a slight noise, a watery vapour. In some of these breathing-holes the sulphur is condensed into crystals; and in many places the ground is covered with pulverized pumice or black cinders, which renders one fearful of approaching them. The earth presents crevices the depth of which the eye dares scarcely measure, and some of which must be at least 200 metres deep; they gather the rain and snow-waters, and become the beds of rivulets which flow westward. These ravines or *quebrados*, as the Indians call them, frequently obliged us to make lengthy detours, and to waste a considerable amount of time. The sides of those of moderate depth were carpeted with verdure and ornamented with various shrubs. Here it was that I had the pleasure of discovering the *Trochilus Stanleyi*, a lovely species, which rifles flowers of the *Chuquiraga insignis*, a plant so named by the illustrious Humboldt, of which it appears to be an ardent lover conjointly with *T. pichincha*, with which species it is continually at war.

"I found that it would require a week completely to explore these wilds; but how were we to carry the necessary provisions? and how could I hope for fine weather at such an elevation for an entire week? On the present occasion I had been favoured in this latter respect beyond my hopes; to a magnificent day succeeded a beautiful night, during which the moon, at its full, lent a fairy charm to the scene. Wrapped in our ponchos, a kind of cloak of the country, we awaited the return of day; reclining upon stones and sheltered by a rock, in spite of the inconvenience of the position, we enjoyed the pleasures of a deep sleep upon this menacing soil, which in a moment of caprice might launch us into eternity. The next day unexpected troubles attended us; in order to regain the rocks which formed the barrier to the enclosure, we were obliged to pick our way along a sandy soil, in which we sank up to our knees; and in order to creep along this quicksand, which continually yielded to our feet, we had to support ourselves by rocks, which as frequently broke away beneath our fingers. Arriving at last, after many long and weary hours, on the highest elevation of this impracticable ground, we found ourselves in the Paramos mountains, used for the pasturage of cattle. Here the winds blew at once with such violence, and a blast so keen, that I twice essayed to continue my route before I could muster courage so to do. We now took again to our horses, with which we soon gained the woody slopes, clothed with various species of *Datura*, and the rich plains which lead to the city. On re-entering Quito, all my troubles were forgotten, while the pleasure of having made a new discovery remained impressed upon my memory, coupled with a vivid recollection of the interesting scene in which it had occurred."

Head, sides, and back of the neck and wing-coverts greenish bronze; back and rump deep violet blue; wings purplish brown; upper tail-coverts and tail dark bluish green; down the centre of the throat is a series of scale-like feathers, broad at the chin and gradually tapering to a point on the breast, the upper part of which is of a brilliant metallic emerald green, which passes into the amethystine blue, tinged on the margins with red of the lower portion; under surface dark sooty-brown washed on the flanks with bronze; under tail-coverts of a greyish white, with a streak of steel-blue down the centre of each feather; bill and feet black.

The female differs in having no trace of the brilliant gorget.

The Plate represents two males and a female of the natural size on the *Siphocampylus giganteus*.

Gould, John. 1857. "Ramphomicron stanleyi, Stanley's Thorn-bill. [Pl. 185]." *A monograph of the Trochilidæ, or family of humming-birds* 3,
<https://doi.org/10.5962/p.316994>.

View This Item Online: <https://www.biodiversitylibrary.org/item/109468>

DOI: <https://doi.org/10.5962/p.316994>

Permalink: <https://www.biodiversitylibrary.org/partpdf/316994>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.