PHAOLAIMA ÆQUATORIALIS, *Coult.*

PHAIOLAIMA ÆQUATORIALIS, *Gould.*

Equatorial Lilac-throat.

Pheolema rubinoides, Sclat. in Proc. Zool. Soc., part xxvii. p. 145.

——— *æquatorialis*, Gould, Sclat. in Proc. Zool. Soc., part xxviii. p. 70.

I HAVE for many years been aware of the existence of a second species of *Phaiolaima*, M. Bourcier having pointed it out soon after his return from Ecuador, and assigned to me the privilege of giving it a name, which I have neglected to do until the present year, 1860. Besides the examples procured in Ecuador which grace M. Bourcier's collection, I have received others from Quito, and I have three more which were collected at Pallatanga by Mr. Fraser, from whose notes we learn that the bill is black, the irides hazel, and the legs and feet dark flesh-colour. Ecuador, then, is evidently the true habitat of this fine species—a species which is nearly allied to, but quite distinct from, the *Phaiolaima rubinoides* of Santa Fé de Bogota. It is not only distinguished from its northern ally by its greater size, but also by the total absence of any glittering feathers on the crown, which are always present in *P. rubinoides*; the luminous lilac patch on the throat, too, is not so extensive; neither is it so brilliant. A peculiar style of colouring pervades both species, by which they are distinguished from all other Humming-Birds,—a character which cannot be rendered very clear by description, but which is readily perceptible on reference to the plates in which they are depicted; they have indeed a sombre-brown look quite peculiar to themselves.

The sexes probably do not differ in colouring. It is true that I have seen examples without the luminous throat-mark, which might be supposed to be females; but as they offer no other difference in their colouring or markings, I am induced to believe them to be young birds.

Crown of the head, face, sides of the neck, and chin pure green; on the centre of the throat a small patch of pale metallic lilac; all the upper surface bronzy green, the bronzy hue predominating on the wing- and tail-coverts; wings purplish brown, edged on the shoulders and outer primary with deep buff; tail pure bronze with deep buff shafts; under surface buff, washed with green on the flanks.

The figures are of the natural size. The Plant is *Thibaudia Pichinchensis*.

Gould, John. 1859. "Phæolæma æquatorialis, Gould., Equatorial Lilac-throat. [Pl. 269]." *A monograph of the Trochilidæ, or family of humming-birds* 4, <https://doi.org/10.5962/p.317078>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108334>

DOI: <https://doi.org/10.5962/p.317078>

Permalink: <https://www.biodiversitylibrary.org/partpdf/317078>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.