THAUMATIAS MILLERI.

THAUMATIAS MILLERI.

Miller's Emerald.

Trochilus Milleri, Lodd. MSS. ined. Bourc. in Proc. Zool. Soc., pt. xv. p. 43.

Polytmus Milleri, Gray and Mitch. Gen. of Birds, vol. i. p. 108, *Polytmus*, sp. 46.

Thaumatias milleri, Bonap. Consp. Gen. Av., p. 78, *Thaumatias*, sp. 4.

Aggyrtia Milleri, Reich. Auf. der Col., p. 10.

Thaumantias milleri, Bonap. Rev. Zool. 1854, p. 255.

MILLER'S Emerald was first described by M. Bourcier of Paris in the "Proceedings of the Zoological Society of London," the specific name of *Milleri* being adopted by him from the late Mr. George Loddiges' MSS. It is not known why Mr. Loddiges gave the bird this specific appellation, unless it was from a desire to perpetuate the name of a surgeon in the Royal Navy, who paid considerable attention to the Humming-birds while stationed on the Pacific side of South America.

The *Thaumatias Milleri* is a diminutive and chastely-coloured species, distinguished by its snow-white breast and glittering crown; in size it is very similar to *T. brevirostris*, with which it might be confounded had it a less brilliant forehead. The native habitat of the species is the Andes of Columbia and the countries lying to the westward towards the upper part of the Rivers Negro and Amazon: most of the specimens which have reached this country are from Bogota. It is by no means a common bird, and but few collectors are in possession of examples; we are, indeed, so little acquainted with it that it would be unsafe to say if the female differs in plumage from her mate; in all probability she is very similarly coloured, but her hues are less bright and contrasted in all their tints, particularly in the glittering portion of the crown and occiput.

The crown, sides of the head, and the sides and back of the neck rich glittering grass-green; upper surface and wing-coverts bronzy green; wings purplish black; tail-feathers greyish green with a transversal mark of brown near the tip of all but the two central ones; throat snow-white; under surface of the body greyish white washed with green on the flanks, the green meeting and forming a band across the breast; upper mandible and tip of the lower mandible black; the basal portion fleshy.

The Plate represents the birds of the natural size. The plant is the *Comparettia falcata*.

Gould, John. 1860. "Thaumatias milleri, Miller's Emerald. [Pl. 296]." *A monograph of the Trochilidæ, or family of humming-birds* 5,
<https://doi.org/10.5962/p.317105>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108806>

DOI: <https://doi.org/10.5962/p.317105>

Permalink: <https://www.biodiversitylibrary.org/partpdf/317105>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.