AMAZILIA VIRIDIGASTER.

AMAZILIA VIRIDIGASTER.

Green-bellied Amazili.

Trochilus viridigaster, Bourc. Rev. Zool. 1843, p. 105.—Ib. Ann. Sci. Phys. de Lyon, 1843, p. 42.

Hylocharis viridigaster, Bonap. Consp. Gen. Av., p. 74, *Hylocharis*, sp. 5.—Gray and Mitch. Gen. of Birds, vol. i. p. 115, *Hylocharis*, sp. 41.

Saucerottia viridiventris, Reichenb. Auf. der Col., p. 8.

——— *viridigaster*, Bonap. Rev. et Mag. de Zool. 1854, p. 255.—Sclat. in Proc. of Zool. Soc., part xxv. p. 16.

It will be admitted by every ornithologist that a more appropriate specific appellation than the one applied to this species might easily have been selected—one having reference, for instance, to its greyish-brown back or its nearly pure violet tail, whereas the term *viridigaster* will apply equally well to at least a hundred species. So far, however, as I am aware, no other term has ever been applied to it, except *viridiventris*, which has the same meaning; *viridigaster* must therefore be retained.

The native country of this bird is New Grenada, and numerous are the examples which have from time to time been sent to Europe from Bogota. M. Bourcier once told me that, upon reconsidering the subject, he feared that the bird to which he had given the name of *Trochilus Mariæ* might prove to be identical with the present species; but on carefully comparing his description with the examples from which the figures in the accompanying Plate were taken, I find it to differ so much, especially in the colouring of the wings, as to preclude the possibility of such being the case; in all probability it is identical with *A. Devillei*.

I regret that I have not a single note to add respecting the habits and economy of this bird. I believe that all the specimens which have reached Europe have been killed and skinned by the Indians, who, as is well known, pay little or no attention to habits: all they desire is to approach sufficiently near to obtain them by means of the blow-pipe, with which they take almost unerring aim, using round balls of clay for their pellets.

Like the other *Amazilia*, the sexes resemble each other in colour; if any difference occurs, it consists in the somewhat brighter hues of the male.

Head, upper part of the back, and wing-coverts dull green; lower part of the back and upper tail-coverts greyish brown; wings purplish brown; tail deep violet; under surface shining grass-green, passing into greyish brown; under tail-coverts greyish buff, with lighter edges; thighs somewhat thickly clothed and white; bill black, except the basal two-thirds of the lower mandible, which appears to have been fleshy.

The figures are of the size of nature. The plant is the *Lælia acuminata*.

Gould, John. 1861. "Amazilia viridigaster, Green-bellied Amazili. [Pl. 314]." *A monograph of the Trochilidæ, or family of humming-birds* 5,
<https://doi.org/10.5962/p.317123>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108806>

DOI: <https://doi.org/10.5962/p.317123>

Permalink: <https://www.biodiversitylibrary.org/partpdf/317123>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.