SAPPHIRONIA CERULEOGULARIS, Gould

J. Gould and H. C. Richter, del. et lith.

Hellmuth & Walton, Imp.

SAPPHIRONIA CÆRULEOGULARIS, Gould.

Blue-throated Sapphironia.

Trochilus (——?) *cæruleogularis*, Gould in Proc. of Zool. Soc., part xviii. p. 163.

——— *Duchassaigui*, Bourc. Compt. Rend. de l'Acad. des Sci., tom. xxxii. p. 187.

Thalurania Caelina, Boire. Rev. et Mag. de Zool. 1856, p. 552.

Cyanochloris cæruleigularis, Reichenb. Auf. der Col., p. 10.

Sapphironia cæruleigularis, Bonap. Rev. et Mag. de Zool. 1854, p. 256.

Hylocharis (?) *cæruleigularis*, Sclat. in Proc. of Zool. Soc., part xxiv. p. 140.

I HAVE not placed the *Trochilus Duchassaigui* and *Thalurania Caelina* of M. Bourcier among the synonyms of the present species without due consideration and a careful comparison of the typical specimens one with the other. The *T. Duchassaigui* is a young male of the bird here represented which had not fully attained the blue colouring of the throat. The specific name *Caelina* not having been given till a year later than my own of *cæruleogularis*, it must give place to that term; besides which, the bird is not a *Thalurania*.

In form, this species and *Sapphironia Goudoti* are precisely alike, and these birds form a minute but well-marked section of the *Trochilidæ*, the species of which are characterized by great elegance of form, as well as by the beauty of their colouring: the under tail-coverts of both species are very conspicuous and extremely pretty.

M. Verreaux has received this bird in tolerable abundance from Santa Martha; my own were obtained in Costa Rica, and M. Bourcier's from Panama; it is clear, therefore, that New Grenada, Panama, and Costa Rica are the countries in which this bird flies. Mr. Bridges informs us that it is found in the very streets of the town of David, feeding on the *Tamarindus Indicus* and orange-trees, and that, like all other Humming-Birds, it is very pugnacious, and constantly fighting with others of its species.

A marked difference is observable in the colouring of the sexes,—the female, although having the same lengthened form as the male, being devoid of the bright blue of the throat and the fine green of the abdomen—those parts being greyish white; these, as well as other differences, will be readily perceived on an inspection of the accompanying figures, or a perusal of the following descriptions:—

The male has the upper surface, shoulders, abdomen, and under tail-coverts grass-green; throat, sides of the neck and chest rich violet-blue; wings purplish brown; central tail-feathers bronzy green; lateral feathers purplish black; upper mandible and tip of the lower black, the basal portion of the latter ashy white; the under tail-coverts, which are largely developed, green, margined with silvery grey.

The female has the upper surface bronzy green; tail purplish black, the two central feathers glossed with bronze, the four lateral feathers on each side tipped with white; centre of the throat, abdomen, and under tail-coverts white.

The figures are those of the male and female of the size of life. The plant is the *Cuphea silenoïdes*.

Gould, John. 1861. "Sapphironia cæruleogularis, Gould., Blue-throated Sapphironia. [Pl. 346]." *A monograph of the Trochilidæ, or family of humming-birds* 5, <https://doi.org/10.5962/p.317155>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108806>

DOI: <https://doi.org/10.5962/p.317155>

Permalink: <https://www.biodiversitylibrary.org/partpdf/317155>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.