CHLOROSTILBON PHÆTON.

CHLOROSTILBON PHAËTHON.

Glittering Emerald.

Trochilus phaëthon, Bourc. Rev. Zool. 1848, p. 274.

——— *phaeton*, Gray and Mitch. Gen. of Birds, vol. iii. App. p. 300, App. to p. 103.

Chlorestes Phaëthon, Reich. Auf. der Col., p. 7.—Ib. Troch. Enum., p. 4.

Hylocharis phaëton, Bonap. Rev. et Mag. de Zool. 1854, p. 255.

——— *aureiventris*, Bonap. Rev. et Mag. de Zool. 1854, p. 255.

——— *similis*, Bonap. Consp. Gen. Av., tom. i. p. 74, *Hylocharis*, sp. 4?

Chlorolampis phaëthon, Cab. Mus. Hein., Theil iii. p. 48, note.

Trochilus flavifrons, Gould, Zool. of Beagle, part iii. Birds, p. 110.

——— *metallicus*, Gould, MS.

Ornismya aureiventris, D'Orb. and La Fresn. Syn. Av., p. 28.

Trochilis similis, Bourc. ?

IF we consider the metallic-plumaged Humming-Birds of Bolivia, Peru, and La Plata to be one and the same species, then this bird enjoys a range extending across the continent of South America from east to west. It is true that the specimens from the three localities above mentioned have received distinct specific appellations, but, except in size, they offer no marked difference. The bird frequenting the high lands of Bolivia was named *aureiventris* by MM. D'Orbigny and La Fresnaye; the Peruvian bird so common at Chuquesaca had the name *Phaëthon* assigned to it by M. Bourcier, and the one from La Plata that of *flavifrons*, by myself.

The typical specimen from which M. Bourcier took his description of *Phaëthon* is now before me; it is evidently the Chuquesacan bird, and is by far the largest in size when compared with the examples from the other localities mentioned; those from Monte Video, where specimens were procured by Mr. Darwin, are the next in size, and the *aureiventris* is the least; all are precisely alike in form and colour, their only difference being, as already mentioned, in their admeasurements.

My figures, which represent both sexes of the size of life, were taken from Chuquesacan birds. Time, and the acquisition of a greater number of specimens from every locality, can alone determine whether we are warranted in considering these birds as one or three species: what is the *Trochilis similis* of M. Bourcier?

The male has the head, all the upper surface and wing-coverts rich golden bronze, but inclining to green on the upper tail-coverts; wings purplish brown; tail black, glossed with deep green; throat and breast glittering emerald-green, merging into the glittering coppery bronze of the sides of the neck and abdomen; under tail-coverts green; bill apparently fleshy red at the base, with a darker tip.

The female is bronzy green above and grey beneath, washed with bronze on the flanks; wings purplish brown; tail bluish black, the two lateral feathers tipped with greyish white.

The Plate represents the two sexes of the natural size. The plant is the *Calliandra Tweediei*.

Gould, John. 1861. "Chlorostilbon phaëthon, Glittering Emerald. [Pl. 354]." *A monograph of the Trochilidæ, or family of humming-birds* 5,
<https://doi.org/10.5962/p.317163>.

View This Item Online: <https://www.biodiversitylibrary.org/item/108806>

DOI: <https://doi.org/10.5962/p.317163>

Permalink: <https://www.biodiversitylibrary.org/partpdf/317163>

Holding Institution

Smithsonian Libraries

Sponsored by

Smithsonian Institution Libraries

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.